

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

## **1. NAME OF THE MEDICINAL PRODUCT**

Spinraza 12 mg solution for injection

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each 5 ml vial contains nusinersen sodium equivalent to 12 mg nusinersen.  
Each ml contains 2.4 mg of nusinersen.

For the full list of excipients, see section 6.1.

## **3. PHARMACEUTICAL FORM**

Solution for injection.

Clear and colourless solution with pH of approximately 7.2.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Spinraza is indicated for the treatment of 5q Spinal Muscular Atrophy.

### **4.2 Posology and method of administration**

Treatment with Spinraza should only be initiated by a physician with experience in the management of spinal muscular atrophy (SMA).

The decision to treat should be based on an individualised expert evaluation of the expected benefits of treatment for that individual, balanced against the potential risk of treatment with Spinraza. Patients with profound hypotonia and respiratory failure at birth, where Spinraza has not been studied, may not experience a clinically meaningful benefit due to severe SMN protein deficiency.

#### Posology

Spinraza is for intrathecal use by lumbar puncture.

The recommended dosage is 12 mg (5 ml) per administration.

Spinraza treatment should be initiated as early as possible after diagnosis with 4 loading doses on Days 0, 14, 28 and 63. A maintenance dose should be administered once every 4 months thereafter.

#### *Duration of treatment*

Information on long term efficacy of this medicinal product is not available. The need for continuation of therapy should be reviewed regularly and considered on an individual basis depending on the patient's clinical presentation and response to the therapy.

#### *Missed or delayed doses*

If a loading dose is delayed or missed Spinraza should be administered as soon as possible, with at least 14 days between doses, and continue dosing at the prescribed frequency. If a maintenance dose is

delayed or missed, Spinraza should be administered as soon as possible and dosing continued every 4 months.

### Special populations

#### *Renal impairment*

Spinraza has not been studied in patients with renal impairment. The safety and efficacy in patients with renal impairment has not been established and they should be closely observed.

#### *Hepatic impairment*

Spinraza has not been studied in patients with hepatic impairment. Spinraza is not metabolised via the cytochrome P450 enzyme system in the liver, therefore dose adjustment is unlikely to be required in patients with hepatic impairment (see sections 4.5 and 5.2).

### Method of administration

Treatment should be administered by health care professionals experienced in performing lumbar punctures.

Spinraza is administered as an intrathecal bolus injection over 1 to 3 minutes, using a spinal anaesthesia needle. The injection must not be administered in areas of the skin where there are signs of infection or inflammation. It is recommended that the volume of cerebral spinal fluid (CSF), equivalent to the volume of Spinraza to be injected, is removed prior to administration of Spinraza.

Sedation may be required to administer Spinraza, as indicated by the clinical condition of the patient. Ultrasound (or other imaging techniques) may be considered to guide intrathecal administration of Spinraza, particularly in younger patients and in patients with scoliosis. Aseptic technique should be used when preparing and administering Spinraza; see instructions for use in section 6.6.

### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

### **4.4 Special warnings and precautions for use**

#### Lumbar puncture procedure

There is a risk of adverse reactions occurring as part of the lumbar puncture procedure (e.g. headache, back pain, vomiting; see section 4.8). Potential difficulties with this route of administration may be seen in very young patients and those with scoliosis. The use of ultrasound or other imaging techniques to assist with intrathecal administration of Spinraza, can be considered at the physician's discretion.

#### Thrombocytopenia and coagulation abnormalities

Thrombocytopenia and coagulation abnormalities, including acute severe thrombocytopenia, have been observed after administration of other subcutaneously or intravenously administered antisense oligonucleotides. If clinically indicated, platelet and coagulation laboratory testing is recommended prior to administration of Spinraza.

#### Renal toxicity

Renal toxicity has been observed after administration of other subcutaneously and intravenously administered antisense oligonucleotides. If clinically indicated, urine protein testing (preferably using a first morning urine specimen) is recommended. For persistent elevated urinary protein, further evaluation should be considered.

## Hydrocephalus

There have been reports of communicating hydrocephalus not related to meningitis or bleeding in patients treated with nusinersen in the post-marketing setting. Some patients were implanted with a ventriculo-peritoneal shunt. In patients with decreased consciousness, an evaluation for hydrocephalus should be considered. The benefits-and risks of nusinersen treatment in patients with a ventriculo-peritoneal shunt are unknown at present and the maintenance of treatment needs to be carefully considered.

### **4.5 Interaction with other medicinal products and other forms of interaction**

No interaction studies have been performed. *In vitro* studies indicated that nusinersen is not an inducer or inhibitor of CYP450 mediated metabolism. *In vitro* studies indicate that the likelihood for interactions with nusinersen due to competition for plasma protein binding, or competition with or inhibition of transporters is low.

### **4.6 Fertility, pregnancy and lactation**

#### Pregnancy

There are no or limited amount of data from the use of nusinersen in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3). As a precautionary measure, it is preferable to avoid the use of Spinraza during pregnancy.

#### Breast-feeding

It is unknown whether nusinersen/metabolites are excreted in human milk.

A risk to the newborn/infants cannot be excluded. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from Spinraza therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

#### Fertility

In toxicity studies in animals no effects on male or female fertility were observed (see section 5.3). There are no data available on the potential effects on fertility in humans.

### **4.7 Effects on ability to drive and use machines**

Spinraza has no or negligible influence on the ability to drive and use machines.

### **4.8 Undesirable effects**

#### Summary of safety profile

The safety assessment of Spinraza was based on two Phase 3 clinical studies in infants (CS3B) and children (CS4) with SMA, together with one Phase 2 study in infants and children with SMA (CS7) and open-label studies including pre-symptomatic infants (CS5) genetically diagnosed with SMA and infants and children with SMA. Study CS11 enrolled infantile and later-onset patients including those who had completed studies CS3B, CS4 and CS12. Of the 346 patients who received Spinraza up to a maximum of 5 years, 258 patients received treatment for at least 1 year.

#### Tabulated list of adverse reactions

The assessment of undesirable effects is based on the following frequency data:  
Very common ( $\geq 1/10$ )

Not known (cannot be estimated from the available data)

Table 1: Adverse reactions related to lumbar puncture procedure reported in CS4 (later onset SMA) with an incidence at least 5% higher in patients treated with Spinraza than sham-control

MedDRA System Organ Class	MedDRA preferred term	Spinraza frequency category, n=84
Nervous system disorders	Headache*	Very common
Gastrointestinal disorders	Vomiting*	Very common
Musculoskeletal and connective tissue disorders	Back pain*	Very common

\*Adverse events considered related to the lumbar puncture procedure. These events can be considered manifestations of post-lumbar puncture syndrome.

### Post-marketing experience

Adverse reactions have been identified during post-approval use of Spinraza. Among patients treated with Spinraza by lumbar puncture, serious infection, such as meningitis, has been observed. Communicating hydrocephalus, aseptic meningitis and hypersensitivity (e.g. angioedema, urticaria and rash) have also been reported. The frequency of these reactions is not known as they have been reported from the post marketing setting.

### Description of selected adverse reactions

Adverse reactions associated with the administration of Spinraza by lumbar puncture have been observed. The majority of these are reported within 72 hours of the procedure. The incidence and severity of these events were consistent with events expected to occur with lumbar puncture. No serious complications of lumbar puncture, such as serious infections, have been observed in the clinical trials of Spinraza.

Some adverse events commonly associated with lumbar puncture (e.g. headache and back pain) could not be assessed in the infant population exposed to Spinraza due to the limited communication appropriate for that age group.

### Immunogenicity

The immunogenic response to nusinersen was determined in 346 patients with baseline and post-baseline plasma samples evaluated for anti-drug antibodies (ADA). Overall, the incidence of ADAs was low, with 15 (4%) patients classified as ADA positive overall, of which 4 had a transient response, 5 had a persistent response, and 6 patients had responses which could not be classified as transient or persistent at the time of data cut off. The impact of immunogenicity on safety was not formally analysed as the number of patients with ADAs was low. However, individual safety data for the treatment-emergent ADA-positive cases were reviewed, and no AEs of interest were identified.

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions **via the national reporting system listed in [Appendix V](#)**.

## 4.9 Overdose

No cases of overdose associated with adverse reactions were reported in clinical studies.

In the event of an overdose, supportive medical care should be provided including consulting with a healthcare professional and close observation of the clinical status of the patient.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Other drugs for disorders of the musculo-skeletal system, ATC code: M09AX07

#### Mechanism of action

Nusinersen is an antisense oligonucleotide (ASO) which increases the proportion of exon 7 inclusion in survival motor neuron 2 (SMN2) messenger ribonucleic acid (mRNA) transcripts by binding to an intronic splice silencing site (ISS-N1) found in intron 7 of the SMN2 pre-messenger ribonucleic acid (pre-mRNA). By binding, the ASO displaces splicing factors, which normally suppress splicing. Displacement of these factors leads to retention of exon 7 in the SMN2 mRNA and hence when SMN2 mRNA is produced, it can be translated into the functional full length SMN protein.

SMA is a progressive neuromuscular disease resulting from mutations in chromosome 5q in the SMN1 gene. A second gene SMN2, located near SMN1, is responsible for a small amount of SMN protein production. SMA is a clinical spectrum of disease with disease severity linked to fewer numbers of SMN2 gene copies and younger age of symptom onset.

#### Clinical efficacy and safety

##### Symptomatic patients

###### *Infantile onset*

Study CS3B (ENDEAR) was a Phase 3, randomized, double-blind, sham-procedure controlled study conducted in 121 symptomatic infants  $\leq 7$  months of age, diagnosed with SMA (symptom onset before 6 months of age). CS3B was designed to assess the effect of Spinraza on motor function and survival. Patients were randomized 2:1 to either Spinraza (as per the approved dosing regimen) or sham-control, with a length of treatment ranging from 6 to 442 days.

The median age of onset of clinical signs and symptoms of SMA was 6.5 weeks and 8 weeks for Spinraza treated versus sham-control patients respectively, with 99% of patients having 2 copies of the SMN2 gene and therefore deemed most likely to develop Type I SMA. The median age when patients received their first dose was 164.5 days for treated patients, and 205 days for sham-control. Baseline disease characteristics were largely similar in the Spinraza treated patients and sham-control patients except that Spinraza treated patients at baseline had a higher percentage compared to sham-control patients of paradoxical breathing (89% vs 66%), pneumonia or respiratory symptoms (35% vs 22%), swallowing or feeding difficulties (51% vs 29%) and requirement for respiratory support (26% vs 15%).

At the final analysis, a statistically significant greater percentage of patients achieved the definition of a motor milestone responder in the Spinraza group (51%) compared to the sham-control group (0%) ( $p < 0.0001$ ). Time to death or permanent ventilation ( $\geq 16$  hours ventilation/day continuously for  $> 21$  days in the absence of an acute reversible event or tracheostomy) was assessed as the primary endpoint. Statistically significant effects on event-free survival, overall survival, the proportion of patients achieving the definition of a motor milestone responder, and the percentage of patients with at least a 4-point improvement from baseline in Children's Hospital of Philadelphia

Infant Test for Neuromuscular Disease (CHOP INTEND) score were observed in patients in the Spinraza group compared to those in the sham-control group (Table 2).

In the efficacy set, 18 patients (25%) in the Spinraza group and 12 patients (32%) in the sham-control group required permanent ventilation. Of these patients, 6 (33%) in the Spinraza group and 0 (0%) in the sham-control group met the protocol-defined criteria for a motor-milestone responder.

**Table 2: Primary and secondary endpoints at final analysis – Study CS3B**

<b>Efficacy Parameter</b>	<b>Spinraza treated Patients</b>	<b>Sham-control Patients</b>
<b>Survival</b>		
<b>Event-free survival<sup>2</sup></b> Number of patients who died or received permanent ventilation	31 (39%)	28 (68%)
Hazard ratio (95% CI) p-value <sup>6</sup>	0.53 (0.32 -0.89) p = 0.0046	
<b>Overall survival<sup>2</sup></b> Number of patients who died	13 (16%)	16 (39%)
Hazard Ratio (95% CI) p-value <sup>6</sup>	0.37 (0.18 – 0.77) p=0.0041	
<b>Motor function</b>		
<b>Motor milestones<sup>3</sup></b> Proportion achieving pre-defined motor milestone responder criteria (HINE section 2) <sup>4,5</sup>	37 (51%) <sup>1</sup> p<0.0001	0 (0%)
Proportion at Day 183	41%	5%
Proportion at Day 302	45%	0%
Proportion at Day 394	54%	0%
Proportion with improvement in total motor milestone score	49 (67%)	5 (14%)
Proportion with worsening in total motor milestone score	1 (1%)	8 (22%)
<b>CHOP INTEND<sup>3</sup></b> Proportion achieving a 4-point improvement	52 (71%) p<0.0001	1 (3%)
Proportion achieving a 4-point worsening	2 (3%)	17 (46%)
Proportion with any improvement	53 (73%)	1 (3%)
Proportion with any worsening	5 (7%)	18 (49%)

<sup>1</sup>CS3B was stopped following positive statistical analysis on the primary endpoint at interim analysis (statistically significantly greater percentage of patients achieved the definition of a motor milestone responder in the Spinraza group (41%) compared to the sham-control group (0%), p<0.0001)

<sup>2</sup>At the final analysis, event-free survival and overall survival were assessed using the Intent to Treat population (ITT Spinraza n=80; Sham-control n=41).

<sup>3</sup>At the final analysis, CHOP INTEND and motor milestone analyses were conducted using the Efficacy Set (Spinraza n=73; Sham-control n=37).


<sup>4</sup>Assessed at the later of Day 183, Day 302, and Day 394 Study Visit

<sup>5</sup>According to Hammersmith Infant Neurological Examination (HINE) section 2: ≥2 point increase [or maximal score] in ability to kick, OR ≥1 point increase in the motor milestones of head control, rolling, sitting, crawling, standing or walking, AND improvement in more categories of motor milestones than worsening, defined as a responder for this primary analysis.

<sup>6</sup>Based on log-rank test stratified by disease duration

The extent of improvement in CHOP INTEND is shown in Figure 1 (change from baseline score for each subject).

Figure 1: Change in CHOP INTEND from Baseline to Later of Day 183, Day 302, and Day 394 Study Visit – Endear Study /CS3B (Efficacy Set, ES)



Note 1: Shortest bars at 0 line indicate 0 value.

Note 2: Out of the 110 patients in the efficacy set, 29 died (13 (18%) for Spinraza and 16 (43%) for Control) and 3 withdrew for reason other than death (2 (3%) for Spinraza and 1 (3%) for Control) and were therefore not included in this analysis of the ES.

To allow for long term follow up of these patients, at the end of Study CS3B, 89 patients (Spinraza: n=65; sham-control: n=24) enrolled in Study CS11 (SHINE). Study CS11 is an open label extension study for SMA patients who previously participated in the other Spinraza clinical studies. In Study CS11 all patients received Spinraza, with the length of treatment ranging from 65 to 592 days (median 289 days) at the time of interim analysis. Improvements in motor function were observed among patients continuing Spinraza from Study CS3B, as well as those who initiated Spinraza in Study CS11 (Figure 3), with the greatest benefit observed in those with earlier treatment initiation. Among patients without permanent ventilation at the baseline of Study CS11, a majority were alive and without permanent ventilation at the time of interim analysis.

In patients randomized to Spinraza in Study CS3B and including the experience in Study CS11, the median time to death or permanent ventilation was 73 weeks. At the time of a Study CS11 interim analysis, 61 out of 65 patients (94%) were alive. Of the 45 patients who had not met the definition of permanent ventilation in Study CS3B, 38 patients (84%) were alive without permanent ventilation in Study CS11 at the time of interim analysis. Further improvement in mean total motor milestone (HINE-Section 2) (2.1; SD 4.36; n=22) and CHOP INTEND (4.68; SD 3.993, n=22) scores were observed from baseline to Study Day 304 in Study CS11.

Patients who first initiated Spinraza treatment in Study CS11 (n=24; sham control in Study CS3B) were of a median age of 17.8 months (range 10 - 23 months) and had a mean CHOP INTEND score of 17.25 (range 2.0 - 46.0) at baseline in Study CS11. At the time of interim analysis, 22 out of 24 patients (92%) were alive. Of the twelve patients (50%) who had not met the definition of permanent ventilation in Study CS3B, 7 patients (58%) were alive without permanent ventilation in Study CS11. The median time to death or permanent ventilation was 50.9 weeks after initiation of Spinraza treatment in Study CS11. Improvement in mean total motor milestone (HINE-Section 2) (1.2; SD 1.8; n=12) and CHOP INTEND (3.58; SD 7.051, n=12) scores were observed from baseline to Study Day 304 in Study CS11.

These results are supported by an open-label Phase 2 study in symptomatic patients diagnosed with SMA (CS3A). Median age of onset of clinical signs and symptoms was 56 days and patients had either 2 SMN2 gene copies (n=17) or 3 SMN2 gene copies (n=2) (SMN2 gene copy number unknown for 1 patient). Patients in this study were deemed most likely to develop Type I SMA. The median age at first dose was 162 days.



The primary endpoint was the proportion of patients who improved in one or more categories in motor milestones (according to HINE section 2:  $\geq 2$  point increase [or maximal score] in ability to kick or voluntary grasp OR  $\geq 1$  point increase in the motor milestones of head control, rolling, sitting, crawling, standing or walking). Twelve out of 20 patients (60%) in the study met the primary endpoint with improvement in mean motor milestone achievement over time. An improvement in mean CHOP INTEND score over time was observed from baseline to day 1072 (mean change 21.30). Overall, 11 out of 20 patients (55%) met the endpoint of an increase in total CHOP INTEND score of  $\geq 4$  points as of the last study visit. Of the 20 subjects enrolled, 11 (55%) were alive and free of permanent ventilation at the last visit. Four patients met the criteria for permanent ventilation and five patients died during the study.

#### *Later onset*

Study CS4 (CHERISH) was a Phase 3, randomised, double-blind, sham-procedure controlled study conducted in 126 symptomatic patients with later-onset SMA (symptom onset after 6 months of age). Patients were randomized 2:1 to either Spinraza (dosed with 3 loading doses and maintenance doses every 6 months) or sham-control, with a length of treatment ranging from 324 to 482 days. The median age at screening was 3 years, and the median age of onset of clinical signs and symptoms of SMA was 11 months. The majority of patients (88%) have 3 copies of the SMN2 gene (8% have 2 copies, 2% have 4 copies, and 2% have an unknown copy number). At baseline, patients had a mean Hammersmith Functional Motor Scale Expanded (HFMSE) score of 21.6, a mean revised upper limb module (RULM) of 19.1, all had achieved independent sitting, and no patients had achieved independent walking. Patients in this study were deemed most likely to develop Type II or III SMA. Baseline disease characteristics were generally similar with the exception of an imbalance in the proportion of patients who had ever achieved the ability to stand without support (13% of patients in the Spinraza group and 29% in sham-control) or walk with support (24% of patients in the Spinraza group and 33% in sham-control).

At the final analysis, a statistically significant improvement in HFMSE score from baseline to Month 15 was seen in the Spinraza group compared to the sham-control group (Table 3, Figure 2). The analysis was conducted in the ITT population (Spinraza: n=84; sham-control: n=42), and post-baseline HFMSE data for patients without a Month 15 visit were imputed using the multiple imputation method. An analysis of the subset of patients in the ITT population who had observed values at Month 15 demonstrated consistent, statistically significant results. Of those with observed values at Month 15, a higher proportion of Spinraza treated subjects had improvement (73% vs 41%, respectively) and a lower proportion of Spinraza treated subjects had worsening (23% vs 44%, respectively) in total HFMSE score compared to sham-control. Secondary endpoints including functional measures and WHO motor milestone achievement were formally statistically tested and are described in Table 3.

Initiation of treatment sooner after symptom onset resulted in earlier and greater improvement in motor function than those with delayed treatment initiation; however, both groups experienced benefit compared to sham-control.

**Table 3: Primary and secondary endpoints at final analysis – Study CS4<sup>1</sup>**

	<b>Spinraza treated Patients</b>	<b>Sham-control Patients</b>
<b>HFMSE score</b> Change from baseline in total HFMSE score at 15 months <sup>1,2,3</sup>	3.9 (95% CI: 3.0, 4.9) p=0.0000001	-1.0 (95% CI: -2.5, 0.5)
Proportion of patients who achieved at least a 3 point improvement from baseline to month 15 <sup>2</sup>	56.8% (95% CI:45.6, 68.1) P=0.0006 <sup>5</sup>	26.3% (95% CI: 12.4,40.2)
<b>RULM</b> Mean change from baseline to month 15 in total RULM score <sup>2,3</sup>	4.2(95% CI: 3.4, 5.0) p=0.0000001 <sup>6</sup>	0.5 (95% CI: -0.6, 1.6)
<b>WHO motor milestones</b> Proportion of patients who achieved new motor milestones at 15 months <sup>4</sup>	19.7% (95% CI: 10.9, 31.3) p=0.0811	5.9% (95% CI: 0.7, 19.7)

<sup>1</sup>CS4 was stopped following positive statistical analysis on the primary endpoint at interim analysis (statistically significant improvement from baseline HFMSE score was observed in Spinraza treated patients compared to the sham-control patients (Spinraza vs. sham-control: 4.0 vs. -1.9; p=0.0000002))

<sup>2</sup> Assessed using the Intent to Treat population (Spinraza n=84; Sham-control n=42); data for patients without a Month 15 visit were imputed using the multiple imputation method


<sup>3</sup>Least squares mean

<sup>4</sup> Assessed using the Month 15 Efficacy Set (Spinraza n=66; Sham control n=34); analyses are based on imputed data when there are missing data.

<sup>5</sup> Based on logistic regression with treatment effect and adjustment for each subject's age at screening and HFMSE score at baseline

<sup>6</sup>Nominal p value

**Figure 2: Mean change from baseline in HFMSE score over time at final analysis (ITT) – Study CS4<sup>1,2</sup>**



<sup>1</sup>Data for patients without a Month 15 visit were imputed using the multiple imputation method

<sup>2</sup>Error bars denote +/- standard error

Upon completion of Study CS4 (CHERISH) 125 patients enrolled in Study CS11 (SHINE), where all patients received Spinraza. The length of treatment ranged from 74 to 474 days (median 250 days) at the time of the interim analysis. A majority of Spinraza treated patients experienced stabilization or improvement in motor function, with the greatest benefit observed in those with earlier treatment initiation.

Of patients who initiated Spinraza treatment in Study CS4 (n=39), stabilization or additional improvements in mean HFMSE (0.2; SD 3.06) and RULM (0.7; SD 2.69) scores were observed from baseline to Study Day 265 in Study CS11.

Patients who initiated Spinraza treatment in Study CS11 (n=20) had a median age of 4.0 years (range 3 - 8 years). Of these patients, stabilization or improvement in mean HFMSE (1.4; SD 4.02) and RULM (2.1; SD 2.56) scores were observed from baseline to Study Day 265 in Study CS11.

These results are supported by 2 open label studies (study CS2 and study CS12). The analysis included 28 patients who received their first dose in study CS2, and then transferred to the extension phase, study CS12. The studies enrolled patients who were between 2 to 15 years of age at first dose. Of the 28 patients, 3 were at least 18 years of age at their last study visit. 1 out of 28 patients had 2 SMN2 gene copies, 21 had 3 copies, and 6 had 4 copies.

Patients were assessed over a 3 year treatment period. A sustained improvement was seen in patients with Type II SMA who experienced a mean improvement from baseline HFMSE score of 5.1 (SD 4.05, n=11) at Day 253, and 9.1 (SD 6.61, n=9) at Day 1050. The mean total score was 26.4 (SD 11.91) at Day 253 and 31.3 (SD 13.02) at Day 1050, no plateau was observed. Patients with Type III SMA demonstrated a mean improvement from baseline HFMSE score of 1.3 (SD 1.87, n=16) at Day 253 and 1.2 (SD 4.64, n=11) at Day 1050. The mean total score was 49.8 (SD 12.46) at Day 253 and 52.6 (SD 12.78) at 1050 days.

In patients with Type II SMA the Upper Limb Module test was conducted with mean improvement of 1.9 (SD 2.68, n=11) at Day 253 and 3.5 (SD 3.32, n=9) at Day 1050. The mean total score was 13.8 (SD 3.09) at Day 253 and 15.7 (SD 1.92) at Day 1050.

The 6MWT (six-minute walk test) was conducted for ambulatory patients only. In these patients, a mean improvement of 28.6 meters (SD 47.22, n=12) at Day 253 and 86.5 metres (SD 40.58, n=8) at Day 1050. The mean 6MWT distance was 278.5 meters (SD 206.46) at Day 253 and 333.6 metres (SD 176.47) at Day 1050. Two previously non-independent ambulatory patients (Type III) achieved independent walking, and one non-ambulatory patient (Type II) achieved independent walking.

An additional clinical study, CS7 (EMBRACE) was opened for patients not eligible for participation in Study CS3B or Study CS4 due to screening age or SMN2 copy number. CS7 is a phase 2, randomized, double-blind, sham-procedure study in symptomatic patients diagnosed with infantile-onset SMA ( $\leq 6$  months) or later-onset SMA ( $> 6$  months) and 2 or 3 copies of SMN2 (Part 1), followed by a long-term open label extension phase (Part 2). In Part 1 of the study, patients were followed for a median of 302 days.

All patients who received Spinraza were alive as of the early termination of Part 1, however, one patient in the control arm died at Study Day 289. In addition, no patients in the Spinraza or sham-control group required the use of permanent ventilation. Of the 13 patients with infantile-onset SMA, 7 of out 9 patients (78%; 95%CI: 45, 94) in the Spinraza group and 0 out of 4 patients (0%; 95%CI: 0, 60) in the sham group met the criteria for motor milestone response (according to HINE section 2:  $\geq 2$  point increase [or maximal score] in ability to kick OR  $\geq 1$  point increase in the motor milestones of head control, rolling, sitting, crawling, standing or walking and improvement in more categories of motor milestones than worsening). Of the 8 patients with later-onset SMA, 4 out of 5 patients (80%; 95% CI: 38, 96) in the Spinraza group and 2 out of 3 (67%; 95% CI: 21, 94) in the sham-control group met this definition of response.

### Pre-symptomatic infants


Study CS5 (NURTURE) is an open-label study in pre-symptomatic infants genetically diagnosed with SMA, who were enrolled at 6 weeks of age or younger. Patients in this study were deemed most likely to develop Type I or II SMA. Median age at first dose was 22 days.

An interim analysis was conducted when patients had been on study for median of 27.1 months (15.1 - 35.5 months) and were of a median age at last visit of 26.0 months (14.0 -34.3 months). At the interim analysis, all 25 patients (2 SMN2 gene copies, n=15; 3 SMN2 gene copies, n=10) were alive without permanent ventilation.. The primary endpoint, time to death or respiratory intervention (defined as invasive or non-invasive ventilation for  $\geq 6$  hours/day continuously for  $\geq 7$  consecutive days OR tracheostomy), could not be estimated as there were too few events. Four patients (2 SMN2 copies) required respiratory intervention  $>6$  hours/day continuously for  $\geq 7$  days, all of whom initiated ventilatory support during an acute reversible illness

Patients achieved milestones unexpected in Type I or II SMA and more consistent with normal development. At the interim analysis, all 25 (100%) patients had achieved the WHO motor milestone of sitting without support, 22 (88%) patients were walking with assistance. Among patients older than the WHO defined window for expected age of achievement (95<sup>th</sup> percentile), 17 of 22 (77%) had achieved walking alone. The mean CHOP INTEND score at last assessment was 61.0 (46 - 64) amongst patients with 2 SMN2 copies and 62.6 (58 - 64) amongst those with 3 SMN2 copies. All patients had the ability to suck and swallow at last assessment, with 22 (88%) infants achieving a maximal score on the HINE Section 1.

The proportion of patients developing clinically manifested SMA was assessed amongst patients who reached the Day 700 visit at the interim analysis (n=16). The protocol-defined criteria for clinically manifested SMA included age-adjusted weight below the fifth WHO percentile, a decrease of 2 or more major weight growth curve percentiles, the placement of a percutaneous gastric tube, and/or the inability to achieve expected age-appropriate WHO milestones (sitting without support, standing with assistance, hands-and-knees crawling, walking with assistance, standing alone and walking alone). At day 700, 7 out of 11 patients (64%) with 2 SMN2 gene copies and 0 out of 5 patients (0%) with 3 SMN2 copies, met the protocol-defined criteria of clinically manifested SMA, however, these patients were gaining weight and achieving WHO milestones, inconsistent with Type I SMA. A comparison of motor milestone achievement among the patients with symptomatic infantile-onset SMA and pre-symptomatic SMA is shown in Figure 3.

Figure 3: Change in HINE Motor Milestones versus Study days for Study CS3B (treated and sham-control), CS3A, CS5 and CS11



Population used in figure: CS5 subjects in the ITT set with SMN2 copy number denoted in parentheses, CS3A: SMN2 2 copy subjects, CS3B: Subjects with SMN2 2 copy in ITT set. For CS3B the data were windowed into intervals based on time from baseline. For each study, visits with n < 5 are not plotted.

## 5.2 Pharmacokinetic properties

Single- and multiple-dose pharmacokinetics (PK) of nusinersen, administered via intrathecal injection, were determined in paediatric patients diagnosed with SMA.

### Absorption

Intrathecal injection of nusinersen into the CSF allows nusinersen to be fully available for distribution from the CSF to the target central nervous system (CNS) tissues. Mean CSF trough concentrations of nusinersen accumulated approximately 1.4- to 3-fold after multiple loading and maintenance doses, and reached a steady state within approximately 24 months. Following intrathecal administration trough plasma concentrations of nusinersen were relatively low compared to the trough CSF concentration. Median plasma  $T_{max}$  values ranged from 1.7 to 6.0 hours. Mean plasma  $C_{max}$  and AUC values increased approximately dose proportionally over the evaluated dose range. There is no accumulation in plasma exposure measures ( $C_{max}$  and AUC) after multiple doses.

### Distribution

Autopsy data from patients (n=3) show that nusinersen administered intrathecally is broadly distributed within the CNS achieving therapeutic levels in the target spinal cord tissues. Presence of nusinersen was also demonstrated in neurons and other cell types in the spinal cord and brain, and peripheral tissues such as skeletal muscle, liver, and kidney.

### Biotransformation

Nusinersen is metabolized slowly and predominantly via exonuclease (3'- and 5') -mediated hydrolysis and is not a substrate for, or inhibitor or inducer of CYP450 enzymes.

## Elimination

The mean terminal elimination half-life in CSF is estimated at 135 to 177 days. The primary route of elimination is expected via urinary excretion of nusinersen and its metabolites.

## Interactions

*In vitro* studies indicated that nusinersen is not an inducer or inhibitor of CYP450-mediated oxidative metabolism and therefore should not interfere with other medicinal products for these metabolic pathways. Nusinersen is not a substrate or inhibitor of human BCRP, P-gp, OAT1, OAT3, OCT1, OCT2, OATP1B1, OATP1B3, or BSEP transporters.

## Characteristics in specific patient populations

### *Renal and hepatic impairment*

The pharmacokinetics of nusinersen in patients with renal or hepatic impairment has not been studied. The effect of hepatic or renal insufficiency as covariates could not be thoroughly evaluated in the population PK model given the rarity of patients displaying clinically relevant liver or kidney insufficiencies. Population PK analyses revealed no apparent correlation between hepatic and renal clinical chemistry markers and inter-subject variability.

### *Race*

The majority of patients studied were Caucasian. The population PK analysis suggests that race is unlikely to affect the PK of nusinersen.

## **5.3 Preclinical safety data**

### Carcinogenesis

Long-term studies in animals to evaluate the carcinogenic potential of nusinersen have not been performed.

### Mutagenesis

Nusinersen demonstrated no evidence of genotoxicity.

### Reproductive toxicity

Reproductive toxicology studies were conducted using subcutaneous administration of nusinersen in mice and rabbits. No impact on male or female fertility, or embryo-foetal development, or pre/post-natal development was observed.

### Toxicology

In repeat-dose toxicity studies (14-weeks and 53-weeks) of intrathecal administration to juvenile cynomolgus monkeys, nusinersen was well tolerated. The exception was an acute, transient deficit in lower spinal reflexes which occurred at the highest dose levels in each study (3 or 4 mg per dose; equivalent to 30 or 40 mg per intrathecal dose in patients). These effects were observed within several hours post-dose and generally resolved within 48 hours.

In the 53-week intrathecal dosing study in cynomolgus monkeys no toxicity effects were seen at levels up to 14-fold the recommended annual clinical maintenance dose.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Sodium dihydrogen phosphate dihydrate  
Disodium phosphate  
Sodium chloride  
Potassium chloride  
Calcium chloride dihydrate  
Magnesium chloride hexahydrate  
Sodium hydroxide (for pH adjustment)  
Hydrochloric acid (for pH adjustment)  
Water for injections

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

3 years

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C - 8°C).

Do not freeze.

Keep the vial in the outer carton in order to protect from light.

If no refrigeration is available, Spinraza may be stored in its original carton, protected from light at or below 30°C for up to 14 days.

Prior to administration, unopened vials of Spinraza can be removed from and returned to the refrigerator if necessary. If removed from the original carton, the total combined time out of refrigeration should not exceed 30 hours, at a temperature that does not exceed 25°C.

### **6.5 Nature and contents of container**

5 ml in a Type I glass vial with bromobutyl rubber stopper and an aluminium over-seal and plastic cap.

Pack size of one vial per carton.

### **6.6 Special precautions for disposal and other handling**

For single use only.

#### Instructions for preparation of the medicinal product before administration

1. The Spinraza vial should be inspected for particles prior to administration. If particles are observed and/or the liquid in the vial is not clear and colourless, the vial must not be used.
2. Aseptic technique should be used when preparing Spinraza solution for intrathecal administration.
3. The vial should be taken out of the refrigerator and allowed to warm to room temperature (25°C) without using external heat sources, prior to administration.
4. If the vial remains unopened and the solution is not used, it should be returned back to the refrigerator (see section 6.4).
5. Just prior to administration, remove the plastic cap and insert the syringe needle into the vial through the centre of the over-seal to remove the appropriate volume. Spinraza must not be diluted. The use of external filters is not required.
6. Once drawn in to the syringe, if the solution is not used within 6 hours, it must be discarded.

7. Any unused product or waste material must be disposed of in accordance with local requirements.

**7. MARKETING AUTHORISATION HOLDER**

Biogen Netherlands B.V.  
Prins Mauritslaan 13  
1171 LP Badhoevedorp  
The Netherlands

**8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/17/1188/001

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 30 May 2017

**10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.



## **ANNEX II**

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturer(s) responsible for batch release

Biogen (Denmark) Manufacturing ApS  
Biogen Allé 1  
DK - 3400 Hillerød  
Denmark

## B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

## C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

### • Periodic Safety Update Reports

The requirements for submission of periodic safety update reports for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

The marketing authorisation holder shall submit the first periodic safety update report for this product within 6 months following authorisation.

## D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

### • Risk Management Plan (RMP)

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

### • Obligation to conduct post-authorisation measures

The MAH shall complete, within the stated timeframe, the below measures:

Description	Due date
Post-authorisation efficacy study (PAES): In order to evaluate the long term efficacy and safety of nusinersen in symptomatic patients with spinal muscular atrophy, the MAH should conduct and submit the results of the Phase 3, open-label extension study (SHINE, CS11)	Submission of study results: Q2 2024.

Post-authorisation efficacy study (PAES): In order to evaluate the long term efficacy and safety of nusinersen in pre-symptomatic patients with spinal muscular atrophy, the MAH should conduct and submit the results of the Phase 2, open-label study (NURTURE (SM201))	Submission of study results: April 2023.
---	--

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Spinraza 12 mg solution for injection  
nusinersen

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each 5 ml vial contains nusinersen sodium equivalent to 12 mg nusinersen (2.4 mg/ml).

**3. LIST OF EXCIPIENTS**

Sodium dihydrogen phosphate dihydrate, disodium phosphate, sodium chloride, potassium chloride, calcium chloride dihydrate, magnesium chloride hexahydrate, sodium hydroxide, hydrochloric acid, water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Solution for injection  
1 vial

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
Intrathecal use.  
For single use only.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator.  
Do not freeze.  
Store in the original carton in order to protect from light.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Biogen Netherlands B.V.  
Prins Mauritslaan 13  
1171 LP Badhoevedorp  
The Netherlands

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/17/1188/001

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Justification for not including Braille accepted.

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC:  
SN:  
NN:

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**VIAL**

**1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

Spinraza 12 mg solution for injection  
nusinersen  
Intrathecal use

**2. METHOD OF ADMINISTRATION**

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

5 ml

**6. OTHER**



**B. PACKAGE LEAFLET**

## Package leaflet: Information for the user

### Spinraza 12 mg solution for injection

nusinersen

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

**Read all of this leaflet carefully before you or your child receives this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or nurse.
- If you or your child gets any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. What Spinraza is and what it is used for
2. What you need to know before you or your child are given Spinraza
3. How Spinraza is given
4. Possible side effects
5. How to store Spinraza
6. Contents of the pack and other information

#### 1. What Spinraza is and what it is used for

Spinraza contains the active substance *nusinersen* which belongs to a group of medicines known as *antisense oligonucleotides*. Spinraza is used to treat a genetic disease called *spinal muscular atrophy* (SMA).

**Spinal muscular atrophy** is caused by a shortage of a protein called *survival motor neuron* (SMN) in the body. This results in the loss of nerve cells in the spine, leading to weakness of the muscles in the shoulders, hips, thighs and upper back. It may also weaken the muscles used for breathing and swallowing.

Spinraza works by helping the body to produce more of the SMN protein that people with SMA are lacking. This reduces the loss of nerve cells and so may improve muscle strength.

#### 2. What you need to know before you or your child are given Spinraza

##### Spinraza must not be given:

- If you or your child are **allergic to nusinersen** or any of the other ingredients of this medicine (listed in section 6).

If you are not sure, talk to your doctor or nurse before you or your child are given Spinraza.

#### Warnings and precautions

There is a risk of side effects occurring after Spinraza is given by a lumbar puncture procedure (see section 3). This can include headaches, vomiting and back pain. There may also be difficulties with giving a medicine by this method in very young patients and those with scoliosis (twisted and curved spine).

Other products that are in the same group of medicines as Spinraza have been shown to affect the cells in the blood which help clotting. Before you or your child are given Spinraza your doctor may decide to do a blood test to check that your or your child's blood can clot properly. This may not be required every time you or your child are given Spinraza.

Other products that are in the same group of medicines as Spinraza have been shown to affect the kidneys. Before you or your child are given Spinraza your doctor may decide to do a urine test to check that your or your child's kidneys are working normally. This may not be required every time you or your child are given Spinraza.

There have been a small number of reports of patients developing hydrocephalus (a build-up of too much fluid around the brain) after Spinraza is given. Some of these patients had needed to have a device called a ventriculo-peritoneal shunt implanted to treat the hydrocephalus. If you notice any symptoms of increase in head size, decreased consciousness, persistent nausea, vomiting or headache; or other symptoms that cause you concern, please inform your or your child's doctor to seek necessary treatment. The benefits and risks of continuing Spinraza whilst having a "ventriculo-peritoneal shunt" in place are not known at present.

Talk to your doctor before you or your child are given Spinraza.

#### **Other medicines and Spinraza**

Tell your doctor if you or your child are taking, have recently taken any, or might take any other medicines in future.

#### **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before being given this medicine. It is preferable to avoid the use of Spinraza during pregnancy and breast-feeding.

#### **Driving and using machines**

Spinraza has no or negligible influence on the ability to drive and use machines.

#### **Spinraza contains a small amount of sodium**

Each dose of Spinraza contains less than 1 mmol (23 mg) sodium. It is essentially 'sodium-free' and can be used by people on a sodium-restricted diet.

### **3. How Spinraza is given**

The usual dose of Spinraza is 12 mg.

Spinraza is given;

- On the first day of treatment, day 0
- Then around day 14, day 28 and day 63
- Then once every 4 months.

Spinraza is given by injection into the lower back. This injection, called a lumbar puncture, is done by inserting a needle into the space around the spinal cord. This will be done by a doctor experienced in doing lumbar punctures. You or your child may also be given a medicine to make you relax or sleep during the procedure.

#### **How long to use Spinraza**

Your doctor will tell you how long you or your child need to receive Spinraza. Don't stop treatment with Spinraza unless your doctor tells you to.

### **If you or your child misses an injection**

If you or your child miss a dose of Spinraza, speak with your doctor so that Spinraza can be given as soon as possible.

If you have any questions about how Spinraza is given, ask your doctor.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Side effects related to the lumbar puncture may occur while Spinraza is being given or afterwards. The majority of these side effects are reported within 72 hours of the procedure.

### Very common (may affect more than 1 in 10 people)

- Back pain
- Headache
- Vomiting

### **Additional side effects not seen in clinical trials:**

#### Not known (frequency cannot be estimated from the available data)

- Serious infection related to lumbar puncture (e.g. meningitis)
- Hydrocephalus (a build-up of too much fluid around the brain)
- Meningitis not caused by an infection (inflammation of the membrane around the spinal cord and brain, which may present as neck stiffness, headache, fever, nausea and vomiting)
- Hypersensitivity (an allergic or allergic-like reaction that may include swelling of your face, lips or tongue, rash, or itching)

### **Reporting of side effects**

**If you or your child get any side effects, talk to your doctor or nurse.** This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Spinraza**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton after “EXP”. The expiry date refers to the last day of that month.

Store in a refrigerator (2°C to 8°C). Do not freeze.

Keep the vial in the outer carton in order to protect from light.

If no refrigeration is available, Spinraza may be stored in its original carton, protected from light at or below 30°C for up to 14 days.

Unopened vials of Spinraza can be removed from and returned to the refrigerator if necessary. If removed from the original carton, the total time out of refrigeration should not exceed 30 hours, at a temperature that does not exceed 25°C.

## 6. Contents of the pack and other information

### What Spinraza contains

- The active substance is nusinersen.
- Each 5 ml vial contains nusinersen sodium equivalent to 12 mg nusinersen.
- Each ml contains 2.4 mg of nusinersen.
- The other ingredients are sodium dihydrogen phosphate dihydrate, disodium phosphate, sodium chloride, potassium chloride, calcium chloride dihydrate, magnesium chloride hexahydrate, sodium hydroxide, hydrochloric acid, water for injections.

### What Spinraza looks like and contents of the pack

Spinraza is a clear colourless solution for injection.

Each carton of Spinraza contains one vial.

Each vial is for single use.

### Marketing Authorisation Holder

Biogen Netherlands B.V.  
Prins Mauritslaan 13  
1171 LP Badhoevedorp  
The Netherlands

### Manufacturer

Biogen (Denmark) Manufacturing ApS  
Biogen Allé 1  
DK - 3400 Hillerød  
Denmark

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

### België/Belgique/Belgien

Biogen Belgium N.V./S.A.  
Tél/Tel: +32 2 219 12 18

### Lietuva

Ewopharma AG Atstovybė  
Tel: +370 52 14 02 60

### България

ТП ЕВОФАРМА  
Тел.: +359 2 962 12 00

### Luxembourg/Luxemburg

Biogen Belgium N.V./S.A.  
Tél/Tel: +32 2 219 12 18

### Česká republika

Biogen (Czech Republic) s.r.o.  
Tel: +420 255 706 200

### Magyarország

Biogen Hungary Kft.  
Tel.: +36 (1) 899 9883

### Danmark

Biogen (Denmark) A/S  
Tlf: +45 77 41 57 57

### Malta

Pharma MT limited  
Tel: +356 213 37008/9

### Deutschland

Biogen GmbH  
Tel: +49 (0) 89 99 6170

### Nederland

Biogen Netherlands B.V.  
Tel: +31 20 542 2000

### Eesti

Ewopharma AG Eesti filiaal  
Tel: + 372 6 68 30 56

### Norge

Biogen Norway AS  
Tlf: +47 23 40 01 00

**Ελλάδα**

Genesis Pharma SA  
Τηλ: +30 210 8771500

**España**

Biogen Spain SL  
Tel: +34 91 310 7110

**France**

Biogen France SAS  
Tél: +33 (0)1 41 37 95 95

**Hrvatska**

Medis Adria d.o.o.  
Tel: +385 (0) 1 230 34 46

**Ireland**

Biogen Idec (Ireland) Ltd.  
Tel: +353 (0)1 463 7799

**Ísland**

Icepharma hf  
Sími: +354 540 8000

**Italia**

Biogen Italia s.r.l.  
Tel: +39 02 584 9901

**Κύπρος**

Genesis Pharma Cyprus Ltd  
Τηλ: +357 22 769946

**Latvija**

Ewopharma AG pārstāvniecība  
Tel: + 371 66 16 40 32

**Österreich**

Biogen Austria GmbH  
Tel: +43 1 484 46 13

**Polska**

Biogen Poland Sp. z o.o.  
Tel.: +48 22 351 51 00

**Portugal**

Biogen Portugal  
Tel.: +351 21 318 8450

**România**

Ewopharma AG Representative Office  
Tel: + 40 377 881 045

**Slovenija**

Biogen Pharma d.o.o.  
Tel.: +386 1 511 02 90

**Slovenská republika**

Biogen Slovakia s.r.o.  
Tel.: +421 2 323 340 08

**Suomi/Finland**

Biogen Finland Oy  
Puh/Tel: +358 207 401 200

**Sverige**

Biogen Sweden AB  
Tel: +46 8 594 113 60

**United Kingdom**

Biogen Idec Limited  
Tel: +44 (0) 1628 50 1000

**This leaflet was last revised in**

**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>

<----->

**The following information is intended for healthcare professionals only:**

1. The Spinraza vial should be inspected for particles prior to administration. If particles are observed and/or the liquid in the vial is not clear and colourless, the vial must not be used.
2. Aseptic technique should be used when preparing Spinraza solution for intrathecal administration.
3. The vial should be taken out of the refrigerator and allowed to warm to room temperature (25°C) without using external heat sources, prior to administration.

4. If the vial remains unopened and the solution is not used, it should be returned back to the refrigerator.
5. Just prior to administration, remove the plastic cap and insert the syringe needle into the vial through the center of the over-seal to remove the appropriate volume. Spinraza must not be diluted. The use of external filters is not required.
6. Spinraza is administered as an intrathecal bolus injection over 1 to 3 minutes, using a spinal anaesthesia needle.
7. The injection must not be administered in areas of the skin where there are signs of infection or inflammation.
8. It is recommended that the volume of CSF, equivalent to the volume of Spinraza to be injected, is removed prior to administration of Spinraza.
9. Once drawn into the syringe, if the solution is not used within 6 hours, it must be discarded.
10. Any unused product or waste material must be disposed of in accordance with local requirements.

**ANNEX IV**

**SCIENTIFIC CONCLUSIONS AND GROUNDS FOR THE VARIATION TO THE TERMS  
OF THE MARKETING AUTHORISATION(S)**



## **Scientific conclusions**

Taking into account the PRAC Assessment Report on the PSUR(s) for nusinersen, the scientific conclusions of CHMP are as follows:

As of 19 October 2018, 100 AEs of hypersensitivity due to the administration of nusinersen were identified from the clinical trial database. Ninety-eight of these were considered not or unlikely related to nusinersen. The remaining 2 events (rash and eosinophil count increased, both nonserious) were possibly related to nusinersen. A cumulative search of postmarketing data identified 37 possible hypersensitivity events; 20 medically confirmed. Time to onset was reported in 11 of these 20, and ranged from directly following the injection to 1 week after the injection. Two events, angioedema and serum sickness, were considered serious and related. There were no reports of a patient discontinuing treatment due to these events. Taken together, there is sufficient evidence to support a casual association between with nusinersen administration and hypersensitivity; and the PI should be updated accordingly. Prevention and treatment will be managed by routine clinical practice and no additional risk minimization measures or evaluations are required as part of the pharmacovigilance plan.

The CHMP agrees with the scientific conclusions made by the PRAC.

## **Grounds for the variation to the terms of the marketing authorisation(s)**

On the basis of the scientific conclusions for nusinersen the CHMP is of the opinion that the benefit-risk balance of the medicinal product(s) containing nusinersen is unchanged subject to the proposed changes to the product information

The CHMP recommends that the terms of the marketing authorisation(s) should be varied.