

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

## 1. NAME OF THE MEDICINAL PRODUCT

Symkevi 100 mg/150 mg film-coated tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 100 mg of tezacaftor and 150 mg of ivacaftor.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Film-coated tablet (tablet)

Yellow, capsule-shaped tablet debossed with “V100” on one side and plain on the other (dimensions 15.9 mm x 8.5 mm).

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Symkevi is indicated in a combination regimen with ivacaftor 150 mg tablets for the treatment of patients with cystic fibrosis (CF) aged 12 years and older who are homozygous for the *F508del* mutation or who are heterozygous for the *F508del* mutation and have one of the following mutations in the cystic fibrosis transmembrane conductance regulator (*CFTR*) gene: *P67L*, *R117C*, *L206W*, *R352Q*, *A455E*, *D579G*, *711+3A→G*, *S945L*, *S977F*, *R1070W*, *D1152H*, *2789+5G→A*, *3272-26A→G*, and *3849+10kbC→T*.

### 4.2 Posology and method of administration

Symkevi should only be prescribed by physicians with experience in the treatment of CF. If the patient's genotype is unknown, an accurate and validated genotyping method should be performed to confirm the presence of an indicated mutation using a genotyping assay.

#### Posology

*Adults and adolescents aged 12 years and older*

*In a combination regimen with ivacaftor 150 mg tablets*

The recommended dose is one Symkevi (tezacaftor 100 mg/ivacaftor 150 mg) tablet taken in the morning and one ivacaftor 150 mg tablet taken in the evening, approximately 12 hours apart with fat-containing food (see Method of administration).

#### *Missed dose*

If 6 hours or less have passed since the missed morning or evening dose, the patient should take the missed dose as soon as possible and continue on the original schedule.

If more than 6 hours have passed since the missed morning or evening dose, the patient should not take the missed dose. The next scheduled dose can be taken at the usual time.

More than one dose of either tablet should not be taken at the same time.

*Concomitant use of CYP3A inhibitors*

The dose of Symkevi and ivacaftor should be adjusted when co-administered with moderate and strong CYP3A inhibitors.

When co-administered with moderate CYP3A inhibitors (*e.g.*, fluconazole, erythromycin), the dose should be adjusted according to Table 1 (see sections 4.4 and 4.5).

<b>Table 1: Dosing recommendations for concomitant use with moderate CYP3A inhibitors</b>				
	<b>Day 1</b>	<b>Day 2</b>	<b>Day 3</b>	<b>Day 4*</b>
<b>Morning Dose</b>				
Tezacaftor 100 mg/ivacaftor 150 mg tablet	✓	-	✓	-
Ivacaftor 150 mg tablet	-	✓	-	✓
<b>Evening Dose</b>				
Ivacaftor 150 mg tablet	-	-	-	-
*Continue dosing with tezacaftor 100 mg/ivacaftor 150 mg or ivacaftor 150 mg tablets on alternate days.				

When co-administered with strong CYP3A inhibitors (*e.g.*, ketoconazole, itraconazole, posaconazole, voriconazole, telithromycin, and clarithromycin), the dose should be adjusted to one Symkevi tablet twice a week, taken approximately 3 to 4 days apart. The evening dose of ivacaftor should not be taken (see sections 4.4 and 4.5).

Special populations

*Elderly people*

The safety, efficacy and pharmacokinetics of Symkevi in combination with ivacaftor have been examined in a limited number of elderly patients. No dose adjustment specific to this patient population is required (see section 5.2).

*Renal impairment*

No dose adjustment is recommended for patients with mild or moderate renal impairment. Caution is recommended in patients with severe renal impairment or end-stage renal disease (see sections 4.4 and 5.2).

*Hepatic impairment*

For dose adjustment for patients with hepatic impairment, see Table 2. There is no experience of the use of Symkevi in patients with severe hepatic impairment (Child-Pugh Class C); therefore, its use is not recommended unless the benefits outweigh the risks. In such cases, Symkevi should be used at a reduced dose (see sections 4.4 and 5.2).

<b>Table 2: Dosing recommendations for patients with hepatic impairment</b>			
	<b>Mild (Child-Pugh Class A)</b>	<b>Moderate (Child-Pugh Class B)</b>	<b>Severe (Child-Pugh Class C)</b>
<b>Morning</b>	No dose adjustment	One tablet of tezacaftor 100 mg/ivacaftor 150 mg once daily	Starting dose: One tablet of tezacaftor 100 mg/ivacaftor 150 mg once daily. Dosing intervals should be modified according to clinical response and tolerability.
<b>Evening</b>	No dose adjustment	No ivacaftor 150 mg dose	No ivacaftor 150 mg dose

*Paediatric population*

The safety and efficacy of Symkevi in children aged less than 12 years has not yet been established. No data are available (see section 5.1).

### Method of administration

For oral use. Patients should be instructed to swallow the tablets whole. The tablets should not be chewed, crushed, or broken before swallowing.

Both Symkevi and ivacaftor tablets should be taken with fat-containing food, such as food recommended in standard nutritional guidelines (see section 5.2).

Food or drink containing grapefruit or Seville oranges should be avoided during treatment (see section 4.5).

### **4.3 Contraindications**

Hypersensitivity to the active substance(s) or to any of the excipients listed in section 6.1.

### **4.4 Special warnings and precautions for use**

Symkevi in combination with ivacaftor should not be prescribed in patients with CF who are heterozygous for the *F508del* mutation and have a second *CFTR* mutation not listed in section 4.1.

### Effect on liver function tests

Elevated transaminases are common in patients with CF, and have been observed in some patients treated with Symkevi in combination with ivacaftor, as well as with ivacaftor monotherapy. Assessments of transaminases (ALT or AST) are recommended for all patients prior to initiating treatment, every 3 months during the first year of treatment, and annually thereafter. For patients with a history of transaminase elevations, more frequent monitoring should be considered. In the event of significant elevations of transaminases (*e.g.*, patients with ALT or AST >5 x the upper limit of normal (ULN), or ALT or AST >3 x ULN with bilirubin >2 x ULN), dosing should be interrupted and laboratory tests closely followed until the abnormalities resolve. Following resolution of transaminase elevations, the benefits and risks of resuming treatment should be considered (see section 4.8).

### Hepatic impairment

The use of Symkevi is not recommended in patients with severe hepatic impairment unless the benefits are expected to outweigh the risks (see sections 4.2 and 5.2).

### Renal impairment

Caution is recommended while using Symkevi in combination with ivacaftor in patients with severe renal impairment or end-stage renal disease (see sections 4.2 and 5.2).

### Patients after organ transplantation

Symkevi in combination with ivacaftor has not been studied in patients with CF who have undergone organ transplantation. Therefore, use in transplanted patients is not recommended. See section 4.5 for interactions with ciclosporin or tacrolimus.

### Interactions with medicinal products

#### *CYP3A inducers*

Exposure to tezacaftor and ivacaftor may be reduced by the concomitant use of CYP3A inducers, potentially resulting in reduced efficacy of Symkevi and ivacaftor. Therefore, co-administration with strong CYP3A inducers is not recommended (see section 4.5).

#### *CYP3A inhibitors*

The dose of Symkevi and ivacaftor should be adjusted when used concomitantly with strong or moderate CYP3A inhibitors (see section 4.5 and Table 1 in section 4.2).

### Cataracts

Cases of non-congenital lens opacities without impact on vision have been reported in paediatric patients treated with Symkevi in combination with ivacaftor, as well as with ivacaftor monotherapy.

Although other risk factors were present in some cases (such as corticosteroid use and exposure to radiation), a possible risk attributable to treatment cannot be excluded. Baseline and follow-up ophthalmological examinations are recommended in paediatric patients initiating treatment with Symkevi in combination with ivacaftor (see section 5.3).

#### Sodium

This medicine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

### **4.5 Interaction with other medicinal products and other forms of interaction**

#### Medicinal products affecting the pharmacokinetics of tezacaftor and ivacaftor

##### *CYP3A inducers*

Tezacaftor and ivacaftor are substrates of CYP3A (ivacaftor is a sensitive substrate of CYP3A). Concomitant use of CYP3A inducers may result in reduced exposures and thus reduced efficacy of Symkevi and ivacaftor. Co-administration of ivacaftor with rifampicin, a strong CYP3A inducer, significantly decreased ivacaftor exposure [area under the curve (AUC)] by 89%. Tezacaftor exposures can also be expected to decrease significantly during co-administration with strong CYP3A inducers; therefore, co-administration with strong CYP3A inducers is not recommended.

Examples of strong CYP3A inducers include:

- rifampicin, rifabutin, phenobarbital, carbamazepine, phenytoin, and St. John's wort (*Hypericum perforatum*)

##### *CYP3A inhibitors*

Co-administration with itraconazole, a strong CYP3A inhibitor, increased tezacaftor exposure (measured as AUC) by 4-fold and increased ivacaftor AUC by 15.6-fold. The dose of Symkevi should be adjusted when co-administered with strong CYP3A inhibitors (see Table 1 in section 4.2).

Examples of strong CYP3A inhibitors include:

- ketoconazole, itraconazole, posaconazole, and voriconazole
- telithromycin and clarithromycin

Physiologically based pharmacokinetic modeling suggested co-administration with fluconazole, a moderate CYP3A inhibitor, may increase tezacaftor exposure (AUC) by approximately 2-fold. Co-administration of fluconazole increased ivacaftor AUC by 3-fold. The dose of Symkevi and ivacaftor should be adjusted when co-administered with moderate CYP3A inhibitors (see Table 1 in section 4.2).

Examples of moderate CYP3A inhibitors include:

- fluconazole
- erythromycin

Co-administration with grapefruit juice, which contains one or more components that moderately inhibit CYP3A, may increase exposure of ivacaftor and tezacaftor; therefore, food or drink containing grapefruit or Seville oranges should be avoided during treatment (see section 4.2).

##### *Potential for tezacaftor/ivacaftor to interact with transporters*

*In vitro* studies showed that tezacaftor is a substrate for the uptake transporter OATP1B1, and efflux transporters P-gp and Breast Cancer Resistance Protein (BCRP). Tezacaftor is not a substrate for OATP1B3. Exposure to tezacaftor is not expected to be affected significantly by concomitant inhibitors of OATP1B1, P-gp, or BCRP due to its high intrinsic permeability and low likelihood of being excreted intact. However, exposure to M2-TEZ (tezacaftor metabolite) may be increased by inhibitors of P-gp. Therefore, caution should be used when P-gp inhibitors are used with Symkevi.

*In vitro* studies showed that ivacaftor is not a substrate for OATP1B1, OATP1B3, or P-gp. Ivacaftor and its metabolites are substrates of BCRP *in vitro*. Due to its high intrinsic permeability and low likelihood of being excreted intact, co-administration of BCRP inhibitors is not expected to alter exposure of ivacaftor and M1-IVA, while any potential changes in M6-IVA exposures are not expected to be clinically relevant.

Symkevi may inhibit OATP1B1 and increase the exposures of medicinal products that are substrates of OATP1B1. When used concomitantly with OATP1B1 substrates, caution should be used.

#### *Ciprofloxacin*

Co-administration of ciprofloxacin did not affect the exposure of ivacaftor or tezacaftor. No dose adjustment is required when Symkevi is co-administered with ciprofloxacin.

#### Medicinal products affected by tezacaftor and ivacaftor

##### *CYP2C9 substrates*

Ivacaftor may inhibit CYP2C9; therefore, monitoring of the international normalized ratio (INR) is recommended during co-administration of warfarin with Symkevi given in combination with ivacaftor. Other medicinal products for which exposure may be increased include glimepiride and glipizide; these medicinal products should be used with caution.

##### *CYP3A, digoxin and other P-gp Substrates*

##### *CYP3A substrates*

Co-administration with (oral) midazolam, a sensitive CYP3A substrate, did not affect midazolam exposure. No dose adjustment of CYP3A substrates is required when co-administered with Symkevi in combination with ivacaftor.

##### *Digoxin and other P-gp substrates*

Co-administration with digoxin, a sensitive P-gp substrate, increased digoxin exposure by 1.3-fold, consistent with weak inhibition of P-gp by ivacaftor. Administration of Symkevi in combination with ivacaftor may increase systemic exposure of medicinal products that are sensitive substrates of P-gp, which may increase or prolong their therapeutic effect and adverse reactions. When used concomitantly with digoxin or other substrates of P-gp with a narrow therapeutic index, such as ciclosporin, everolimus, sirolimus, and tacrolimus, caution and appropriate monitoring should be used.

##### *Hormonal contraceptives*

Symkevi in combination with ivacaftor has been studied with an estrogen/progesterone oral contraceptive and was found to have no significant effect on the exposures of the hormonal contraceptive. Symkevi and ivacaftor are not expected to modify the efficacy of hormonal contraceptives.

#### Paediatric population

Interaction studies have only been performed in adults.

## **4.6 Fertility, pregnancy and lactation**

### Pregnancy

There are no or limited amount of data (less than 300 pregnancy outcomes) from the use of tezacaftor or ivacaftor in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3). As a precautionary measure, it is preferable to avoid the use of therapy during pregnancy.

### Breast-feeding

It is unknown whether tezacaftor, ivacaftor, or their metabolites are excreted in human milk. Available pharmacokinetic/toxicological data in animals have shown excretion of tezacaftor and ivacaftor into the milk of lactating female rats (see section 5.3). A risk to the newborns/infants cannot be excluded.

A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

#### Fertility

##### *Tezacaftor*

There are no data available on the effect of tezacaftor on fertility in humans. Tezacaftor had no effects on fertility and reproductive performance indices in male and female rats at doses up to 100 mg/kg/day.

##### *Ivacaftor*

There are no data available on the effect of ivacaftor on fertility in humans. Ivacaftor had an effect on fertility in rats (see section 5.3).

#### **4.7 Effects on ability to drive and use machines**

Symkevi in combination with ivacaftor has a minor influence on the ability to drive and use machines. Dizziness has been reported in patients receiving Symkevi in combination with ivacaftor, as well as ivacaftor monotherapy (see section 4.8). Patients experiencing dizziness should be advised not to drive or use machines until symptoms abate.

#### **4.8 Undesirable effects**

##### Summary of the safety profile

The most common adverse reactions experienced by patients aged 12 years and older who received Symkevi in combination with ivacaftor in the pooled, placebo-controlled, Phase 3 studies were headache (14% versus 11% on placebo) and nasopharyngitis (12% versus 10% on placebo).

##### Tabulated list of adverse reactions

Table 3 reflects adverse reaction observed with Symkevi in combination with ivacaftor in patients aged 12 years and older based on pooled data from three double-blind, placebo-controlled Phase 3 clinical studies. Table 4 reflects adverse reactions observed in clinical studies in patients treated with ivacaftor monotherapy. Adverse reactions are listed by MedDRA system organ class and frequency: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); not known (cannot be estimated from the available data).

<b>MedDRA System Organ Class</b>	<b>Adverse Reaction</b>	<b>Frequency</b>
Infections and infestations	Nasopharyngitis	very common
Nervous system disorder	Headache	very common
	Dizziness	common
Respiratory, thoracic and mediastinal disorders	Sinus congestion	common
Gastrointestinal disorders	Nausea	common

<b>MedDRA System Organ Class</b>	<b>Adverse Reactions</b>	<b>Frequency</b>
Infections and infestations	Upper respiratory tract infection, Nasopharyngitis	very common
	Rhinitis	common
Nervous system disorders	Headache, Dizziness	very common

<b>Table 4: Adverse reactions in patients treated with ivacaftor monotherapy</b>		
<b>MedDRA System Organ Class</b>	<b>Adverse Reactions</b>	<b>Frequency</b>
Ear and labyrinth disorders	Ear pain, Ear discomfort, Tinnitus, Tympanic membrane hyperaemia, Vestibular disorder	common
	Ear congestion	uncommon
Respiratory, thoracic and mediastinal disorders	Oropharyngeal pain, Nasal congestion	very common
	Sinus congestion, Pharyngeal erythema	common
Gastrointestinal disorders	Abdominal pain, Diarrhoea	very common
Hepatobiliary disorders	Transaminase elevations	very common
Skin and subcutaneous tissue disorders	Rash	very common
Reproductive system and breast disorders	Breast mass	common
	Breast inflammation, Gynaecomastia, Nipple disorder, Nipple pain	uncommon
Investigations	Bacteria in sputum	very common

Safety data from an interim safety analysis performed on 867 patients in a long-term safety and efficacy rollover study (Study 3), including 326 patients with  $\geq 48$  weeks of cumulative treatment with Symkevi given in combination with ivacaftor, were consistent with the safety data from the placebo-controlled Phase 3 studies.

#### Description of selected adverse reactions

##### Laboratory abnormalities

##### *Transaminase elevations*

During the placebo-controlled Phase 3 studies (up to 24 weeks), the incidence of maximum transaminase (ALT or AST)  $>8$ ,  $>5$ , or  $>3$  x ULN were similar between Symkevi in combination with ivacaftor- and placebo-treated patients; 0.2%, 1.0%, and 3.4% in Symkevi in combination with ivacaftor-treated patients, and 0.4%, 1.0%, and 3.4% in placebo-treated patients. One patient (0.2%) on therapy and two patients (0.4%) on placebo permanently discontinued treatment for elevated transaminases. No patients treated with Symkevi in combination with ivacaftor experienced a transaminase elevation  $>3$  x ULN associated with elevated total bilirubin  $>2$  x ULN.

##### Paediatric population

The safety profile is generally consistent among adolescents and adult patients.

##### Other special populations

The safety profile of Symkevi in combination with ivacaftor, including respiratory events (*e.g.*, chest discomfort, dyspnea, and respiration abnormal), was generally similar across all subgroups of patients, including analysis by age, gender, and baseline percent predicted FEV<sub>1</sub> (ppFEV<sub>1</sub>).

##### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).



## 4.9 Overdose

There are no known risks due to overdose with Symkevi and there is no specific antidote available in the event of overdose. Treatment of overdose consists of general supportive measures including monitoring of vital signs and observation of the clinical status of the patient.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Other respiratory system products; ATC code: R07AX31

#### Mechanism of action

Tezacaftor is a selective CFTR corrector that binds to the first Membrane Spanning Domain (MSD-1) of CFTR. Tezacaftor facilitates the cellular processing and trafficking of normal or multiple mutant forms of CFTR (including F508del-CFTR) to increase the amount of CFTR protein delivered to the cell surface, resulting in increased chloride transport *in vitro*.

Ivacaftor is a CFTR potentiator that potentiates the channel-open probability (or gating) of CFTR at the cell surface to increase chloride transport. For ivacaftor to function CFTR protein must be present at the cell surface. Ivacaftor can potentiate the CFTR protein delivered to the cell surface by tezacaftor, leading to a further enhancement of chloride transport than either active substance alone. The combination targets the abnormal CFTR protein by increasing the quantity and function of CFTR at the cell surface and subsequently increasing airway surface liquid height, and ciliary beat frequency *in vitro* in human bronchial epithelial (HBE) cells from homozygous F508del CF patients. The exact mechanisms by which tezacaftor improves cellular processing and trafficking of F508del-CFTR and ivacaftor potentiates F508del-CFTR are not known.

#### Pharmacodynamic effects

##### *Effects on sweat chloride*

In Study 1 (patients homozygous for the *F508del* mutation), the treatment difference between Symkevi in combination with ivacaftor and placebo in mean absolute change from baseline in sweat chloride through Week 24, was -10.1 mmol/L (95% CI: -11.4, -8.8; nominal  $P < 0.0001^*$ ).

In Study 2 (patients heterozygous for the *F508del* mutation and a second mutation associated with residual CFTR activity), the treatment difference in mean absolute change from baseline in sweat chloride through Week 8 was -9.5 mmol/L (95% CI: -11.7, -7.3; nominal  $P < 0.0001^*$ ) between Symkevi in combination with ivacaftor and placebo, and -4.5 mmol/L (95% CI: -6.7, -2.3; nominal  $P < 0.0001^*$ ) between ivacaftor and placebo.

\*Nominal p-value, based on hierarchical testing procedure.

##### *ECG evaluation*

Neither tezacaftor nor ivacaftor prolong the QTcF interval in healthy subjects at 3 times the therapeutic dose.

#### Clinical efficacy

The efficacy of Symkevi in combination with ivacaftor 150 mg tablet in patients with CF was demonstrated in two Phase 3, double-blind, controlled studies (Study 1 and Study 2), and one Phase 3, open-label extension study (Study 3).

Study 1 was a 24-week, randomized, double-blind, placebo-controlled study. A total of 504 patients aged 12 years and older (mean age 26.3 years) who were homozygous for the *F508del* mutation in the *CFTR* gene were randomized (1:1 randomization: 248 Symkevi in combination with ivacaftor, 256 placebo). Patients had a percent predicted forced expiratory volume in one second (ppFEV<sub>1</sub>) at screening between 40 to 90%. The mean ppFEV<sub>1</sub> at baseline was 60.0% (range: 27.8% to 96.2%).

Study 2 was a randomized, double-blind, placebo-controlled, 2-period, 3-treatment, 8-week crossover study. A total of 244 patients aged 12 years and older (mean age 34.8 years) who were heterozygous for the *F508del* mutation and a second mutation associated with residual CFTR activity were randomized to and received sequences of treatment that included Symkevi in combination with ivacaftor, ivacaftor, and placebo. Patients had a ppFEV<sub>1</sub> at screening between 40 to 90%. The mean ppFEV<sub>1</sub> at baseline was 62.3% (range: 34.6% to 93.5%).

Patients in Studies 1 and 2 continued on their standard-of-care CF therapies during the studies (*e.g.*, bronchodilators, inhaled antibiotics, dornase alfa, and hypertonic saline), and were eligible to roll over into a 96-week open-label extension study (Study 3). Patients had a confirmed genotype of a protocol-specified *CFTR* mutation, and a confirmed diagnosis of CF.

Patients with a history of colonization with organisms associated with a more rapid decline in pulmonary status such as *Burkholderia cenocepacia*, *Burkholderia dolosa*, or *Mycobacterium abscessus*, or who had two or more abnormal liver function tests at screening (ALT, AST, AP, GGT  $\geq 3$  x ULN or total bilirubin  $\geq 2$  x ULN) or AST or ALT  $\geq 5$  x ULN, were excluded from both studies.

***Study in patients with CF who were homozygous for the F508del mutation in the CFTR gene (Study 1)***

In Study 1 treatment with Symkevi in combination with ivacaftor resulted in a statistically significant improvement in ppFEV<sub>1</sub> (Table 5). The treatment difference between Symkevi (in combination with ivacaftor) and placebo for the primary endpoint of mean absolute change (95% CI) in ppFEV<sub>1</sub> from baseline through Week 24 was 4.0 percentage points (95% CI: 3.1, 4.8;  $P < 0.0001$ ). Mean improvement in ppFEV<sub>1</sub> was observed at the first assessment on Day 15 and sustained throughout the 24-week treatment period. Improvements in ppFEV<sub>1</sub> were observed regardless of age, sex, baseline ppFEV<sub>1</sub>, colonization with *Pseudomonas*, concomitant use of standard-of-care medications for CF, and geographic region. See Table 5 for a summary of primary and key secondary outcomes.

<b>Table 5: Primary and key secondary efficacy analyses, full analysis set (Study 1)</b>			
<b>Analysis</b>	<b>Statistic</b>	<b>Placebo N=256</b>	<b>Symkevi in combination with Ivacaftor N=248</b>
<b>Primary</b>			
ppFEV <sub>1</sub> Baseline value	n/N Mean (SD)	256/256 60.4 (15.7)	247/248 59.6 (14.7)
	n/N Within-group change LS mean (95% CI)	256/256 -0.6 (-1.3, 0.0)	245/248 3.4 (2.7, 4.0)
Average absolute change from baseline through Week 24 (percentage points)**	Treatment difference LS mean (95% CI)	4.0 (3.1, 4.8)	
	<i>P</i> value	$P < 0.0001^*$	
<b>Key Secondary</b>			
ppFEV <sub>1</sub> Baseline value	n/N Mean (SD)	256/256 60.4 (15.7)	247/248 59.6 (14.7)
	n/N Within-group change LS mean (95% CI)	256/256 -0.5 (-1.7, 0.6)	245/248 6.3 (5.1, 7.4)
Relative change from baseline through Week 24 (%)**	Treatment difference LS mean (95% CI)	6.8 (5.3, 8.3)	
	<i>P</i> value	$P < 0.0001^*$	

<b>Analysis</b>	<b>Statistic</b>	<b>Placebo N=256</b>	<b>Symkevi in combination with Ivacaftor N=248</b>
Pulmonary exacerbations	Number of subjects with events (n)/N	88/256	62/248
	Number of events (estimated event rate per year <sup>†</sup> )	122 (0.99)	78 (0.64)
Number of pulmonary exacerbations from baseline through Week 24	Rate ratio (RR) (95% CI) <i>P</i> value	0.65 (0.48, 0.88) <i>P</i> =0.0054*	
BMI Baseline value	n/N Mean (SD)	256/256 21.12 (2.88)	248/248 20.96 (2.95)
	n/N Within-group change LS mean (95% CI)	245/256 0.12 (0.03, 0.22)	237/248 0.18 (0.08, 0.28)
Absolute change from baseline at Week 24 (kg/m <sup>2</sup> )**	Treatment difference LS mean (95% CI) <i>P</i> value	0.06 (-0.08, 0.19) <i>P</i> =0.4127 <sup>#</sup>	
CFQ-R respiratory domain score Baseline value	n/N Mean (SD)	256/256 69.9 (16.6)	248/248 70.1 (16.8)
	n/N Within-group change LS mean (95% CI)	256/256 -0.1 (-1.6, 1.4)	246/248 5.0 (3.5, 6.5)
Absolute change from baseline through Week 24 (points)**	Treatment difference LS mean (95% CI) <i>P</i> value	5.1 (3.2, 7.0) nominal <i>P</i> <0.0001 <sup>±</sup>	
ppFEV <sub>1</sub> : percent predicted forced expiratory volume in 1 second; SD: Standard Deviation; LS mean: least squares mean; CI: confidence interval; BMI: body mass index; CFQ-R: Cystic Fibrosis Questionnaire-Revised. **Mixed Effect model for repeated measures with treatment, visit, treatment-by-visit interaction, sex, age group (<18, >=18 years) at screening, baseline value, and baseline value-by-visit interaction as fixed effect. *Indicates statistical significance confirmed in the hierarchical testing procedure. †Estimated event rate per year calculated using 48 weeks per year. # <i>P</i> value not statistically significant. ±Nominal <i>p</i> value, based on hierarchical testing procedure.			

Symkevi in combination with ivacaftor was associated with a lower event rate per year of severe pulmonary exacerbations requiring hospitalization or IV antibiotic therapy (0.29) compared to placebo (0.54). The rate ratio versus placebo was 0.53 (95% CI: 0.34, 0.82; nominal *P*= 0.0042). Pulmonary exacerbations requiring IV antibiotic therapy were lower in the treatment group compared to placebo (RR: 0.53 [95% CI: 0.34, 0.82]; nominal *P*=0.0042). Pulmonary exacerbations requiring hospitalizations were similar between treatment groups (RR: 0.78 [95% CI: 0.44, 1.36]; *P*=0.3801).

BMI increased in both treatment groups (Symkevi in combination with ivacaftor: 0.18 kg/m<sup>2</sup>, placebo: 0.12 kg/m<sup>2</sup>). The treatment difference of 0.06 kg/m<sup>2</sup> for mean change in BMI from baseline to Week 24 (95% CI: -0.08, 0.19) was not statistically significant (*P*=0.4127).

For CFQ-R respiratory domain score (a measure of respiratory symptoms relevant to patients with CF including cough, sputum production, and difficulty breathing) the percentage of subjects with at least a

4 point-increase from baseline (minimal clinically important difference) was 51.1% for Symkevi and 35.7% for placebo at Week 24.

***Study in patients with CF who were heterozygous for the F508del mutation and a second mutation associated with residual CFTR activity (Study 2)***

Of the 244 patients enrolled in Study 2 the following indicated mutations associated with residual CFTR activity were represented: *P67L, R117C, L206W, R352Q, A455E, D579G, 711+3A→G, S945L, S977F, R1070W, D1152H, 2789+5G→A, 3272-26A→G, and 3849+10kbC→T.*

In Study 2 treatment with Symkevi in combination with ivacaftor resulted in a statistically significant improvement in ppFEV<sub>1</sub> (Table 6). The treatment difference between Symkevi in combination with ivacaftor- and placebo-treated patients for the primary endpoint of mean absolute change in ppFEV<sub>1</sub> from study baseline to the average of Week 4 and Week 8 was 6.8 percentage points (95% CI: 5.7, 7.8; *P*<0.0001). The treatment difference between ivacaftor alone- and placebo-treated patients was 4.7 percentage points (95% CI: 3.7, 5.8; *P*<0.0001) and 2.1 percentage points (95% CI: 1.2, 2.9) between Symkevi in combination with ivacaftor- and ivacaftor alone-treated patients. Mean improvement in ppFEV<sub>1</sub> was observed at the first assessment on Day 15 and sustained throughout the 8-week treatment period. Improvements in ppFEV<sub>1</sub> were observed regardless of age, disease severity, sex, mutation class, colonization with *Pseudomonas*, concomitant use of standard-of-care medications for CF, and geographic region. See Table 6 for a summary of primary and key secondary outcomes.

<b>Table 6: Primary and key secondary efficacy analyses, full analysis set (Study 2)</b>				
<b>Analysis</b>	<b>Statistic</b>	<b>Placebo N=161</b>	<b>Ivacaftor N=156</b>	<b>Symkevi in Combination with Ivacaftor N=161</b>
ppFEV <sub>1</sub> Baseline value	n/N Mean (SD)	161/161 62.2 (14.3)	156/156 62.1 (14.6)	161/161 62.1 (14.7)
	n/N Within-group change LS mean (95% CI)	160/161 -0.3 (-1.2, 0.6)	156/156 4.4 (3.5, 5.3)	159/161 6.5 (5.6, 7.3)
Absolute change from baseline to the average of Week 4 and Week 8 (percentage points)**	Treatment difference versus placebo LS mean (95% CI) P value	NA NA	4.7 (3.7, 5.8) P<0.0001*	6.8 (5.7, 7.8) P<0.0001*
	Treatment difference versus IVA LS mean (95% CI)	NA	NA	2.1 (1.2, 2.9)
CFQ-R respiratory domain score Baseline value	n/N Mean (SD)	161/161 68.7 (18.3)	156/156 67.9 (16.9)	161/161 68.2 (17.5)
	n/N Within-group change LS mean (95% CI)	160/161 -1.0 (-2.9, 1.0)	156/156 8.7 (6.8, 10.7)	161/161 10.1 (8.2, 12.1)
Absolute change from baseline to the average of Week 4 and Week 8 (points)**	Treatment difference versus placebo LS mean (95% CI) P value	NA NA	9.7 (7.2, 12.2) P<0.0001*	11.1 (8.7, 13.6) P<0.0001*
	Treatment difference versus IVA LS mean (95% CI)	NA	NA	1.4 (-1.0, 3.9)
ppFEV <sub>1</sub> : percent predicted forced expiratory volume in 1 second; SD: Standard Deviation; LS mean: least squares mean; CI: confidence interval; NA: not applicable; IVA: ivacaftor; CFQ-R: Cystic Fibrosis Questionnaire-Revised. ** Linear Mixed Effects model with treatment, period, and study baseline ppFEV <sub>1</sub> as fixed effects and subject as a random effect. *Indicates statistical significance confirmed in the hierarchical testing procedure.				

### **Subgroup analysis of patients with severe lung dysfunction (ppFEV<sub>1</sub> <40)**

Study 1 and Study 2 included a total of 39 patients treated with Symkevi in combination with ivacaftor with ppFEV<sub>1</sub> <40. There were 23 patients with ppFEV<sub>1</sub> <40 at baseline receiving Symkevi and 24 patients receiving placebo in Study 1. The mean treatment difference between Symkevi and placebo-treated patients for absolute change in ppFEV<sub>1</sub> through Week 24 in this subgroup was 3.5 percentage points (95% CI: 1.0, 6.1). There were 16 patients with ppFEV<sub>1</sub> <40 at baseline receiving Symkevi, 13 receiving ivacaftor and 15 receiving placebo in Study 2. The mean treatment difference between Symkevi and placebo-treated patients for absolute change in ppFEV<sub>1</sub> through the average of Week 4 and Week 8 was 4.4 percentage points (95% CI: 1.1, 7.8). The mean treatment difference between ivacaftor and placebo-treated patients was 4.4 percentage points (95% CI: 0.9, 7.9).

### **Long-term safety and efficacy rollover study (Study 3)**

An ongoing, Phase 3, open-label, multicenter, rollover, 96-week study to evaluate the safety and efficacy of long-term treatment with Symkevi in combination with ivacaftor is being conducted with patients from Studies 1 (n=459) and 2 (n=226). An **interim analysis** was conducted through Week 24

for patients from Study 1 and Week 16 for patients from Study 2. As efficacy was a secondary objective for Study 3 there was no multiplicity adjustment.

Patients who received placebo in both Study 1 and Study 2 demonstrated improvements in ppFEV<sub>1</sub> when treated with Symkevi in combination with ivacaftor in Study 3 [Study 1: within-group change=4.2 (0.5) percentage points, Study 2: within-group change=4.9 (0.6) percentage points]. Patients who received Symkevi in combination with ivacaftor in Study 1 and Study 2, and continued on treatment, showed sustained improvements in ppFEV<sub>1</sub> through 48 weeks (Week 24 Study 3) and through 24 weeks (Week 16 Study 3), respectively.

Similar trends were observed for CFQ-R respiratory domain score, pulmonary exacerbation rate and BMI.

### ***Paediatric population***

#### **Adolescents aged 12 years and older**

Adolescents were included together with adults in the trials.

#### ***Adolescent patients with CF who were homozygous for the F508del mutation in the CFTR gene***

The mean absolute change (SE) from baseline in ppFEV<sub>1</sub> was 3.5 (0.6) percentage points in the Symkevi in combination with ivacaftor group and -0.4 (0.6) percentage points in the placebo group in Study 1. Patients who received Symkevi in combination with ivacaftor in Study 1 and continued on treatment showed sustained improvements in ppFEV<sub>1</sub> through 48 weeks [within-group change=-0.8 (0.8) percentage points from Study 3 baseline]. Patients who were previously treated with placebo and received Symkevi in combination with ivacaftor in Study 3 showed an increase of 5.3 (0.7) percentage points.

The mean absolute change (SE) from baseline in BMI z-value was -0.01(0.05) kg/m<sup>2</sup> in the Symkevi in combination with ivacaftor group and 0.00 (0.05) kg/m<sup>2</sup> in the placebo group in Study 1. In Study 3, the change in BMI z-value in the Symkevi in combination with ivacaftor group was maintained and patients previously treated with placebo showed an increase of 0.10 (0.05) kg/m<sup>2</sup>.

#### ***Adolescent patients with CF who were heterozygous for the F508del mutation and a second mutation associated with residual CFTR activity***

The mean absolute change (SE) from baseline in ppFEV<sub>1</sub> was 11.7 (1.2) percentage points in the Symkevi in combination with ivacaftor group, 7.6 (1.2) percentage points in the ivacaftor group and -0.4 (1.2) percentage points in the placebo group in Study 2. Patients who received Symkevi in combination with ivacaftor in Study 2 and continued on treatment showed sustained improvements in ppFEV<sub>1</sub> through 24 weeks [within-group change=0.7 (1.5) percentage points from Study 3 baseline]. Patients who were previously treated with ivacaftor and placebo and received Symkevi in combination with ivacaftor in Study 3 showed an increase of 1.6 (1.6) percentage points and 7.2 (1.2) percentage points, respectively.

The mean absolute change (SE) from baseline in BMI z-value was 0.24 (0.07) kg/m<sup>2</sup> in the Symkevi in combination with ivacaftor group, 0.20 (0.07) kg/m<sup>2</sup> in the ivacaftor group and 0.04 (0.07) kg/m<sup>2</sup> in the placebo group in Study 2. In Study 3, the change in BMI z-value in the Symkevi in combination with ivacaftor group was maintained.

#### **Children aged less than 12 years**

The European Medicines Agency has deferred the obligation to submit the results of studies with Symkevi in combination with ivacaftor in children aged less than 12 years in cystic fibrosis, as per PIP decision in the granted indication. See 4.2 for information on paediatric use.

## 5.2 Pharmacokinetic properties

The pharmacokinetics of tezacaftor and ivacaftor are similar between healthy adult volunteers and patients with CF. Following once daily dosing of tezacaftor and twice-daily dosing of ivacaftor in patients with CF, plasma concentrations of tezacaftor and ivacaftor reach steady-state within 8 days and within 3 to 5 days, respectively, after starting treatment. At steady-state, the accumulation ratio is approximately 2.3 for tezacaftor and 3.0 for ivacaftor. Exposures of tezacaftor (administered alone or in combination with ivacaftor) increase in an approximately dose-proportional manner with increasing doses from 10 mg to 300 mg once daily. Key pharmacokinetic parameters for tezacaftor and ivacaftor at steady-state are shown in Table 7.

	<b>Drug</b>	<b>C<sub>max</sub> (mcg/mL)</b>	<b>t<sub>1/2</sub> (h)</b>	<b>AUC<sub>0-24h</sub> or AUC<sub>0-12h</sub> (mcg·h/mL)*</b>
<b>Tezacaftor 100 mg once daily/ivacaftor 150 mg every 12 hours</b>	Tezacaftor	6.52 (1.83)	156 (52.7)	82.7 (23.3)
	Ivacaftor	1.28 (0.440)	9.3 (1.7)	10.9 (3.89)

\*AUC<sub>0-24h</sub> for tezacaftor and AUC<sub>0-12h</sub> for ivacaftor

### Absorption

After a single dose in healthy subjects in the fed state, tezacaftor was absorbed with a median (range) time to maximum concentration (t<sub>max</sub>) of approximately 4 hours (2 to 6 hours). The median (range) t<sub>max</sub> of ivacaftor was approximately 6 hours (3 to 10 hours) in the fed state. The AUC of tezacaftor did not change when given with fat-containing food relative to fasted conditions. The AUC of ivacaftor when given in combination with tezacaftor increased approximately 3-fold when given with fat-containing food; therefore, Symkevi and ivacaftor should be administered with fat-containing food.

### Distribution

Tezacaftor is approximately 99% bound to plasma proteins, primarily to albumin. Ivacaftor is approximately 99% bound to plasma proteins, primarily to alpha 1-acid glycoprotein and albumin. After oral administration of tezacaftor 100 mg once daily in combination with ivacaftor 150 mg every 12 hours in patients with CF in the fed state, the mean (±SD) for apparent volume of distribution of tezacaftor and ivacaftor was 271 (157) L and 206 (82.9) L, respectively. Neither tezacaftor nor ivacaftor partition preferentially into human red blood cells.

### Biotransformation

Tezacaftor is metabolized extensively in humans. *In vitro* data suggested that tezacaftor is metabolized mainly by CYP3A4 and CYP3A5. Following oral administration of a single dose of 100 mg <sup>14</sup>C-tezacaftor to healthy male subjects, M1-TEZ, M2-TEZ, and M5-TEZ were the three major circulating metabolites of tezacaftor in humans, contributing to 15%, 31%, and 33% of total radioactivity, respectively. Under steady-state, for each of the metabolites, exposure to M1-TEZ, M2-TEZ and M5-TEZ is approximately 1.5-fold higher than for tezacaftor. M1-TEZ has similar potency to that of tezacaftor and is considered pharmacologically active. M2-TEZ is much less pharmacologically active than tezacaftor or M1-TEZ, and M5-TEZ is not considered pharmacologically active. Another minor circulating metabolite, M3-TEZ, is formed by direct glucuronidation of tezacaftor.

Ivacaftor is also metabolized extensively in humans. *In vitro* and *in vivo* data indicate that ivacaftor is metabolized primarily by CYP3A4 and CYP3A5. M1-IVA and M6-IVA are the two major metabolites of ivacaftor in humans. M1-IVA has approximately one-sixth the potency of ivacaftor and is considered pharmacologically active. M6-IVA is not considered pharmacologically active.

The effect of the potentially reduced activity of CYP3A4 in patients carrying the CYP3A4\*22 variant on tezacaftor and ivacaftor exposures is not known.

### Elimination

After oral administration of tezacaftor 100 mg once daily in combination with ivacaftor 150 mg every 12 hours in patients with CF in the fed state, the mean ( $\pm$ SD) for apparent clearance values of tezacaftor and ivacaftor were 1.31 (0.41) and 15.7 (6.38) L/h, respectively. After steady-state dosing of tezacaftor in combination with ivacaftor in CF patients, the mean (SD) terminal half-lives of tezacaftor and ivacaftor were approximately 156 (52.7) and 9.3 (1.7) hours, respectively. The mean (SD) elimination half-lives for M1-TEZ, M2-TEZ and M5-TEZ were similar to that of the parent compound. The mean (SD) elimination half-lives for M1-IVA and M6-IVA were 11.3 (2.12) h and 14.4 (6.14) h, respectively.

Following oral administration of  $^{14}$ C-tezacaftor, the majority of the dose (72%) was excreted in the faeces (unchanged or as the M2-TEZ metabolite) and about 14% was recovered in urine (mostly as M2-TEZ metabolite), resulting in a mean overall recovery of 86% up to 21 days after the dose. Less than 1% of the administered dose was excreted in urine as unchanged tezacaftor, showing that renal excretion is not the major pathway of tezacaftor elimination in humans.

Following oral administration of ivacaftor alone, the majority of ivacaftor (87.8%) is eliminated in the faeces after metabolic conversion. There was negligible urinary excretion of ivacaftor as unchanged drug.

### Hepatic impairment

Following multiple doses of tezacaftor and ivacaftor for 10 days, subjects with moderately impaired hepatic function (Child-Pugh Class B, score 7 to 9) had an approximately 36% increase in AUC and a 10% increase in  $C_{max}$  for tezacaftor, and a 50% increase in ivacaftor AUC compared with healthy subjects matched for demographics. Based on these results, a modified regimen of Symkevi is recommended for patients with moderate hepatic impairment (see Table 2 in section 4.2).

The impact of severe hepatic impairment (Child-Pugh Class C, score 10 to 15) on the pharmacokinetics of tezacaftor and ivacaftor has not been studied. The magnitude of increase in exposure in these patients is unknown but is expected to be higher than that observed in patients with moderate hepatic impairment. The use of Symkevi in patients with severe hepatic impairment is therefore not recommended unless the benefits outweigh the risks (see Table 2 in section 4.2).

No dose adjustment is considered necessary for patients with mild hepatic impairment.

### Renal impairment

Symkevi in combination with ivacaftor has not been studied in patients with moderate or severe renal impairment (creatinine clearance  $\leq$ 30 mL/min) or in patients with end-stage renal disease. In a human pharmacokinetic study with tezacaftor alone, there was minimal elimination of tezacaftor and its metabolites in urine (only 13.7% of total radioactivity was recovered in the urine with 0.79% as unchanged medicinal product).

In a human pharmacokinetic study with ivacaftor alone, there was minimal elimination of ivacaftor and its metabolites in urine (only 6.6% of total radioactivity was recovered in the urine).

In population pharmacokinetic analysis, data from 147 patients on tezacaftor or tezacaftor in combination with ivacaftor in Phase 2/3 clinical studies indicated that mild renal impairment (estimated glomerular filtration rate, estimated by the modification of diet in renal disease method, 60 to  $\leq$ 89 mL/min/1.73 m<sup>2</sup>) did not affect the clearance of tezacaftor significantly. No dose adjustment is recommended for mild and moderate renal impairment. Caution is recommended when administering Symkevi in combination with ivacaftor to patients with severe renal impairment or end-stage renal disease.

### Gender

The pharmacokinetic parameters of tezacaftor and ivacaftor are similar in males and females.



### Race

Very limited PK data indicate comparable exposure to tezacaftor in white (n=652) and non-white (n=8) patients. Race had no clinically meaningful effect on the PK of ivacaftor in white (n=379) and non-white (n=29) patients based on a population PK analysis.

### Elderly

Clinical trials of Symkevi in combination with ivacaftor did not include patients over 75 years of age. The pharmacokinetic parameters of tezacaftor in combination with ivacaftor in the elderly patients (65 to 72 years) are comparable to those in younger adults.

### Paediatric Population

As presented in Table 8, the pharmacokinetic parameters of tezacaftor and ivacaftor in adolescents (12 to 17 years) are comparable to those in adults when given in combination. The pharmacokinetics of tezacaftor/ivacaftor in children below 12 years of age has not been investigated.

<b>Table 8. Mean (SD) tezacaftor and ivacaftor exposure by age group</b>			
<b>Age group</b>	<b>Dose</b>	<b>Tezacaftor Mean (SD) AUC<math>\tau</math>,ss (mcg·h/mL)</b>	<b>Ivacaftor Mean (SD) AUC<math>\tau</math>,ss (mcg·h/mL)</b>
Adolescents	TEZ 100 mg qd/IVA 150 mg q12h	97.1 (35.8)	11.4 (5.5)
Adults	TEZ 100 mg qd/IVA 150 mg q12h	85.9 (28.0)	11.4 (4.14)

## **5.3 Preclinical safety data**

### Tezacaftor

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential and toxicity to reproduction and development. Placental transfer of tezacaftor was observed in pregnant rats.

### Ivacaftor

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, and carcinogenic potential.

Ivacaftor was associated with slight decreases of the seminal vesicle weights, a decrease of overall fertility index and number of pregnancies in females mated with treated males and significant reductions in number of corpora lutea and implantation sites with subsequent reductions in the average litter size and average number of viable embryos per litter in treated females. The No Observed Adverse Effect Level (NOAEL) for fertility findings provides an exposure level of approximately 5 times the systemic exposure of ivacaftor and its metabolites when administered as tezacaftor/ivacaftor in adult humans at the maximum recommended human dose (MRHD).

In the pre- and post-natal study ivacaftor decreased survival and lactation indices and caused a reduction in pup body weights. The NOAEL for viability and growth in the offspring provides an exposure level of approximately 4 times the systemic exposure of ivacaftor and its metabolites when administered as tezacaftor/ivacaftor in adult humans at the MRHD. Placental transfer of ivacaftor was observed in pregnant rats and rabbits.

Findings of cataracts were observed in juvenile rats dosed from postnatal Day 7 through 35 at ivacaftor exposure levels of 0.25 times the MRHD based on systemic exposure of ivacaftor and its metabolites when administered as tezacaftor/ivacaftor. This finding has not been observed in fetuses derived from rat dams treated with ivacaftor on gestation Days 7 to 17, in rat pups exposed to ivacaftor through milk ingestion up to postnatal Day 20, in 7-week-old rats, nor in 3.5- to 5-month-old dogs treated with ivacaftor. The potential relevance of these findings in humans is unknown.

### Tezacaftor/ivacaftor

Combination repeat-dose toxicity studies in rats and dogs involving the co-administration of tezacaftor

and ivacaftor to assess the potential for additive and/or synergistic toxicity did not produce any unexpected toxicities or interactions.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Hypromellose acetate succinate  
Sodium laurilsulfate (E487)  
Hypromellose (E464)  
Microcrystalline cellulose (E460(i))  
Croscarmellose sodium (E468)  
Magnesium stearate (E470b)

#### Tablet film coat

Hypromellose (E464)  
Hydroxypropyl cellulose (E463)  
Titanium dioxide (E171)  
Talc (E553b)  
Iron oxide yellow (E172)

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

30 months

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and contents of container**

Blister consisting of PCTFE (polychlorotrifluoroethylene)/PVC (polyvinyl chloride) with a paper-backed aluminum foil lidding.

Pack size of 28 tablets (4 blister cards of 7 tablets each).

### **6.6 Special precautions for disposal**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Vertex Pharmaceuticals (Ireland) Limited  
28-32 Pembroke Street Upper  
Dublin 2, D02 EK84  
Ireland

**8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/18/1306/001

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 31 October 2018

**10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

## **ANNEX II**

- A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## **A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer responsible for batch release

Almac Pharma Services (Ireland) Limited  
Finnabair Industrial Estate  
Dundalk  
Co. Louth  
A91 P9KD  
Ireland

Almac Pharma Services Ltd  
20 Seagoe Industrial Estate  
Portadown  
Craigavon  
BT63 5UA  
United Kingdom

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- **Periodic safety update reports**

The requirements for submission of periodic safety update reports for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal. The marketing authorisation holder shall submit the first periodic safety update report for this product within 6 months following authorisation.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk Management Plan (RMP)**

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Symkevi 100 mg/150 mg film-coated tablets  
tezacaftor/ivacaftor

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each film-coated tablet contains 100 mg of tezacaftor and 150 mg of ivacaftor.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

28 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.  
Oral use

Take the tablets with fat-containing food.

Open  
Insert tab below to close

You may start taking Symkevi on any day of the week.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNINGS, IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**



**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Vertex Pharmaceuticals (Ireland) Limited  
28-32 Pembroke Street Upper  
Dublin 2, D02 EK84  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/18/1306/001

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Symkevi 100 mg, 150 mg

**17. UNIQUE IDENTIFIER - 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC:  
SN:  
NN:

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING**

**BLISTER CARD**

**1. NAME OF THE MEDICINAL PRODUCT**

Symkevi 100 mg/150 mg film-coated tablets  
tezacaftor/ivacaftor

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each film-coated tablet contains 100 mg of tezacaftor and 150 mg of ivacaftor.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

7 tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.  
Oral use

Take the tablets with fat-containing food.

Mon. Tue. Wed. Thu. Fri. Sat. Sun.

You may start taking Symkevi on any day of the week.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNINGS, IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Vertex Pharmaceuticals (Ireland) Limited  
28-32 Pembroke Street Upper  
Dublin 2, D02 EK84  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/18/1306/001

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER - 2D BARCODE**

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTERS**

**1. NAME OF THE MEDICINAL PRODUCT**

Symkevi 100 mg/150 mg tablets  
tezacaftor/ivacaftor

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Vertex

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

**B. PACKAGE LEAFLET**

## Package leaflet: Information for the patient

### Symkevi 100 mg/150 mg film-coated tablets tezacaftor/ivacaftor

- ▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

#### **Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Symkevi is and what it is used for
2. What you need to know before you take Symkevi
3. How to take Symkevi
4. Possible side effects
5. How to store Symkevi
6. Contents of the pack and other information

#### **1. What Symkevi is and what it is used for**

**Symkevi contains two active substances**, tezacaftor and ivacaftor. The medicine helps lung cells to work better in some patients with cystic fibrosis (CF). CF is an inherited condition in which the lungs and the digestive system can become clogged with thick, sticky mucus.

Symkevi works on a protein called CFTR (*cystic fibrosis transmembrane conductance regulator*), which is damaged in some people with CF (who have a mutation in the *CFTR* gene). Ivacaftor causes the protein to work better while tezacaftor increases the amount of protein at the cell surface. Symkevi is normally taken with ivacaftor, another medicine.

Symkevi taken with ivacaftor is for long-term treatment of patients aged 12 and over who have CF with certain genetic mutations that result in reduced amount and/or function of the CFTR protein.

Symkevi taken with ivacaftor helps your breathing by improving your lung function. You may also notice that you do not get ill as often and/or that it is easier to gain weight.

#### **2. What you need to know before you take Symkevi**

##### **Do not take Symkevi:**

- **If you are allergic** to tezacaftor, ivacaftor, or any of the other ingredients of this medicine (listed in section 6).

**Talk to your doctor**, without taking the tablets, if this applies to you.

### Warnings and precautions

- **Talk to your doctor if you have liver problems**, or have had them previously. Your doctor may need to adjust your dose.
- Your doctor will do some **blood tests to check your liver** before and during treatment with Symkevi, especially if your blood tests showed high liver enzymes in the past. Increased liver enzymes in the blood have been seen in patients with CF receiving Symkevi.

**Tell your doctor right away** if you have any symptoms of liver problems. These are listed in section 4.

- **Your doctor may do eye examinations** before and during treatment with Symkevi. Cloudiness of the eye lens (cataract) without any effect on vision has occurred in some children and adolescents receiving this treatment.
- **Talk to your doctor if you have kidney problems**, or you have previously had them.
- **Talk to your doctor** before starting treatment if you have received **an organ transplant**.

### Children under 12

Symkevi is not to be used in children under the age of 12 years. It is not known if Symkevi is safe and effective in children under 12.

### Other medicines and Symkevi

**Tell your doctor or pharmacist** if you are taking, have recently taken, or might take any other medicines. Some medicines can affect how Symkevi works or may make side effects more likely. In particular, tell your doctor if you take any of the medicines listed below. Your doctor may change the dose of one of the medicines if you take any of these.

- **Antifungal medicines** (used for the treatment of fungal infections). These include ketoconazole, itraconazole, posaconazole, voriconazole and fluconazole.
- **Antibiotic medicines** (used for the treatment of bacterial infections). These include telithromycin, clarithromycin, erythromycin, rifampicin and rifabutin.
- **Anticonvulsant medicines** (used for the treatment of epilepsy and epileptic seizures or fits). These include phenobarbital, carbamazepine and phenytoin.
- **Herbal medicines**. These include St. John's wort (*Hypericum perforatum*).
- **Immunosuppressants** (used after an organ transplantation). These include ciclosporin, tacrolimus, sirolimus and everolimus.
- **Cardiac glycosides** (used for the treatment of some heart conditions). These include digoxin.
- **Anticoagulant medicines** (used to prevent blood clots). These include warfarin.
- **Medicines for diabetes**. These include glimepiride and glipizide.

### Symkevi with food and drink

Avoid food or drinks containing grapefruit or Seville oranges (marmalade oranges) during treatment as these may increase the side effects of Symkevi by increasing the amount of Symkevi in your body.

### Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

- **Pregnancy:** It may be better to avoid using this medicine during pregnancy. Your doctor will help you decide what is best for you and your child.
- **Breast-feeding:** It is not known if tezacaftor or ivacaftor pass into breast milk. Your doctor will consider the benefit of breast-feeding for the child and the benefit of treatment for you to help you decide whether to stop breast-feeding or to stop treatment.

### **Driving and using machines**

Symkevi in combination with ivacaftor can make you dizzy. If you feel dizzy, do not drive, cycle or use machines unless you are not affected.

### **Important information about the contents of Symkevi**

**This medicine contains** less than 1 mmol sodium (23 mg) per dose, that is to say essentially ‘sodium-free’.

## **3. How to take Symkevi**

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

### **The recommended dose:**

Symkevi is usually taken with ivacaftor.

- **In the morning, take one yellow Symkevi tablet.** This is stamped with “V100”.
- **In the evening, take one ivacaftor 150 mg tablet.**

Take the tablets about 12 hours apart.

**Take both Symkevi and ivacaftor tablets with food that contains fat.** Meals or snacks that contain fat include those prepared with butter or oils or those containing eggs. Other fat-containing foods are:

- Cheese, whole milk, whole milk dairy products, yogurt, chocolate
- Meats, oily fish
- Avocados, hummus, soy-based products (tofu)
- Nuts, fat-containing nutritional bars or drinks

The tablets are for oral use.

**Swallow the tablet whole.** Do not chew, crush or break the tablets before swallowing.

You must keep using all your other medicines, unless your doctor tells you to stop.

**If you have liver problems**, either moderate or severe, your doctor may need to reduce the dose of your tablets, because your liver will not process the medicine as fast as usual.

### **If you take more Symkevi than you should**

**Contact your doctor or pharmacist** for advice. If possible, take your medicine and this leaflet with you. You may get side effects, including those mentioned in section 4 below.

### **If you forget to take Symkevi**

- If you forget to take either your morning Symkevi or evening ivacaftor tablet, and you remember **within 6 hours** of the scheduled time you should have taken the tablet, take the forgotten tablet at once.
- If **more than 6 hours** have passed, do not take the forgotten tablet. Just wait, and take your next tablet at the usual time.
- **Do not** take 2 tablets to make up for a missed dose.

### **If you stop taking Symkevi**

Your doctor will tell you how long you need to keep using Symkevi. It is important to take this medicine regularly. Do not make changes unless your doctor tells you.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.



#### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

##### **Possible signs of liver problems**

Increased liver enzymes in the blood are very common in patients with CF. These may be a sign of liver problems:

- Pain or discomfort in the upper right area of the stomach (abdominal) area
- Yellowing of the skin or the white part of the eyes
- Loss of appetite
- Nausea or vomiting
- Dark urine

**Tell your doctor straight away** if you have any of these symptoms.

##### **Side effects seen with Symkevi in combination with ivacaftor:**

###### **Very common side effects**

(may affect more than 1 in 10 people)

- Headache
- Common cold

###### **Common side effects**

(may affect up to 1 in 10 people)

- Feeling sick (nausea)
- Blocked nose (sinus congestion)
- Dizziness

##### **Side effects seen with ivacaftor:**

###### **Very common side effects**

- Upper respiratory tract infection (the common cold), including sore throat and nasal congestion
- Headache
- Dizziness
- Stomach (abdominal) ache
- Diarrhoea
- Increased liver enzymes in the blood
- Rash
- Changes in the type of bacteria in mucus

###### **Common side effects**

- Runny nose
- Ear pain, ear discomfort
- Ringing in the ears
- Redness inside the ear
- Inner ear disorder (feeling dizzy or spinning)
- Sinus congestion
- Redness in the throat
- Breast mass

###### **Uncommon side effects**

(may affect up to 1 in 100 people)

- Ear congestion
- Breast inflammation
- Enlargement of the breast in males
- Nipple changes or pain

### **Additional side effects in children and adolescents**

Side effects in children and adolescents are similar to those observed in adults.

### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Symkevi**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date that is stated on the package after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

## **6. Contents of the pack and other information**

### **What Symkevi contains**

- The active substances are tezacaftor and ivacaftor. Each film-coated tablet contains 100 mg of tezacaftor and 150 mg of ivacaftor.
- The other ingredients are:
  - Tablet core: Hypromellose acetate succinate, sodium laurilsulfate (E487), hypromellose (E464), microcrystalline cellulose (E460(i)), croscarmellose sodium (E468), and magnesium stearate (E470b).
  - Tablet film coating: Hypromellose (E464), hydroxypropyl cellulose (E463), titanium dioxide (E171), talc (E553b), and iron oxide yellow (E172).

See the end of section 2 for important information about the contents of Symkevi.

### **What Symkevi looks like and contents of the pack**

Symkevi 100 mg tezacaftor/150 mg ivacaftor film-coated tablets are yellow, oval-shaped tablets stamped with “V100” on one side and plain on the other.

Symkevi is available in the following pack size:

Pack size of 28 tablets (4 blister cards, each with 7 tablets).

### **Marketing Authorisation Holder and Manufacturer**

Marketing Authorisation Holder:

Vertex Pharmaceuticals (Ireland) Limited

28-32 Pembroke Street Upper

Dublin 2, D02 EK84

Ireland

Tel: +353 (0)1 761 7299

Manufacturer:  
Almac Pharma Services (Ireland) Limited  
Finnabair Industrial Estate  
Dundalk  
Co. Louth  
A91 P9KD  
Ireland

Almac Pharma Services Limited  
Seagoe Industrial Estate  
Craigavon  
County Armagh  
BT63 5UA  
United Kingdom

**This leaflet was last revised in**

**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency website:  
<http://www.ema.europa.eu>. There are also links to other websites about rare diseases and treatments.