

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

1. NAME OF THE MEDICINAL PRODUCT

Yesafili 40 mg/mL solution for injection in a vial

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

1 mL solution for injection contains 40 mg aflibercept*.

One vial contains an extractable volume of at least 0.1 mL, equivalent to at least 4 mg aflibercept. This provides a usable amount to deliver a single dose of 0.05 mL containing 2 mg aflibercept.

*Fusion protein consisting of portions of human VEGF (vascular endothelial growth factor) receptors 1 and 2 extracellular domains fused to the Fc portion of human IgG1 and produced in Chinese hamster ovary (CHO) K1 cells by recombinant DNA technology.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Solution for injection (injection).

Clear, colourless to pale yellow and iso-osmotic solution.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Yesafili is indicated for adults for the treatment of

- neovascular (wet) age-related macular degeneration (AMD) (see section 5.1),
- visual impairment due to macular oedema secondary to retinal vein occlusion (branch RVO or central RVO) (see section 5.1),
- visual impairment due to diabetic macular oedema (DME) (see section 5.1),
- visual impairment due to myopic choroidal neovascularisation (myopic CNV) (see section 5.1).

4.2 Posology and method of administration

Yesafili is for intravitreal injection only.

Yesafili must only be administered by a qualified physician experienced in administering intravitreal injections.

Posology

wet AMD

The recommended dose for Yesafili is 2 mg aflibercept, equivalent to 0.05 mL.

Yesafili treatment is initiated with one injection per month for three consecutive doses. The treatment interval is then extended to two months.

Based on the physician's judgement of visual and/or anatomic outcomes, the treatment interval may be maintained at two months or further extended using a treat-and-extend dosing regimen, where injection intervals are increased in 2- or 4-weekly increments to maintain stable visual and/or anatomic outcomes.

If visual and/or anatomic outcomes deteriorate, the treatment interval should be shortened accordingly.

There is no requirement for monitoring between injections. Based on the physician's judgement the schedule of monitoring visits may be more frequent than the injection visits. Treatment intervals greater than four months or shorter than 4 weeks between injections have not been studied (see section 5.1).

Macular oedema secondary to RVO (branch RVO or central RVO)

The recommended dose for Yesafili is 2 mg aflibercept equivalent to 0.05 mL.

After the initial injection, treatment is given monthly. The interval between two doses should not be shorter than one month.

If visual and anatomic outcomes indicate that the patient is not benefiting from continued treatment, Yesafili should be discontinued.

Monthly treatment continues until maximum visual acuity is achieved and/or there are no signs of disease activity. Three or more consecutive, monthly injections may be needed.

Treatment may then be continued with a treat-and-extend regimen with gradually increased treatment intervals to maintain stable visual and/or anatomic outcomes, however there are insufficient data to conclude on the length of these intervals. If visual and/or anatomic outcomes deteriorate, the treatment interval should be shortened accordingly.

The monitoring and treatment schedule should be determined by the treating physician based on the individual patient's response.

Monitoring for disease activity may include clinical examination, functional testing or imaging techniques (e.g. optical coherence tomography or fluorescein angiography).

Diabetic macular oedema

The recommended dose for Yesafili is 2 mg aflibercept equivalent to 0.05 mL.

Yesafili treatment is initiated with one injection per month for five consecutive doses, followed by one injection every two months.

Based on the physician's judgement of visual and/or anatomic outcomes, the treatment interval may be maintained at 2 months or individualized, such as with a treat-and-extend dosing regimen, where the treatment intervals are usually increased by 2-week increments to maintain stable visual and/or anatomic outcomes. There are limited data for treatment intervals longer than 4 months. If visual and/or anatomic outcomes deteriorate, the treatment interval should be shortened accordingly.

Treatment intervals shorter than 4 weeks between injections have not been studied (see section 5.1).

The schedule for monitoring should be determined by the treating physician.

If visual and anatomic outcomes indicate that the patient is not benefiting from continued treatment, Yesafili should be discontinued.

Myopic choroidal neovascularisation

The recommended dose for Yesafili is a single intravitreal injection of 2 mg aflibercept equivalent to 0.05 mL.

Additional doses may be administered if visual and/or anatomic outcomes indicate that the disease persists. Recurrences should be treated as a new manifestation of the disease.

The schedule for monitoring should be determined by the treating physician.

The interval between two doses should not be shorter than one month.

Special populations

Hepatic and/or renal impairment

No specific studies in patients with hepatic and/or renal impairment have been conducted with aflibercept.

Available data do not suggest a need for a dose adjustment with Yesafili in these patients (see section 5.2).

Elderly population

No special considerations are needed. There is limited experience in patients older than 75 years with DME.

Paediatric population

The safety and efficacy of aflibercept have not been established in children and adolescents. There is no relevant use of aflibercept in the paediatric population for the indications of wet AMD, CRVO, BRVO, DME and myopic CNV.

Method of administration

Intravitreal injections must be carried out according to medical standards and applicable guidelines by a qualified physician experienced in administering intravitreal injections. In general, adequate anaesthesia and asepsis, including topical broad-spectrum microbicide (e.g. povidone iodine applied to the periocular skin, eyelid and ocular surface), have to be ensured. Surgical hand disinfection, sterile gloves, a sterile drape, and a sterile eyelid speculum (or equivalent) are recommended.

The injection needle should be inserted 3.5 - 4.0 mm posterior to the limbus into the vitreous cavity, avoiding the horizontal meridian and aiming towards the centre of the globe. The injection volume of 0.05 mL is then delivered; a different scleral site should be used for subsequent injections.

Immediately following the intravitreal injection, patients should be monitored for elevation in intraocular pressure. Appropriate monitoring may consist of a check for perfusion of the optic nerve head or tonometry. If required, sterile equipment for paracentesis should be available.

Following intravitreal injection patients should be instructed to report any symptoms suggestive of endophthalmitis (e.g. eye pain, redness of the eye, photophobia, blurring of vision) without delay.

Each vial should only be used for the treatment of a single eye. Extraction of multiple doses from a single vial may increase the risk of contamination and subsequent infection.

The vial contains more than the recommended dose of 2 mg aflibercept (equivalent to 0.05 mL solution for injection). The extractable volume of the vial is the amount that can be withdrawn from the vial and is not to be used in total. For the Yesafili vial, the extractable volume is at least 0.1 mL. **(The excess volume must be expelled before injecting the recommended dose** (see section 6.6).

Injecting the entire volume of the vial could result in overdose. To expel the air bubbles along with excess medicinal product, slowly depress the plunger so that the flat plunger edge aligns with the line that marks 0.05 mL on the syringe (equivalent to 0.05 mL i.e. 2 mg aflibercept) (see sections 4.9 and

6.6).

After injection any unused product must be discarded.

For handling of the medicinal product before administration, see section 6.6.

4.3 Contraindications

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- Active or suspected ocular or periocular infection.
- Active severe intraocular inflammation.

4.4 Special warnings and precautions for use

Traceability

In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded.

Intravitreal injection-related reactions

Intravitreal injections, including those with aflibercept, have been associated with endophthalmitis, intraocular inflammation, rhegmatogenous retinal detachment, retinal tear and iatrogenic traumatic cataract (see section 4.8). Proper aseptic injection techniques must always be used when administering aflibercept. In addition, patients should be monitored during the week following the injection to permit early treatment if an infection occurs. Patients should be instructed to report any symptoms suggestive of endophthalmitis or any of the above mentioned events without delay.

The vial contains more than the recommended dose of 2 mg aflibercept (equivalent to 0.05 mL). The excess volume must be discarded prior to administration (see sections 4.2 and 6.6.).

Increases in intraocular pressure have been seen within 60 minutes of intravitreal injection, including those with aflibercept (see section 4.8). Special precaution is needed in patients with poorly controlled glaucoma (do not inject Yesafili while the intraocular pressure is ≥ 30 mmHg). In all cases, both the intraocular pressure and the perfusion of the optic nerve head must therefore be monitored and managed appropriately.

Immunogenicity

As this is a therapeutic protein, there is a potential for immunogenicity with Yesafili (see section 4.8). Patients should be instructed to report any signs or symptoms of intraocular inflammation, e.g. pain, photophobia, or redness, which may be a clinical sign attributable to hypersensitivity.

Systemic effects

Systemic adverse events including non-ocular haemorrhages and arterial thromboembolic events have been reported following intravitreal injection of VEGF inhibitors and there is a theoretical risk that these may relate to VEGF inhibition. There are limited data on safety in the treatment of patients with CRVO, BRVO, DME or myopic CNV with a history of stroke or transient ischaemic attacks or myocardial infarction within the last 6 months. Caution should be exercised when treating such patients.

Other

As with other intravitreal anti-VEGF treatments for AMD, CRVO, BRVO, DME and myopic CNV the following also applies:

- The safety and efficacy of aflibercept therapy administered to both eyes concurrently have not been systematically studied (see section 5.1). If bilateral treatment is performed at the same

time this could lead to an increased systemic exposure, which could increase the risk of systemic adverse events.

- Concomitant use of other anti-VEGF (vascular endothelial growth factor). There is no data available on the concomitant use of aflibercept with other anti-VEGF medicinal products (systemic or ocular).
- Risk factors associated with the development of a retinal pigment epithelial tear after anti-VEGF therapy for wet AMD, include a large and/or high pigment epithelial retinal detachment.
When initiating aflibercept therapy, caution should be used in patients with these risk factors for retinal pigment epithelial tears.
- Treatment should be withheld in patients with rhegmatogenous retinal detachment or stage 3 or 4 macular holes.
- In the event of a retinal break the dose should be withheld and treatment should not be resumed until the break is adequately repaired.
- The dose should be withheld and treatment should not be resumed earlier than the next scheduled treatment in the event of:
 - a decrease in best-corrected visual acuity (BCVA) of ≥ 30 letters compared with the last assessment of visual acuity;
 - a subretinal haemorrhage involving the centre of the fovea, or, if the size of the haemorrhage is $\geq 50\%$, of the total lesion area.
- The dose should be withheld within the previous or next 28 days in the event of a performed or planned intraocular surgery.
- Aflibercept should not be used in pregnancy unless the potential benefit outweighs the potential risk to the foetus (see section 4.6).
- Women of childbearing potential have to use effective contraception during treatment and for at least 3 months after the last intravitreal injection of aflibercept (see section 4.6).
- There is limited experience with treatment of patients with ischaemic CRVO and BRVO. In patients presenting with clinical signs of irreversible ischaemic visual function loss, the treatment is not recommended.

Populations with limited data

There is only limited experience in the treatment of subjects with DME due to type I diabetes or in diabetic patients with an HbA1c over 12% or with proliferative diabetic retinopathy.

Aflibercept has not been studied in patients with active systemic infections or in patients with concurrent eye conditions such as retinal detachment or macular hole. There is also no experience of treatment with aflibercept in diabetic patients with uncontrolled hypertension. This lack of information should be considered by the physician when treating such patients.

In myopic CNV there is no experience with aflibercept in the treatment of non-Asian patients, patients who have previously undergone treatment for myopic CNV, and patients with extrafoveal lesions.

Information about excipients

This medicinal product contains less than 1 mmol sodium (23 mg) per dosage unit, that is to say essentially 'sodium-free'

4.5 Interaction with other medicinal products and other forms of interaction

No interaction studies have been performed.

Adjunctive use of verteporfin photodynamic therapy (PDT) and aflibercept has not been studied, therefore, a safety profile is not established.

4.6 Fertility, pregnancy and lactation

Women of childbearing potential

Women of childbearing potential have to use effective contraception during treatment and for at least 3

months after the last intravitreal injection of aflibercept (see section 4.4).

Pregnancy

There are no data on the use of aflibercept in pregnant women. Studies in animals have shown embryo-foetal toxicity (see section 5.3).

Although the systemic exposure after ocular administration is very low, aflibercept should not be used during pregnancy unless the potential benefit outweighs the potential risk to the foetus.

Breast-feeding

It is unknown whether aflibercept is excreted in human milk. A risk to the breast-fed child cannot be excluded.

Aflibercept is not recommended during breast-feeding. A decision must be made whether to discontinue breast-feeding or to abstain from aflibercept therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

Fertility

Results from animal studies with high systemic exposure indicate that aflibercept can impair male and female fertility (see section 5.3). Such effects are not expected after ocular administration with very low systemic exposure

4.7 Effects on ability to drive and use machines

Injection with aflibercept has a minor influence on the ability to drive and use machines due to possible temporary visual disturbances associated either with the injection or the eye examination. Patients should not drive or use machines until their visual function has recovered sufficiently.

4.8 Undesirable effects

Summary of the safety profile

A total of 3 102 patients constituted the safety population in the eight phase III studies. Among those, 2 501 patients were treated with the recommended dose of 2 mg.

Serious ocular adverse reactions in the study eye related to the injection procedure have occurred in less than 1 in 1 900 intravitreal injections with aflibercept and included blindness, endophthalmitis, retinal detachment, cataract traumatic, cataract, vitreous haemorrhage, vitreous detachment, and intraocular pressure increased (see section 4.4).

The most frequently observed adverse reactions (in at least 5% of patients treated with aflibercept) were conjunctival haemorrhage (25%), retinal haemorrhage (11%), visual acuity reduced (11%), eye pain (10%), cataract (8%), intraocular pressure increased (8%), vitreous detachment (7%), and vitreous floaters (7%).

Tabulated list of adverse reactions

The safety data described below include all adverse reactions from the eight phase III studies in the indications wet AMD, CRVO, BRVO, DME and myopic CNV with a reasonable possibility of causality to the injection procedure or medicinal product.

The adverse reactions are listed by system organ class and frequency using the following convention: Very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1\ 000$ to $< 1/100$), rare ($\geq 1/10\ 000$ to $< 1/1\ 000$)

Within each frequency grouping, adverse drug reactions are presented in order of decreasing seriousness.

Table 1: All treatment-emergent adverse drug reactions reported in patients in phase III studies (pooled data of the phase III studies for the indications wet AMD, CRVO, BRVO, DME and myopic CNV) or during post-marketing surveillance

System Organ Class	Very common	Common	Uncommon	Rare
Immune system disorders			Hypersensitivity***	
Eye disorders	Visual acuity reduced, Retinal haemorrhage, Conjunctival haemorrhage, Eye pain	Retinal pigment epithelial tear*, Detachment of the retinal pigment epithelium, Retinal degeneration, Vitreous haemorrhage, Cataract, Cataract cortical, Cataract nuclear, Cataract subcapsular, Corneal erosion, Corneal abrasion, Intraocular pressure increased, Vision blurred, Vitreous floaters, Vitreous detachment, Injection site pain, Foreign body sensation in eyes, Lacrimation increased, Eyelid oedema, Injection site haemorrhage, Punctate keratitis, Conjunctival hyperaemia, Ocular hyperaemia	Endophthalmitis** , Retinal detachment, Retinal tear, Iritis, Uveitis, Iridocyclitis, Lenticular opacities, Corneal epithelium defect, Injection site irritation, Abnormal sensation in eye, Eyelid irritation, Anterior chamber flare, Corneal oedema	Blindness, Cataract traumatic, Vitritis, Hypopyon

* Conditions known to be associated with wet AMD. Observed in the wet AMD studies only.

** Culture positive and culture negative endophthalmitis

*** During the post-marketing period, reports of hypersensitivity included rash, pruritus, urticaria, and isolated cases of severe anaphylactic/anaphylactoid reactions.

Description of selected adverse reactions

In the wet AMD phase III studies, there was an increased incidence of conjunctival haemorrhage in patients receiving anti-thrombotic agents. This increased incidence was comparable between patients treated with ranibizumab and aflibercept.

Arterial thromboembolic events (ATEs) are adverse events potentially related to systemic VEGF inhibition. There is a theoretical risk of arterial thromboembolic events, including stroke and myocardial infarction, following intravitreal use of VEGF inhibitors.

A low incidence rate of arterial thromboembolic events was observed in the aflibercept clinical trials in patients with AMD, DME, RVO and myopic CNV. Across indications no notable difference between the groups treated with aflibercept and the respective comparator groups were observed.

As with all therapeutic proteins, there is a potential for immunogenicity with Yesafili.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via **the national reporting system listed in [Appendix V](#)**.

4.9 Overdose

In clinical trials, doses of up to 4 mg in monthly intervals have been used and isolated cases of overdoses with 8 mg occurred.

Overdosing with increased injection volume may increase intraocular pressure. Therefore, in case of overdose, intraocular pressure should be monitored and if deemed necessary by the treating physician, adequate treatment should be initiated (see section 6.6).

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Ophthalmologicals / Antineovascularisation agents
ATC code: S01LA05

Yesafili is a biosimilar medicinal product. Detailed information is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

Aflibercept is a recombinant fusion protein consisting of portions of human VEGF receptor 1 and 2 extracellular domains fused to the Fc portion of human IgG1.

Aflibercept is produced in Chinese hamster ovary (CHO) K1 cells by recombinant DNA technology.

Aflibercept acts as a soluble decoy receptor that binds VEGF-A and PlGF with higher affinity than their natural receptors, and thereby can inhibit the binding and activation of these cognate VEGF receptors.

Mechanism of action

Vascular endothelial growth factor-A (VEGF-A) and placental growth factor (PlGF) are members of the VEGF family of angiogenic factors that can act as potent mitogenic, chemotactic, and vascular permeability factors for endothelial cells. VEGF acts via two receptor tyrosine kinases; VEGFR-1 and

VEGFR-2, present on the surface of endothelial cells. PlGF binds only to VEGFR-1, which is also present on the surface of leucocytes. Excessive activation of these receptors by VEGF-A can result in pathological neovascularisation and excessive vascular permeability. PlGF can synergize with VEGF-A in these processes and is also known to promote leucocyte infiltration and vascular inflammation.

Pharmacodynamic effects

wet AMD

Wet AMD is characterised by pathological choroidal neovascularisation (CNV). Leakage of blood and fluid from CNV may cause retinal thickening or oedema and/or sub-/intra-retinal haemorrhage, resulting in loss of visual acuity.

In patients treated with aflibercept (one injection per month for three consecutive months, followed by one injection every 2 months), central retinal thickness [CRT] decreased soon after treatment initiation, and the mean CNV lesion size was reduced, consistent with the results seen with ranibizumab 0.5 mg every month.

In the VIEW1 study there were mean decreases in CRT on optical coherence tomography (OCT) (- 130 and - 129 microns at week 52 for the aflibercept 2 mg every two months and ranibizumab 0.5 mg every month study groups, respectively). Also at the 52 -week time point, in the VIEW2 study there were mean decreases in CRT on OCT (- 149 and - 139 microns for the aflibercept 2 mg every two months and ranibizumab 0.5 mg every month study groups, respectively). The reduction of CNV size and reduction in CRT were generally maintained in the second year of the studies.

The ALTAIR study was conducted in Japanese patients with treatment naïve wet AMD, showing similar outcomes to the VIEW studies using 3 initial monthly aflibercept 2 mg injections, followed by one injection after a further 2 months, and then continued with a treat-and-extend regimen with variable treatment intervals (2 -week or 4 -week adjustments) up to a maximum 16 week interval according to pre-specified criteria. At week 52, there were mean decreases in central retinal thickness (CRT) on OCT of - 134.4 and - 126.1 microns for the 2 -week adjustment group and the 4 -week adjustment group, respectively. The proportion of patients without fluid on OCT at week 52 was 68.3% and 69.1% in the 2 - and 4 -week adjustment groups, respectively. The reduction in CRT was generally maintained in both treatment arms in the second year of the ALTAIR study.

The ARIES study was designed to explore the non-inferiority of an aflibercept 2 mg treat-and-extend dosing regimen initiated immediately after administration of 3 initial monthly injections and one additional injection after 2 months vs. a treat-and-extend dosing regimen initiated after one year of treatment. For patients requiring a more frequent than Q8 dosing at least once over the course of the study, CRT remained higher, but the mean decrease in CRT from baseline to week 104 was - 160.4 microns, similar to the patients treated at Q8 or less frequent intervals.

Macular oedema secondary to CRVO and BRVO

In CRVO and BRVO, retinal ischaemia occurs and signals the release of VEGF which in turn destabilises the tight junctions, and promotes endothelial cell proliferation. Up-regulation of VEGF is associated with the breakdown of the blood retina barrier, increased vascular permeability, retinal oedema, and neovascularisation complications.

In patients treated with 6 consecutive monthly injections of aflibercept 2 mg, there was a consistent, rapid and robust morphologic response (as measured by improvements in mean CRT) observed. At week 24, the reduction in CRT was statistically superior versus control in all three studies (COPERNICUS in CRVO: - 457 vs. - 145 microns; GALILEO in CRVO: - 449 vs. - 169 microns; VIBRANT in BRVO: - 280 vs. - 128 microns). This decrease from baseline in CRT was maintained to the end of each study, week 100 in COPERNICUS, week 76 in GALILEO, and week 52 in VIBRANT.

Diabetic macular oedema

Diabetic macular oedema is a consequence of diabetic retinopathy and is characterised by increased vasopermeability and damage to the retinal capillaries which may result in loss of visual acuity.

In patients treated with aflibercept, the majority of whom were classified as having Type II diabetes, a rapid and robust response in morphology (CRT, DRSS level) was observed.

In the VIVID^{DME} and the VISTA^{DME} studies, a statistically significant greater mean decrease in CRT from baseline to week 52 was observed in patients treated with aflibercept than with the laser control, - 192.4 and - 183.1 microns for the 2Q8 aflibercept groups and - 66.2 and - 73.3 microns for the control groups, respectively. At week 100 the decrease was maintained with - 195.8 and - 191.1 microns for the 2Q8 aflibercept groups and - 85.7 and - 83.9 microns for the control groups, in the VIVID^{DME} and VISTA^{DME} studies, respectively.

A ≥ 2 step improvement in DRSS was assessed in a pre-specified manner in VIVID^{DME} and VISTA^{DME}. The DRSS score was gradable in 73.7% of the patients in VIVID^{DME} and 98.3% of the patients in VISTA^{DME}. At week 52, 27.7% and 29.1% of the aflibercept 2Q8 groups, and 7.5% and 14.3% of the control groups experienced a ≥ 2 step improvement in the DRSS. At week 100, the respective percentages were 32.6% and 37.1% of the aflibercept 2Q8 groups and 8.2% and 15.6% of the control groups.

The VIOLET study compared three different dosing regimens of aflibercept 2 mg for treatment of DME after at least one year of treatment at fixed intervals, where treatment was initiated with 5 consecutive monthly doses followed by dosing every 2 months. At week 52 and week 100 of the study, i.e. second and third year of treatment, the mean changes in CRT were clinically similar for treat-and-extend (2T&E), pro re nata (2PRN) and 2Q8, respectively, - 2.1, 2.2 and - 18.8 microns at week 52, and 2.3, - 13.9 and - 15.5 microns at week 100.

Myopic choroidal neovascularisation

Myopic choroidal neovascularisation (myopic CNV) is a frequent cause of vision loss in adults with pathologic myopia. It develops as a wound healing mechanism consequent to Bruch's membrane ruptures and represents the most vision-threatening event in pathologic myopia.

In patients treated with aflibercept in the MYRROR study (one injection given at start of therapy, with additional injections given in case of disease persistence or recurrence), CRT decreased soon after treatment initiation favouring aflibercept at week 24 (- 79 microns and - 4 microns for the aflibercept 2 mg treatment group and the control group, respectively), which was maintained through week 48. In addition, the mean CNV lesion size decreased.

Clinical efficacy and safety

wet AMD

The safety and efficacy of aflibercept were assessed in two randomised, multi-centre, double-masked, active-controlled studies in patients with wet AMD (VIEW1 and VIEW2) with a total of 2 412 patients treated and evaluable for efficacy (1 817 with aflibercept). Patient ages ranged from 49 to 99 years with a mean of 76 years. In these clinical studies, approximately 89% (1 616/1 817) of the patients randomised to treatment with aflibercept were 65 years of age or older, and approximately 63% (1 139/1 817) were 75 years of age or older. In each study, patients were randomly assigned in a 1:1:1:1 ratio to 1 of 4 dosing regimens:

- 1) Aflibercept administered at 2 mg every 8 weeks following 3 initial monthly doses (aflibercept 2Q8);
- 2) Aflibercept administered at 2 mg every 4 weeks (aflibercept 2Q4);
- 3) Aflibercept administered at 0.5 mg every 4 weeks (aflibercept 0.5Q4); and
- 4) ranibizumab administered at 0.5 mg every 4 weeks (ranibizumab 0.5Q4).

In the second year of the studies, patients continued to receive the initially randomised dose but on a modified dosing schedule guided by assessment of visual and anatomic outcomes with a protocol-defined maximum dosing interval of 12 weeks.

In both studies, the primary efficacy endpoint was the proportion of patients in the Per Protocol Set who maintained vision, i.e. losing fewer than 15 letters of visual acuity at week 52 from baseline.

In the VIEW1 study, at week 52, 95.1% of patients in the aflibercept 2Q8 group maintained vision

compared to 94.4% patients in the ranibizumab 0.5Q4 group. In the VIEW2 study, at week 52, 95.6% of patients in the aflibercept 2Q8 group maintained vision compared to 94.4% patients in the ranibizumab 0.5Q4 group. In both studies aflibercept was shown to be non-inferior and clinically equivalent to the ranibizumab 0.5Q4 group.

Detailed results from the combined analysis of both studies are shown in Table 2 and Figure 1 below

Table 2: Efficacy outcomes at week 52 (primary analysis) and week 96; combined data from the VIEW1 and VIEW2 studies^{B)}

Efficacy Outcome	Aflibercept 2Q8 ^{E)} (aflibercept 2 mg every 8 weeks following 3 initial monthly doses) (N=607)		Ranibizumab 0.5Q4 (ranibizumab 0.5 mg every 4 weeks) (N=595)	
	Week 52	Week 96	Week 52	Week 96
Mean number of injections from baseline	7.6	11.2	12.3	16.5
Mean number of injections from week 52 to 96		4.2		4.7
Proportion of patients with < 15 letters loss from baseline (PPS ^{A)})	95.33% ^{B)}	92.42%	94.42% ^{B)}	91.60%
Difference ^{C)} (95 % CI) ^{D)}	0.9% (- 1.7, 3.5) ^{F)}	0.8% (- 2.3, 3.8) ^{F)}		
Mean change in BCVA as measured by ETDRS ^{A)} letter score from baseline	8.40	7.62	8.74	7.89
Difference in LS ^{A)} mean change (ETDRS letters) ^{C)} (95 % CI) ^{D)}	- 0.32 (- 1.87, 1.23)	- 0.25 (- 1.98, 1.49)		
Proportion of patients with ≥ 15 letters gain from baseline	30.97%	33.44%	32.44%	31.60%
Difference ^{C)} (95 % CI) ^{D)}	- 1.5% (- 6.8, 3.8)	1.8% (- 3.5, 7.1)		

^{A)} BCVA: Best Corrected Visual Acuity

ETDRS: Early Treatment Diabetic Retinopathy Study

LS: Least square means derived from ANCOVA

PPS: Per Protocol Set

^{B)} Full Analysis Set (FAS), Last Observation Carried Forward (LOCF) for all analyses except proportion of patients with maintained visual acuity at week 52 which is PPS


^{C)} The difference is the value of the aflibercept group minus the value of the ranibizumab group. A positive value favours aflibercept

^{D)} Confidence interval (CI) calculated by normal approximation

^{E)} After treatment initiation with three monthly doses

^{F)} A confidence interval entirely above -10% indicates a non-inferiority of aflibercept to ranibizumab

Figure 1: Mean change in visual acuity from baseline to week 96 for the combined data from the view 1 and view 2 studies



In combined data analysis of VIEW1 and VIEW2, aflibercept demonstrated clinically meaningful changes from baseline in pre-specified secondary efficacy endpoint National Eye Institute Visual Function Questionnaire (NEI VFQ-25) without clinically meaningful differences to ranibizumab. The magnitude of these changes was similar to that seen in published studies, which corresponded to a 15 - letter gain in Best Corrected Visual Acuity (BCVA).

In the second year of the studies, efficacy was generally maintained through the last assessment at week 96, and 2-4% of patients required all injections on a monthly basis, and a third of patients required at least one injection with a treatment interval of only one month.

Decreases in mean CNV area were evident in all dose groups in both studies.

Efficacy results in all evaluable subgroups (e.g., age, gender, race, baseline visual acuity, lesion type, lesion size) in each study and in the combined analysis were consistent with the results in the overall populations.

ALTAIR was a 96 -week multicentre, randomised, open-label study in 247 Japanese patients with treatment naïve wet AMD, designed to assess the efficacy and safety of aflibercept following two different adjustment intervals (2 - weeks and 4 - weeks) of a treat-and-extend dosing regimen.

All patients received monthly doses of aflibercept 2 mg for 3 months, followed by one injection after a further 2-month interval. At week 16, patients were randomised 1:1 into two treatment groups: 1) aflibercept treat-and-extend with 2-week adjustments and 2) aflibercept treat-and-extend with 4 -week adjustments. Extension or shortening of the treatment interval was decided based on visual and/or anatomic criteria defined by protocol with a maximum treatment interval of 16 weeks for both groups.

The primary efficacy endpoint was mean change in BCVA from baseline to week 52. The secondary efficacy endpoints were the proportion of patients who did not lose ≥ 15 letters and the proportion of patients who gained at least 15 letters of BCVA from baseline to week 52.

At week 52, patients in the treat-and-extend arm with 2 -week adjustments gained a mean of 9.0 letters from baseline as compared to 8.4 letters for those in the 4 -week adjustment group [LS mean difference in letters (95% CI): - 0.4 (- 3.8,3.0), ANCOVA]. The proportion of patients who did not lose ≥ 15 letters in the two treatment arms was similar (96.7% in the 2 -week and 95.9% in the 4 -week adjustment groups). The proportion of patients who gained ≥ 15 letters at week 52 was 32.5% in the 2 -week adjustment group and 30.9% in the 4 -week adjustment group. The proportion of patients who extended their treatment interval to 12 weeks or beyond was 42.3% in the 2 -week adjustment group and 49.6% in the 4 -week adjustment group. Furthermore, in the 4 -week adjustment group 40.7% of patients were extended to 16 -week intervals. At the last visit up to week 52, 56.8% and 57.8% of

patients in the 2 -week and 4 -week adjustment groups, respectively had their next injection scheduled at an interval of 12 weeks or beyond.

In the second year of the study, efficacy was generally maintained up to and including the last assessment at week 96, with a mean gain from baseline of 7.6 letters for the 2 -week adjustment group and 6.1 letters for the 4 -week adjustment group. The proportion of patients who extended their treatment interval to 12 weeks or beyond was 56.9% in the 2 -week adjustment group and 60.2% in the 4 -week adjustment group. At the last visit prior to week 96, 64.9% and 61.2% of patients in the 2 -week and 4 -week adjustment groups, respectively had their next injection scheduled at an interval of 12 weeks or beyond. During the second year of treatment patients in both the 2 -week and 4 -week adjustment groups received an average of 3.6 and 3.7 injections, respectively. Over the 2-year treatment period patients received an average of 10.4 injections.

Ocular and systemic safety profiles were similar to the safety observed in the pivotal studies VIEW1 and VIEW2.

ARIES was a 104 -week multicentre, randomised, open-label, active-controlled study in 269 patients with treatment naïve wet AMD, designed to assess the non-inferiority in terms of efficacy as well as the safety of a treat-and-extend dosing regimen initiated after 3 consecutive monthly doses followed by extension to a 2 monthly treatment interval vs. a treat-and-extend dosing regimen initiated after the first year of treatment.

The ARIES study also explored the percentage of patients that required more frequent treatment than every 8 weeks based on the investigator's decision. Out of the 269 patients 62 patients received more frequent dosing at least once during the course of the study. Such patients remained in the study and received treatment according to the investigator's best clinical judgement but not more frequently than every 4 weeks and their treatment intervals could be extended again afterwards. The average treatment interval after the decision to treat more frequently was 6.1 weeks. Week 104 BCVA was lower in patients requiring more intensive treatment at least once over the course of the study compared with patients who did not and the mean change in BCVA from baseline to end of the study was $+2.3 \pm 15.6$ letters. Among the patients treated more frequently, 85.5% maintained vision, i.e., lost less than 15 letters, and 19.4% gained 15 letters or more. The safety profile of patients treated more frequently than every 8 weeks was comparable to the safety data in VIEW 1 and VIEW 2.

Macular oedema secondary to CRVO

The safety and efficacy of aflibercept were assessed in two randomised, multi-centre, double-masked, sham-controlled studies in patients with macular oedema secondary to CRVO (COPERNICUS and GALILEO) with a total of 358 patients treated and evaluable for efficacy (217 with aflibercept). Patient ages ranged from 22 to 89 years with a mean of 64 years. In the CRVO studies, approximately 52% (112/217) of the patients randomised to treatment with aflibercept were 65 years of age or older, and approximately 18% (38/217) were 75 years of age or older. In both studies, patients were randomly assigned in a 3:2 ratio to either 2 mg aflibercept administered every 4 weeks (2Q4), or the control group receiving sham injections every 4 weeks for a total of 6 injections.

After 6 consecutive monthly injections, patients received treatment only if they met pre-specified retreatment criteria, except for patients in the control group in the GALILEO study who continued to receive sham (control to control) until week 52. From this timepoint all patients were treated if pre-specified criteria were met.

In both studies, the primary efficacy endpoint was the proportion of patients who gained at least 15 letters in BCVA at week 24 compared to baseline. A secondary efficacy variable was change in visual acuity at week 24 compared to baseline.

The difference between treatment groups was statistically significant in favour of aflibercept in both studies. The maximal improvement in visual acuity was achieved at month 3 with subsequent stabilisation of visual acuity and CRT until month 6. The statistically significant difference was maintained through week 52.

Detailed results from the analysis of both studies are shown in Table 3 and Figure 2 below.

Table 3: Efficacy outcomes at week 24, week 52 and week 76/100 (full analysis set with LOCF^C) in COPERNICUS and GALILEO studies

Efficacy Outcomes	COPERNICUS						GALILEO					
	24 Weeks		52 Weeks		100 Weeks		24 Weeks		52 Weeks		76 Weeks	
	Aflibercept 2 mg Q4 (N=114)	Control (N=73)	Aflibercept 2 mg (N=114)	Control ^E (N=73)	Aflibercept ^F 2 mg (N=114)	Control ^{E,F} (N=73)	Aflibercept 2 mg Q4 (N=103)	Control (N=68)	Aflibercept 2 mg (N=103)	Control (N=68)	Aflibercept ^G 2 mg (N=103)	Control ^G (N=68)
Proportion of patients with ≥ 15 letters gain from baseline	56%	12%	55%	30%	49.1%	23.3%	60%	22%	60%	32%	57.3%	29.4%
Weighted difference ^{A,B,E} (95% CI) p-value	44.8% (33.0, 56.6) p < 0.0001		25.9% (11.8, 40.1) p = 0.0006		26.7% (13.1, 40.3) p = 0.0003		38.3% (24.4, 52.1) p < 0.0001		27.9% (13.0, 42.7) p = 0.0004		28.0% (13.3, 42.6) p = 0.0004	
Mean change in BCVA ^C as measured by ETDRS ^C letter score from baseline (SD)	17.3 (12.8)	- 4.0 (18.0)	16.2 (17.4)	3.8 (17.1)	13.0 (17.7)	1.5 (17.7)	18.0 (12.2)	3.3 (14.1)	16.9 (14.8)	3.8 (18.1)	13.7 (17.8)	6.2 (17.7)
Difference in LS mean ^{A,C,D,E} (95% CI) p-value	21.7 (17.4, 26.0) p < 0.0001		12.7 (7.7, 17.7) p < 0.0001		11.8 (6.7, 17.0) p < 0.0001		14.7 (10.8, 18.7) p < 0.0001		13.2 (8.2, 18.2) p < 0.0001		7.6 (2.1, 13.1) p=0.0070	

A) Difference is aflibercept 2 mg Q4 weeks minus control

B) Difference and confidence interval (CI) are calculated using Cochran-Mantel-Haenszel (CMH) test adjusted for region (America vs. rest of the world for COPERNICUS and Europe vs. Asia/Pacific for GALILEO) and baseline BCVA category ($> 20/200$ and $\leq 20/200$)

C) BCVA: Best Corrected Visual Acuity ETDRS: Early Treatment Diabetic Retinopathy Study LOCF: Last Observation Carried Forward SD: Standard deviation LS: Least square means derived from ANCOVA


D) LS mean difference and confidence interval based on an ANCOVA model with factors treatment group, region (America vs. rest of the world for COPERNICUS and Europe vs. Asia/Pacific for GALILEO) and baseline BCVA category ($> 20/200$ and $\leq 20/200$)

E) In COPERNICUS study, control group patients could receive aflibercept on an as-needed basis as frequently as every 4 weeks during week 24 to week 52; patients had visits every 4 weeks.

F) In COPERNICUS study, both control group and aflibercept 2 mg patients received aflibercept 2 mg on an as-needed basis as frequently as every 4 weeks starting from week 52 to week 96; patients had mandatory quarterly visits but may have been seen as frequently as every 4 weeks if necessary.

G) In GALILEO study, both control group and aflibercept 2 mg patients received aflibercept 2 mg on an as-needed basis every 8 weeks starting from week 52 to week 68; patients had mandatory visits every 8 weeks.

Figure 2: Mean change from baseline to week 76/100 in visual acuity by treatment group for the COPERNICUS and GALILEO studies (full analysis set)



✗ Indicates the switch of the control group to PRN treatment with Aflibercept 2 mg

In GALILEO, 86.4% (n = 89) of the aflibercept group and 79.4% (n = 54) of the sham group had perfused CRVO at baseline. At week 24, this was 91.8% (n = 89) in the aflibercept group and 85.5% (n = 47) in the sham group. These proportions were maintained at week 76, with 84.3% (n = 75) in the aflibercept group and 84.0% (n = 42) in the sham group.

In COPERNICUS, 67.5% (n = 77) of the aflibercept group and 68.5% (n = 50) of the sham group had perfused CRVO at baseline. At week 24, this was 87.4% (n = 90) in the aflibercept group and 58.6% (n = 34) in the sham group. These proportions were maintained at week 100 with 76.8% (n = 76) in the aflibercept group and 78% (n = 39) in the sham group. Patients in the sham group were eligible to receive aflibercept from week 24.

The beneficial effect of aflibercept treatment on visual function was similar in the baseline subgroups of perfused and non-perfused patients. Treatment effects in other evaluable subgroups (e.g., age,

gender, race, baseline visual acuity, CRVO duration) in each study were in general consistent with the results in the overall populations.

In combined data analysis of GALILEO and COPERNICUS, aflibercept demonstrated clinically meaningful changes from baseline in pre-specified secondary efficacy endpoint National Eye Institute Visual Function Questionnaire (NEI VFQ-25). The magnitude of these changes was similar to that seen in published studies, which corresponded to a 15-letter gain in Best Corrected Visual Acuity (BCVA).

Macular oedema secondary to BRVO

The safety and efficacy of aflibercept were assessed in a randomised, multi-centre, double-masked, active-controlled study in patients with macular oedema secondary to BRVO (VIBRANT) which included Hemi-Retinal Vein Occlusion. A total of 181 patients were treated and evaluable for efficacy (91 with aflibercept). Patient ages ranged from 42 to 94 years with a mean of 65 years. In the BRVO study, approximately 58% (53/91) of the patients randomised to treatment with aflibercept were 65 years of age or older, and approximately 23% (21/91) were 75 years of age or older. In the study, patients were randomly assigned in a 1:1 ratio to either 2 mg aflibercept administered every 8 weeks following 6 initial monthly injections or laser photocoagulation administered at baseline (laser control group). Patients in the laser control group could receive additional laser photocoagulation (called 'rescue laser treatment') beginning at week 12 with a minimum interval of 12 weeks. Based on pre-specified criteria, patients in the laser group could receive rescue treatment with aflibercept 2 mg from week 24, administered every 4 weeks for 3 months followed by every 8 weeks.

In the VIBRANT study, the primary efficacy endpoint was the proportion of patients who gained at least 15 letters in BCVA at week 24 compared to baseline and the aflibercept group was superior to laser control.

A secondary efficacy endpoint was change in visual acuity at week 24 compared to baseline, which was statistically significant in favour of aflibercept in the VIBRANT study. The course of visual improvement was rapid and peaked at 3 months with maintenance of the effect until month 12.

In the laser group 67 patients received rescue treatment with aflibercept beginning at week 24 (Active Control/ aflibercept 2 mg group), which resulted in improvement of visual acuity by about 5 letters from week 24 to 52.

Detailed results from the analysis of the VIBRANT study are shown in Table 4 and Figure 3 below.

Table 4: Efficacy outcomes at week 24 and week 52 (full analysis set with LOCF) in VIBRANT study

Efficacy Outcomes	VIBRANT			
	24 Weeks		52 Weeks	
	Aflibercept 2 mg Q4 (N = 91)	Active Control (laser) (N = 90)	Aflibercept 2 mg Q8 (N = 91) ^{D)}	Active Control (laser)/aflibercept 2 mg ^{E)} (N = 90)
Proportion of patients with ≥ 15 letters gain from baseline (%)	52.7%	26.7%	57.1%	41.1%
Weighted Difference ^{A,B)} (%) (95% CI) p-value	26.6% (13.0, 40.1) p = 0.0003		16.2% (2.0, 30.5) p = 0.0296	
Mean change in BCVA as measured by ETDRS letter score from baseline (SD)	17.0 (11.9)	6.9 (12.9)	17.1 (13.1)	12.2 (11.9)
Difference in LS mean ^{A,C)} (95% CI) p-value	10.5 (7.1, 14.0) p < 0.0001		5.2 (1.7, 8.7) p = 0.0035 ^{F)}	

^{A)} Difference is aflibercept 2 mg Q4 weeks minus Laser Control

^{B)} Difference and 95% CI are calculated using Mantel-Haenszel weighting scheme adjusted for region (North America vs. Japan) and baseline BCVA category ($> 20/200$ and $\leq 20/200$)


^{C)} LS mean difference and 95% CI based on an ANCOVA model with treatment group, baseline BCVA category ($> 20/200$ and $\leq 20/200$) and region (North America vs. Japan) as fixed effects, and baseline BCVA as covariate.

^{D)} From week 24 on the treatment interval in the aflibercept treatment group was extended for all subjects from 4 weeks to 8 weeks through week 48.

^{E)} Beginning at week 24 subjects in the Laser Group could receive rescue treatment with aflibercept, if they met at least one pre-specified eligibility criterion. At total of 67 subjects in this group received aflibercept rescue treatment. The fixed regimen for aflibercept rescue was three times aflibercept 2 mg every 4 weeks followed by injections every 8 weeks.

^{F)} Nominal p-value

Figure 3: Mean change in BCVA as measured by ETDRS letter score from baseline to week 52 in VIBRANT Study



At baseline, the proportion of perfused patients in the aflibercept and laser groups was 60% and 68%, respectively. At week 24 these proportions were 80% and 67%, respectively. In the aflibercept group the proportion of perfused patients was maintained through week 52. In the laser group, where patients were eligible for rescue treatment with aflibercept from week 24, the proportion of perfused patients increased to 78% by week 52.

Diabetic macular oedema

The safety and efficacy of aflibercept were assessed in two randomised, multi-centre, double-masked, active-controlled studies in patients with DME (VIVID^{DME} and VISTA^{DME}). A total of 862 patients were treated and evaluable for efficacy, 576 with aflibercept. Patient ages ranged from 23 to 87 years with a mean of 63 years. In the DME studies, approximately 47% (268/576) of the patients randomised to treatment with aflibercept were 65 years of age or older, and approximately 9% (52/576) were 75 years of age or older. The majority of patients in both studies had Type II diabetes.

In both studies, patients were randomly assigned in a 1:1:1 ratio to 1 of 3 dosing regimens:

- 1) Aflibercept administered 2 mg every 8 weeks following 5 initial monthly injections (aflibercept 2Q8);
- 2) Aflibercept administered 2 mg every 4 weeks (aflibercept 2Q4); and
- 3) Macular laser photocoagulation (active control).

Beginning at week 24, patients meeting a pre-specified threshold of vision loss were eligible to receive additional treatment: patients in the aflibercept groups could receive laser and patients in the control group could receive aflibercept.

In both studies, the primary efficacy endpoint was the mean change from baseline in BCVA at week 52 and both aflibercept 2Q8 and aflibercept 2Q4 groups demonstrated statistical significance and were superior to the control group. This benefit was maintained through week 100.

Detailed results from the analysis of the VIVID^{DME} and VISTA^{DME} studies are shown in Table 5 and Figure 4 below.

Table 5: Efficacy outcomes at week 52 and week 100 (full analysis set with LOCF) in VIVID^{DME} and VISTA^{DME} studies

Efficacy Outcomes	VIVID ^{DME}						VISTA ^{DME}					
	52 Weeks			100 Weeks			52 Weeks			100 Weeks		
	Aflibercept 2 mg Q8 ^A (N=135)	Aflibercept 2 mg Q4 (N=136)	Active Control (laser) (N=132)	Aflibercept 2 mg Q8 ^A (N=135)	Aflibercept 2 mg Q4 (N=136)	Active Control (laser) (N=132)	Aflibercept 2 mg Q8 ^A (N=151)	Aflibercept 2 mg Q4 (N=154)	Active Control (laser) (N=154)	Aflibercept 2 mg Q8 ^A (N=151)	Aflibercept 2 mg Q4 (N=154)	Active Control (laser) (N=154)
Mean change in BCVA as measured by ETDRS ^E letter score from Baseline	10.7	10.5	1.2	9.4	11.4	0.7	10.7	12.5	0.2	11.1	11.5	0.9
Difference in LS mean _{B,C,E} (97.5% CI)	9.1 (6.3, 11.8)	9.3 (6.5, 12.0)		8.2 (5.2, 11.3)	10.7 (7.6, 13.8)		10.45 (7.7, 13.2)	12.19 (9.4, 15.0)		10.1 (7.0, 13.3)	10.6 (7.1, 14.2)	
Proportion of patients with ≥ 15 letters gain from Baseline	33%	32%	9%	31.1%	38.2%	12.1%	31%	42%	8%	33.1%	38.3%	13.0%
Adjusted Difference _{D,C,E} (97.5% CI)	24% (13.5, 34.9)	23% (12.6, 33.9)		19.0% (8.0, 29.9)	26.1% (14.8, 37.5)		23% (13.5, 33.1)	34% (24.1, 44.4)		20.1% (9.6, 30.6)	25.8% (15.1, 36.6)	

^A After treatment initiation with 5 monthly injections

^B LS mean and CI based on an ANCOVA model with baseline BCVA measurement as a covariate and a factor for treatment group. Additionally, region (Europe/Australia vs. Japan) had been included as factor for VIVID^{DME}, and history of MI and/or CVA as a factor for VISTA^{DME}


^C Difference is aflibercept group minus active control (laser) group

^D Difference with confidence interval (CI) and statistical test is calculated using Mantel-Haenszel weighting scheme adjusted by region (Europe/Australia vs. Japan) for VIVID^{DME} and medical history of MI or CVA for VISTA^{DME}

^E BCVA: Best Corrected Visual Acuity

ETDRS: Early Treatment Diabetic Retinopathy Study; LOCF: Last Observation Carried Forward; LS: Least square means derived from ANCOVA; CI: Confidence interval

Figure 4: Mean change in BCVA as measured by ETDRS letter score from baseline to week 100 in VIVID^{DME} and VISTA^{DME} studies



Treatment effects in evaluable subgroups (e.g., age, gender, race, baseline HbA1c, baseline visual acuity, prior anti-VEGF therapy) in each study and in the combined analysis were generally consistent with the results in the overall populations.

In the VIVID^{DME} and VISTA^{DME} studies, 36 (9%) and 197 (43%) patients received prior anti-VEGF therapy, respectively, with a 3-month or longer washout period. Treatment effects in the subgroup of patients who had previously been treated with a VEGF inhibitor were similar to those seen in patients who were VEGF inhibitor naïve.

Patients with bilateral disease were eligible to receive anti-VEGF treatment in their fellow eye if assessed necessary by the physician. In the VISTA^{DME} study, 217 (70.7%) of aflibercept patients received bilateral aflibercept injections until week 100; in the VIVID^{DME} study, 97 (35.8%) of aflibercept patients received a different anti-VEGF treatment in their fellow eye.

An independent comparative trial (DRCR.net Protocol T) utilised a flexible dosing regimen based on strict OCT and vision re-treatment criteria. In the aflibercept treatment group (n = 224) at week 52, this treatment regimen resulted in patients receiving a mean of 9.2 injections, which is similar to the administered number of doses in the aflibercept 2Q8 group in VIVID^{DME} and VISTA^{DME}, while overall efficacy of the aflibercept treatment group in Protocol T was comparable to the aflibercept 2Q8 group in VIVID^{DME} and VISTA^{DME}. A 13.3 mean letter gain with 42% of patients gaining at least 15 letters in vision from baseline was observed in Protocol T. Safety outcomes demonstrated that overall

incidences of ocular and non-ocular adverse events (including ATEs) were comparable across all treatment groups in each of the studies and between the studies.

VIOLET, a 100-week multicentre, randomised, open-label, active controlled study in patients with DME compared three different dosing regimens of aflibercept 2 mg for treatment of DME after at least one year of treatment at fixed intervals, where treatment was initiated with 5 consecutive monthly doses followed by dosing every 2 months. The study evaluated non-inferiority of aflibercept 2 mg dosed according to a treat-and-extend regimen (2T&E where injections intervals were kept at a minimum of 8 weeks and gradually extended based on clinical and anatomical outcomes) and aflibercept 2 mg dosed as needed (2PRN where patients were observed every 4 weeks and injected when needed based on clinical and anatomical outcomes), compared to aflibercept 2 mg dosed every 8 weeks (2Q8) for the second and third year of treatment.

The primary efficacy endpoint (change in BCVA from baseline to week 52) was 0.5 ± 6.7 letters in the 2T&E group and 1.7 ± 6.8 letters in the 2PRN group compared to 0.4 ± 6.7 letters in the 2Q8 group, achieving statistical non-inferiority ($p < 0.0001$ for both comparisons; NI margin 4 letters). The changes in BCVA from baseline to week 100 were consistent with the week 52 results: -0.1 ± 9.1 letters in the 2T&E group and 1.8 ± 9.0 letters in the 2PRN group compared to 0.1 ± 7.2 letters in the 2Q8 group. The mean number of injections over 100 weeks were 12.3, 10.0 and 11.5 for 2Q8fix, 2T&E and 2PRN, respectively.

Ocular and systemic safety profiles in all 3 treatment groups were similar to those observed in the pivotal studies VIVID and VISTA.

In the 2T&E group, the increments and decrements for the injection intervals were at the investigator's discretion; increments of 2 weeks were recommended in the study.

Myopic choroidal neovascularisation

The safety and efficacy of aflibercept were assessed in a randomised, multi-centre, double-masked, sham-controlled study in treatment-naïve, Asian patients with myopic CNV. A total of 121 patients were treated and evaluable for efficacy (90 with aflibercept). Patient ages ranged from 27 to 83 years with a mean of 58 years. In the myopic CNV study, approximately 36% (33/91) of the patients randomised to treatment with aflibercept were 65 years of age or older, and approximately 10% (9/91) were 75 years of age or older.

Patients were randomly assigned in a 3:1 ratio to receive either 2 mg aflibercept intravitreally or sham injections administered once at study start with additional injections given monthly in case of disease persistence or recurrence until week 24, when the primary endpoint was assessed. At week 24, patients initially randomised to sham were eligible to receive the first dose of aflibercept. Following this, patients in both groups continued to be eligible for additional injections in case of disease persistence or recurrence.

The difference between treatment groups was statistically significant in favour of aflibercept for the primary endpoint (change in BCVA) and confirmatory secondary efficacy endpoint (proportion of patients who gained 15 letters in BCVA) at week 24 compared to baseline. Differences for both endpoints were maintained through week 48.

Detailed results from the analysis of the MYRROR study are shown in Table 6 and Figure 5 below.

Table 6: Efficacy outcomes at week 24 (primary analysis) and week 48 in MYRROR study (full analysis set with LOCF^{A)})

Efficacy Outcomes	MYRROR			
	24 Weeks		48 Weeks	
	Aflibercept 2 mg (N = 90)	Sham (N = 31)	Aflibercept 2 mg (N = 90)	Sham/Aflibercept 2 mg (N = 31)
Mean change in BCVA ^{B)} as measured by ETDRS letter score from baseline (SD) ^{B)}	12.1 (8.3)	-2.0 (9.7)	13.5 (8.8)	3.9 (14.3)
Difference in LS mean ^{C,D,E)} (95% CI)	14.1 (10.8, 17.4)		9.5 (5.4, 13.7)	
Proportion of patients with ≥ 15 letters gain from baseline	38.9%	9.7%	50.0%	29.0%
Weighted difference ^{D,F)} (95% CI)	29.2% (14.4, 44.0)		21.0% (1.9, 40.1)	

^{A)} LOCF: Last Observation Carried Forward

^{B)} BCVA: Best Corrected Visual Acuity

ETDRS: Early Treatment Diabetic Retinopathy Study

SD: Standard Deviation


^{C)} LS mean: Least square means derived from ANCOVA model

^{D)} CI: Confidence Interval

^{E)} LS mean difference and 95% CI based on an ANCOVA model with treatment group and country (country designations) as fixed effects, and baseline BCVA as covariant.

^{F)} Difference and 95% CI are calculated using Cochran-Mantel-Haenszel (CMH) test adjusted for country (country designations)

Figure 5: Mean change from baseline to week 48 in visual acuity by treatment group for the MYRROR study (full analysis set, LOCF)



Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies with aflibercept in all subsets of the paediatric population in wet AMD, CRVO, BRVO, DME and myopic CNV populations (see section 4.2 for information on paediatric use).

5.2 Pharmacokinetic properties

Aflibercept is administered directly into the vitreous to exert local effects in the eye.

Absorption / Distribution

Aflibercept is slowly absorbed from the eye into the systemic circulation after intravitreal administration and is predominately observed in the systemic circulation as an inactive, stable complex with VEGF; however only “free aflibercept” is able to bind endogenous VEGF.

In a pharmacokinetic sub-study in 6 neovascular wet AMD patients with frequent sampling, maximum plasma concentrations of free aflibercept (systemic C_{max}) were low, with a mean of approximately 0.02 microgram/mL (range 0 to 0.054) within 1 to 3 days after a 2 mg intravitreal injection and were undetectable two weeks following dose in almost all patients. Aflibercept does not accumulate in the plasma when administered intravitreally every 4 weeks.

The mean maximum plasma concentration of free aflibercept is approximately 50 to 500 times below the aflibercept concentration required to inhibit the biologic activity of systemic VEGF by 50% in animal models, in which blood pressure changes were observed after circulating levels of free aflibercept attained approximately 10 microgram/mL and returned to baseline when levels fell below approximately 1 microgram/mL. It is estimated that after intravitreal administration of 2 mg to patients, the mean maximum plasma concentration of free aflibercept is more than 100-fold lower than the concentration of aflibercept required to half-maximally bind systemic VEGF (2.91 microgram/mL) in a study of healthy volunteers. Therefore, systemic pharmacodynamic effects such as blood pressure changes are unlikely.

In pharmacokinetic sub-studies in patients with CRVO, BRVO, DME or myopic CNV mean C_{max} of free aflibercept in plasma were similar with values in the range of 0.03 to 0.05 microgram/mL and individual values not exceeding 0.14 microgram/mL. Thereafter, plasma concentrations of free aflibercept declined to values below or close to the lower limit of quantitation generally within one week; undetectable concentrations were reached before the next administration after 4 weeks in all patients.

Elimination

As aflibercept is a protein-based therapeutic, no metabolism studies have been conducted.

Free aflibercept binds VEGF to form a stable, inert complex. As with other large proteins, both free and bound aflibercept are expected to be cleared by proteolytic catabolism.

Renal impairment

No special studies in patients with renal impairment have been conducted with aflibercept.

Pharmacokinetic analysis of patients in the VIEW2 study, of which 40% had renal impairment (24% mild, 15% moderate, and 1% severe), revealed no differences with respect to plasma concentrations of active substance after intravitreal administration every 4 or 8 weeks.

Similar results were seen in patients with CRVO in the GALILEO study, in patients with DME in the VIVID^{DME} study, and in patients with myopic CNV in the MYRROR study.

5.3 Preclinical safety data

Effects in non-clinical studies on repeated dose toxicity were observed only at systemic exposures considered substantially in excess of the maximum human exposure after intravitreal administration at the intended clinical dose indicating little relevance to clinical use.

Erosions and ulcerations of the respiratory epithelium in nasal turbinates in monkeys treated with aflibercept intravitreally were observed at systemic exposures in excess of the maximum human exposure. The systemic exposure based on C_{max} and AUC for free aflibercept were approximately 200 - and 700 -fold higher, respectively, when compared to corresponding values observed in humans after an intravitreal dose of 2 mg. At the No Observed Adverse Effect Level (NOAEL) of 0.5 mg/eye in monkeys the systemic exposure was 42 - and 56 -fold higher based on C_{max} and AUC, respectively.

No studies have been conducted on the mutagenic or carcinogenic potential of aflibercept.

An effect of aflibercept on intrauterine development was shown in embryo-foetal development studies in pregnant rabbits with intravenous (3 to 60 mg/kg) as well as subcutaneous (0.1 to 1 mg/kg) administration. The maternal NOAEL was at the dose of 3 mg/kg or 1 mg/kg, respectively. A developmental NOAEL was not identified. At the 0.1 mg/kg dose, the systemic exposures based on C_{max} and cumulative AUC for free aflibercept were approximately 17 - and 10 -fold higher, respectively, when compared to corresponding values observed in humans after an intravitreal dose of 2 mg.

Effects on male and female fertility were assessed as part of a 6 -month study in monkeys with intravenous administration of aflibercept at doses ranging from 3 to 30 mg/kg. Absent or irregular menses associated with alterations in female reproductive hormone levels and changes in sperm morphology and motility were observed at all dose levels. Based on C_{max} and AUC for free aflibercept observed at the 3 mg/kg intravenous dose, the systemic exposures were approximately 4 900 -fold and 1 500 -fold higher, respectively, than the exposure observed in humans after an intravitreal dose of 2 mg. All changes were reversible.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Histidine
Histidine hydrochloride monohydrate
Polysorbate 20 (E 432)
Trehalose dihydrate
Water for injections

6.2 Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

Store in a refrigerator (2 °C - 8 °C). Do not freeze.
Store in the original package in order to protect from light.

The unopened vial may be stored outside the refrigerator below 25 °C for up to 24 hours. After opening the vial, proceed under aseptic conditions.

6.5 Nature and contents of container

Solution in a vial (type I glass) with a stopper (chlorobutyl rubber). Each vial contains an extractable volume of at least 0.1 mL.

Yesafili is available in 2 different packs:

One pack containing 1 vial and a 5-micron sterile filter needle (18 G ×1½-inch).

One pack containing 1 vial, a 5-micron sterile filter needle (18 G ×1½-inch), a 1 mL Luer-lock syringe and an injection needle (30 G ×½-inch).

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

The vial is for single use in one eye only.

The vial contains more than the recommended dose of 2 mg aflibercept (equivalent to 0.05 mL). The excess volume must be discarded prior to administration.

The solution should be inspected visually for any foreign particulate matter and/or discoloration or any variation in physical appearance prior to administration. In the event of either being observed, discard the medicinal product.

Preparation for intravitreal injection requires the following medical devices for single use:

Filter needle

BD blunt 5-micron sterile filter (fill) needle, not for skin injection.

Do not autoclave the BD blunt filter (fill) needle.

The filter needle is non-pyrogenic. Do not use it if individual packaging is damaged.

Discard the used BD blunt filter (fill) needle in approved sharps collector.

Caution: Re-use of the filter needle may lead to infection or other illness/injury.

Syringe (where supplied)

BD 1 mL, Luer-lock syringe.

Do not autoclave the BD 1 mL, Luer-lock syringe.

The syringe is non-pyrogenic. Do not use it if individual packaging is damaged.

Discard the used BD 1 mL, Luer-lock syringe, with injection needle attached, in approved sharps collector.

Caution: Re-use of the syringe may lead to infection or other illness/injury.

Injection needle (where supplied)

BD 30 G ×½-inch injection needle.

Do not autoclave the BD 30 G ×½-inch injection needle.

The injection needle is non-pyrogenic. Do not use it if individual packaging is damaged.

Discard the used BD 30 G ×½-inch injection needle, attached to the syringe, in approved sharps collector.

Caution: Re-use of the injection needle may lead to infection or other illness/injury.

For the intravitreal injection, a 30 G ×½-inch injection needle should be used.

- Yesafili is available in 2 different packs

Figure A: Pack size of 1 vial + 1 filter needle + 1 syringe + 1 injection needle








Figure B: Pack size of 1 vial + 1 filter needle




Instructions for use

<p>1. Remove the plastic cap and disinfect the outer part of the rubber stopper of the vial.</p>	
<p>2. The 18 G × 1½-inch, 5-micron, filter needle is used for withdrawal of drug product from the vial.</p> <p>Remove the 18 G × 1½-inch, 5-micron, filter needle and the 1 mL syringe from their packaging. Attach the filter needle to the syringe by twisting it onto the Luer lock syringe tip.</p>	
<p>3. Push the filter needle into the centre of the vial stopper until the needle is completely inserted into the vial and the tip touches the bottom or bottom edge of the vial.</p> <p>4. Using aseptic technique withdraw all of the Yesafili vial contents into the syringe, keeping the vial in an upright position, slightly inclined to ease complete withdrawal. To deter the introduction of air, ensure the bevel of the filter needle is submerged into the liquid. Continue to tilt the vial during withdrawal keeping the bevel of the filter needle submerged in the liquid.</p>	
<p>5. Ensure that the plunger rod is drawn sufficiently back when emptying the vial in order to completely empty the filter needle.</p>	
<p>6. Remove the filter needle and properly dispose of it.</p> <p>Note: Filter needle is not to be used for intravitreal injection.</p>	


7. Ensure the 30 G \times 1/2-inch injection needle is used for the intravitreal injection.

Remove the 30 G \times 1/2-inch injection needle from its packaging and using aseptic technique, firmly twist onto the Luer-lock syringe tip.





8. When ready to administer, remove the plastic needle shield from the needle.

Holding the syringe with the needle pointing up, check the syringe for bubbles. If there are bubbles, gently tap the syringe with your finger until the bubbles rise to the top.



9. Eliminate all bubbles and expel excess medicinal product by slowly depressing the plunger so that the flat plunger edge aligns with the line that marks 0.05 mL on the syringe.



10. The vial is for single use only. Extraction of multiple doses from a single vial may increase the risk of contamination and subsequent infection.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Biosimilar Collaborations Ireland Limited
Unit 35/36
Grange Parade,
Baldoyle Industrial Estate,
Dublin 13
DUBLIN
Ireland
D13 R20R

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/23/1751/001
EU/1/23/1751/002

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorization: 15 September 2023

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency (EMA): <http://www.ema.europa.eu>

ANNEX II

- A. MANUFACTURER OF THE BIOLOGICAL ACTIVE SUBSTANCE AND MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

A. MANUFACTURER OF THE BIOLOGICAL ACTIVE SUBSTANCE AND MANUFACTURER RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturer of the biological active substance

WuXi Biologics Co. Ltd.
108 Meiliang Road,
Mashan, Binhu District,
Wuxi, Jiangsu 214092
China

Name and address of the manufacturer responsible for batch release

McDermott Laboratories Limited T/A Mylan Dublin Biologics
Newenham Court, Northern Cross, Malahide Road
Dublin 17
Ireland

Mylan Germany GmbH
Zweigniederlassung Bad Homburg v. d. Hoehe, Benzstrasse 1
Bad Homburg v. d. Hoehe
Hessen, 61352
Germany

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

- **Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being

received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

If the dates for submission of a PSUR and the update of a RMP coincide, they can be submitted at the same time.

Pharmacovigilance system

The MAH must ensure that the system of pharmacovigilance presented in Module 1.8.1. of the Marketing Authorisation is in place and functioning before and whilst the medicinal product is on the market.

- Additional risk minimisation measures

The MAH has agreed to provide EU educational material for Yesafili. Prior to launch and during the product's lifecycle in each Member State the MAH will agree the final educational material with the National Competent Authority.

The MAH ensures that, following discussions and agreement with the National Competent Authorities in each Member State where Yesafili is marketed, ophthalmological clinics where Yesafili is expected to be used are provided with an updated physician information pack containing the following elements:

- Physician information
- Intravitreal injection procedure video
- Intravitreal injection procedure pictogram
- Patient information packs

The physician information in the educational material contains the following key elements:

- Techniques for the intravitreal injection including use of a 30 G needle and angle of injection
- Confirmation that the vial is for single use only
- The need to expel excess volume of the syringe before injecting Yesafili to avoid overdose
- Patient monitoring after intravitreal injection including monitoring for visual acuity and increase of intraocular pressure post-injection
- Key signs and symptoms of intravitreal injection related adverse events including endophthalmitis, intraocular inflammation, increased intraocular pressure, retinal pigment epithelial tear and cataract
- Female patients of childbearing potential have to use effective contraception and pregnant women should not use Yesafili

The patient information pack of the educational material includes a patient information guide and its audio version. The patient information guide contains following key elements:

- Patient information leaflet
- Who should be treated with Yesafili
- How to prepare for Yesafili treatment
- What are the steps following treatment with Yesafili
- Key signs and symptoms of serious adverse events including endophthalmitis, intraocular inflammation, intraocular pressure increased, retinal pigment epithelial tear and cataract
- When to seek urgent attention from their health care provider
- Female patients of childbearing potential have to use effective contraception and pregnant women should not use Yesafili

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING
CARTON**

Vial

1. NAME OF THE MEDICINAL PRODUCT

Yesafili 40 mg/mL solution for injection in a vial
aflibercept

2. STATEMENT OF ACTIVE SUBSTANCE

1 vial contains 4 mg aflibercept in 0.1 mL solution (40 mg/mL).

3. LIST OF EXCIPIENTS

Histidine; histidine hydrochloride monohydrate; polysorbate 20 (E 432); trehalose dihydrate; water for injections.

4. PHARMACEUTICAL FORM AND CONTENTS

Solution for injection

1 vial contains 4 mg aflibercept in 0.1 mL solution (40 mg/mL).
4 mg/0.1 mL.

1 x 18 G × 1½-inch filter needle
1 x 18 G × 1½-inch filter needle
1 x 1 mL Luer-lock syringe
1 x 30 G × ½ inch injection needle

Delivers 1 single dose of 2 mg/0.05 mL.

5. METHOD AND ROUTE OF ADMINISTRATION

Intravitreal use.
For single use only.
Read the package leaflet before use.
Excess volume to be expelled before injecting.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT
OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Store in a refrigerator. Do not freeze.

Store in the original package in order to protect from light.

The unopened vial may be stored outside the refrigerator below 25 °C for up to 24 hours.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Biosimilar Collaborations Ireland Limited
Unit 35/36
Grange Parade,
Baldoyle Industrial Estate,
Dublin 13
DUBLIN
Ireland
D13 R20R

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/23/1751/001
EU/1/23/1751/002

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS LABEL

Vial

1. NAME OF THE MEDICINAL PRODUCT AND ROUTE OF ADMINISTRATION

Yesafili 40 mg/mL injection
aflibercept
Intravitreal use

2. METHOD OF ADMINISTRATION

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT

Extractable volume 0.1 mL

6. OTHER

B. PACKAGE LEAFLET

Package Leaflet: Information for the patient

Yesafili 40 mg/mL solution for injection in a vial aflibercept

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

Read all of this leaflet carefully before you are given this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor.
- If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Yesafili is and what it is used for
2. What you need to know before you are given Yesafili
3. How you will be given Yesafili
4. Possible side effects
5. How to store Yesafili
6. Contents of the pack and other information

1. What Yesafili is and what it is used for

Yesafili is a solution which is injected into the eye to treat eye conditions in adults called

- neovascular (wet) age-related macular degeneration (wet AMD),
- impaired vision due to macular oedema secondary to retinal vein occlusion (branch RVO (BRVO) or central RVO (CRVO)),
- impaired vision due to diabetic macular oedema (DME),
- impaired vision due to myopic choroidal neovascularisation (myopic CNV).

Aflibercept, the active substance in Yesafili, blocks the activity of a group of factors, known as vascular endothelial growth factor A (VEGF-A) and placental growth factor (PlGF).

In patients with wet AMD and myopic CNV, these factors, in excess are involved in the abnormal formation of new blood vessels in the eye. These new blood vessels can cause the leak of blood components into the eye and eventual damage to tissues in the eye responsible for vision.

In patients with CRVO, a blockage occurs in the main blood vessel that transports blood away from the retina. VEGF levels are elevated in response causing the leakage of fluid into the retina and thereby causing a swelling of the macula, (the portion of the retina responsible for fine vision), which is called macular oedema. When the macula swells with fluid, central vision becomes blurry.

In patients with BRVO, one or more branches of the main blood vessel that transports blood away from the retina is blocked. VEGF levels are elevated in response causing the leakage of fluid into the retina and thereby causing macular oedema.

Diabetic macular oedema is a swelling of the retina occurring in patients with diabetes due to leaking of fluid from blood vessels within the macula. The macula is the portion of retina responsible for fine vision. When the macula swells with fluid, central vision becomes blurry.

Aflibercept has been shown to stop the growth of new abnormal blood vessels in the eye which often

leak fluid or bleed. Aflibercept can help to stabilise, and in many cases, improve the vision loss related to wet AMD, CRVO, BRVO, DME and myopic CNV.

2. What you need to know before you are given Yesafili

You will not be given Yesafili

- if you are allergic to aflibercept or any of the other ingredients of this medicine (listed in section 6).
- if you have an active or suspected infection in or around the eye (ocular or periocular infection).
- if you have severe inflammation of the eye (indicated by pain or redness).

Warnings and precautions

Talk to your doctor before you are given Yesafili:

- if you have glaucoma.
- if you have a history of seeing flashes of light or floaters and if you have a sudden increase of size and number of floaters.
- if surgery was performed or is planned on your eye within the previous or next four weeks.
- if you have a severe form of CRVO or BRVO (ischaemic CRVO or BRVO), treatment with Yesafili is not recommended.

Furthermore, it is important for you to know that:

- the safety and efficacy of Aflibercept when administered to both eyes at the same time has not been studied and if used in this way may lead to an increased risk of experiencing side effects.
- injections with Yesafili may cause an increase in eye pressure (intraocular pressure) in some patients within 60 minutes of the injection. Your doctor will monitor this after each injection.
- if you develop an infection or inflammation inside the eye (endophthalmitis) or other complications, you may have eye pain or increased discomfort, worsening eye redness, blurred or decreased vision, and increased sensitivity to light. It is important to have any symptoms diagnosed and treated as soon as possible.
- your doctor will check whether you have other risk factors that may increase the chance of a tear or detachment of one of the layers at the back of the eye (retinal detachment or tear, and retinal pigment epithelial detachment or tear), in which case Aflibercept must be given with caution.
- Aflibercept should not be used in pregnancy unless the potential benefit outweighs the potential risk to the unborn child.
- women of childbearing potential have to use effective contraception during treatment and for at least three further months after the last injection of Yesafili.

The systemic use of VEGF inhibitors, substances similar to those contained in Yesafili, is potentially related to the risk of blood clots blocking blood vessels (arterial thromboembolic events) which may lead to heart attack or stroke. There is a theoretical risk of such events following injection of Yesafili into the eye. There are limited data on safety in treating patients with CRVO, BRVO, DME and myopic CNV who have had a stroke or a mini-stroke (transient ischaemic attack) or a heart attack within the last 6 months. If any of these apply to you, Yesafili will be given with caution.

There is only limited experience in the treatment of

- patients with DME due to type I diabetes.
- diabetics with very high average blood sugar values (HbA1c over 12%).
- diabetics with an eye disease caused by diabetes called proliferative diabetic retinopathy.

There is no experience in the treatment of

- patients with acute infections.

- patients with other eye conditions such as a detachment of the retina or a hole in the macula.
- diabetics with uncontrolled high blood pressure.
- non-Asian patients with myopic CNV.
- patients previously treated for myopic CNV.
- patients with damage outside the central part of the macula (extrafoveal lesions) for myopic CNV.

If any of the above applies to you, your doctor will consider this lack of information when treating you with Yesafili.

Children and adolescents

The use of aflibercept in children or adolescents under 18 has not been studied because wet AMD, CRVO, BRVO, DME and myopic CNV occur mainly in adults. Therefore, its use in this age group is not relevant.

Other medicines and Yesafili

Tell your doctor if you are using, have recently used or might use any other medicines.

Pregnancy and breast-feeding

Women of childbearing potential have to use effective contraception during treatment and for at least three further months after the last injection of Yesafili.

There is no experience of using aflibercept in pregnant women. Yesafili should not be used during pregnancy unless the potential benefit outweighs the potential risk to the unborn child. If you are pregnant or planning to become pregnant, discuss this with your doctor before treatment with Yesafili.

Yesafili is not recommended during breast-feeding as it is not known whether aflibercept passes into human milk. Ask your doctor for advice before starting Yesafili treatment.

Driving and using machines

After your injection with Yesafili, you may experience some temporary visual disturbances. Do not drive or use machines as long as these last.

Important information about some of the ingredients of Yesafili

This medicine contains less than 1 mmol sodium (23 mg) per dosage unit, that is to say essentially 'sodium-free'.

3. How you will be given Yesafili

A doctor experienced in giving eye injections will inject Yesafili into your eye under aseptic (clean and sterile) conditions.

The recommended dose is 2 mg aflibercept (0.05 mL).
Yesafili is given as an injection into your eye (intravitreal injection).

Before the injection your doctor will use a disinfectant eyewash to clean your eye carefully to prevent infection. Your doctor will also give you a local anaesthetic to reduce or prevent any pain you might have with the injection.

wet AMD

Patients with wet AMD will be treated with one injection per month for three consecutive doses, followed by another injection after a further two months.

Your doctor will then decide whether the treatment interval between injections may be kept at every two months or be gradually extended in 2 - or 4 -weekly intervals if your condition has been stable.

If your condition worsens, the interval between injections can be shortened.

Unless you experience any problems or are advised differently by your doctor, there is no need for you to see your doctor between the injections.

Macular oedema secondary to RVO (branch RVO or central RVO)

Your doctor will determine the most appropriate treatment schedule for you. You will start your treatment with a series of monthly Yesafili injections.

The interval between two injections should not be shorter than one month.

Your doctor may decide to stop treatment with Yesafili, if you are not benefiting from continued treatment.

Your treatment will continue with monthly injections until your condition is stable. Three or more monthly injections may be needed.

Your doctor will monitor your response to treatment and may continue your treatment by gradually increasing the interval between your injections to maintain a stable condition. If your condition starts to worsen with a longer treatment interval, your doctor will shorten the interval accordingly.

Based on your response to treatment your doctor will decide on the schedule for follow up examinations and treatments.

Diabetic macular oedema (DME)

Patients with DME will be treated with one injection per month for the first five consecutive doses followed by one injection every two months thereafter.

Treatment interval may be kept at every two months or adjusted to your condition, based on your doctor's examination. Your doctor will decide on the schedule for follow up examinations.

Your doctor may decide to stop treatment with Yesafili if it is determined that you are not benefiting from continued treatment.

Myopic CNV

Patients with myopic CNV will be treated with one single injection. You will receive further injections only if your doctor's examinations reveal that your condition has not improved.

The interval between two injections should not be shorter than one month.

If your condition goes away and then comes back, your doctor may re-start the treatment. Your doctor will decide on the schedule for follow up examinations.

If a dose of Yesafili is missed

Make a new appointment for an examination and injection.

Stopping treatment with Yesafili

Consult your doctor before stopping the treatment.

If you have any further questions on the use of this medicine, ask your doctor.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Allergic reactions (hypersensitivity) could potentially occur. **These may be serious and require that you contact your doctor immediately.**

With administration of aflibercept, there may be some side effects affecting the eyes which are due to the injection procedure. Some of these may be **serious** and include **blindness, a serious infection or inflammation inside the eye** (endophthalmitis), **detachment, tear or bleeding of the light-sensitive layer at the back of the eye** (retinal detachment or tear), **clouding of the lens** (cataract), **bleeding in the eye** (vitreous haemorrhage), **detachment of the gel-like substance inside the eye from the retina** (vitreous detachment) and **increase of pressure inside the eye**, see section 2. These serious side effects affecting the eyes occurred in less than 1 in 1 900 injections in clinical studies.

If you experience a sudden decrease in vision, or an increase in pain and redness in your eye after your injection, **contact your doctor immediately.**

List of side effects reported

The following is a list of the side effects reported to be possibly related to the injection procedure or to the medicine. Please do not get alarmed, you might not experience any of these. Always discuss any suspected side effects with your doctor.

Very common side effects (*may affect more than 1 in 10 people*):

- deterioration of eyesight
- bleeding in the back of the eye (retinal haemorrhage)
- bloodshot eye caused by bleeding from small blood vessels in the outer layers of the eye
- eye pain

Common side effects (*may affect up to 1 in 10 people*):

- detachment or tear of one of the layers in the back of the eye, resulting in flashes of light with floaters sometimes progressing to a loss of vision (retinal pigment epithelial tear*/detachment, retinal detachment/tear)
- degeneration of the retina (causing disturbed vision)
- bleeding in the eye (vitreous haemorrhage)
- certain forms of clouding of the lens (cataract)
- damage to the front layer of the eyeball (the cornea)
- increase in eye pressure
- moving spots in vision (floaters)
- detachment of the gel-like substance inside the eye from the retina (vitreous detachment, resulting in flashes of light with floaters)
- a feeling of having something in the eye
- increased tear production
- swelling of the eyelid
- bleeding at the injection site
- redness of the eye
- * Conditions known to be associated with wet AMD; observed in wet AMD patients only.

Uncommon side effects (*may affect up to 1 in 100 people*):

- allergic reactions (hypersensitivity)**
- serious inflammation or infection inside the eye (endophthalmitis)
- inflammation in the iris or other parts of the eye (iritis, uveitis, iridocyclitis, anterior chamber flare)
- abnormal sensation in the eye
- eyelid irritation
- swelling of the front layer of the eyeball (cornea)

** Allergic reactions like rash, itching (pruritus), hives (urticaria), and a few cases of severe allergy (anaphylactic/anaphylactoid) reactions were reported.

Rare side effects (*may affect up to 1 in 1 000 people*):

- blindness
- clouding of the lens due to injury (traumatic cataract)
- inflammation of the gel-like substance inside the eye
- pus in the eye

In the clinical trials, there was an increased incidence of bleeding from small blood vessels in the outer layers of the eye (conjunctival haemorrhage) in patients with wet AMD receiving blood thinners. This increased incidence was comparable between patients treated with ranibizumab and aflibercept.

The systemic use of VEGF inhibitors, substances similar to those contained in Yesafili, is potentially related to the risk of blood clots blocking blood vessels (arterial thromboembolic events) which may lead to heart attack or stroke. There is a theoretical risk of such events following injection of aflibercept into the eye.

As with all therapeutic proteins, there is a possibility for an immune reaction (formation of antibodies) with Yesafili.

Reporting of side effects

If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in [Appendix V](#). By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Yesafili

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton and label after EXP. The expiry date refers to the last day of that month.
- Store in a refrigerator (2 °C – 8 °C). Do not freeze.
- The unopened vial may be stored outside the refrigerator below 25 °C for up to 24 hours.
- Store in the original package in order to protect from light.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away any medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Yesafili contains

- The active substance is: aflibercept. One vial contains an extractable volume of at least 0.1 mL, equivalent to at least 4 mg aflibercept. One vial delivers a dose of 2 mg aflibercept in 0.05 mL.
- The other ingredients are: histidine; histidine hydrochloride monohydrate; polysorbate 20 (E 432); trehalose dihydrate and water for injections.

What Yesafili looks like and contents of the pack

Yesafili is a solution for injection (injection) in a vial. The solution is colourless to pale yellow. Yesafili is supplied as packs containing 1 vial and 1 filter needle or as packs containing 1 vial, 1 filter needle, 1 syringe and 1 injection needle.

Marketing Authorisation Holder

Biosimilar Collaborations Ireland Limited
Unit 35/36
Grange Parade,
Baldoyle Industrial Estate,
Dublin 13
DUBLIN
Ireland
D13 R20R

Manufacturer

McDermott Laboratories Limited T/A Mylan Dublin Biologics
Newenham Court,
Northern Cross,
Malahide Road
Dublin 17
Ireland

Mylan Germany GmbH
Zweigniederlassung Bad Homburg v. d. Hoehe, Benzstrasse 1
Bad Homburg v. d. Hoehe
Hessen, 61352
Germany

This leaflet was last revised in

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency web site:
<http://www.ema.europa.eu>

←-----

The following information is intended for healthcare professionals only

The vial should only be used **for the treatment of a single eye**.

The vial contains more than the recommended dose of 2 mg aflibercept (equivalent to 0.05 mL). The excess volume must be discarded prior to administration.

The solution should be inspected visually for any foreign particulate matter and/or discolouration or any variation in physical appearance prior to administration. In the event of either being observed, discard the medicinal product.

The unopened vial may be stored outside the refrigerator below 25 °C for up to 24 hours. After opening the vial, proceed under aseptic conditions.
For the intravitreal injection, a 30 G ×½-inch injection needle should be used.

Figure A: Pack size of 1 vial + 1 filter needle + 1 syringe + 1 injection needle








Figure B: Pack size of 1 vial + 1 filter needle




Instructions for use

<p>1. Remove the plastic cap and disinfect the outer part of the rubber stopper of the vial.</p>	 The first drawing shows a hand holding a vial with the other hand removing a plastic cap. The second drawing shows a hand holding a vial with the other hand applying a disinfectant swab to the rubber stopper.
<p>2. The 18 G × 1½-inch, 5-micron, filter needle is used for withdrawal of drug product from the vial.</p> <p>Remove the 18 G × 1½-inch, 5-micron, filter needle and the 1 mL syringe from their packaging. Attach the filter needle to the syringe by twisting it onto the Luer lock syringe tip.</p>	 A line drawing showing a hand holding a syringe and another hand twisting a filter needle onto the Luer lock tip. A large black arrow points downwards from the top of the syringe.
<p>3. Push the filter needle into the centre of the vial stopper until the needle is completely inserted into the vial and the tip touches the bottom or bottom edge of the vial.</p> <p>4. Using aseptic technique withdraw all of the Yesafili vial contents into the syringe, keeping the vial in an upright position, slightly inclined to ease complete withdrawal. To deter the introduction of air, ensure the bevel of the filter needle is submerged into the liquid. Continue to tilt the vial during withdrawal keeping the bevel of the filter needle submerged in the liquid.</p>	 The top drawing shows a hand holding a syringe with the filter needle inserted into the vial stopper. The bottom drawing shows a hand tilting the vial to withdraw liquid into the syringe. Labels 'Solution' and 'Needle Bevel Pointing Down' are present with arrows pointing to the liquid and the bevel respectively.
<p>5. Ensure that the plunger rod is drawn sufficiently back when emptying the vial in order to completely empty the filter needle.</p>	
<p>6. Remove the filter needle and properly dispose of it.</p> <p>Note: Filter needle is not to be used for intravitreal injection.</p>	


7. Ensure the 30 G \times 1/2-inch injection needle is used for the intravitreal injection.

Remove the 30 G \times 1/2-inch injection needle from its packaging and using aseptic technique, firmly twist onto the Luer-lock syringe tip.





8. When ready to administer, remove the plastic needle shield from the needle.

Holding the syringe with the needle pointing up, check the syringe for bubbles. If there are bubbles, gently tap the syringe with your finger until the bubbles rise to the top.



9. Eliminate all bubbles and expel excess medicinal product by slowly depressing the plunger so that the flat plunger edge aligns with the line that marks 0.05 mL on the syringe.



10. The vial is for single use only. Extraction of multiple doses from a single vial may increase the risk of contamination and subsequent infection.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.