

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Zeffix 100 mg film-coated tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 100 mg lamivudine

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Film-coated tablet (tablet)

Butterscotch coloured, film-coated, capsule shaped, biconvex, approximate dimensions 11mm x 5mm and engraved "GX CG5" on one face.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Zeffix is indicated for the treatment of chronic hepatitis B in adults with:

- compensated liver disease with evidence of active viral replication, persistently elevated serum alanine aminotransferase (ALT) levels and histological evidence of active liver inflammation and/or fibrosis. Initiation of lamivudine treatment should only be considered when the use of an alternative antiviral agent with a higher genetic barrier is not available or appropriate (see section 5.1).
- decompensated liver disease in combination with a second agent without cross-resistance to lamivudine (see section 4.2).

4.2 Posology and method of administration

Therapy with Zeffix should be initiated by a physician experienced in the management of chronic hepatitis B.

Posology

Adults

The recommended dosage of Zeffix is 100 mg once daily.

In patients with decompensated liver disease, lamivudine should always be used in combination with a second agent, without cross-resistance to lamivudine, to reduce the risk of resistance and to achieve rapid viral suppression.

Duration of treatment

The optimal duration of treatment is unknown.

- In patients with HBeAg positive chronic hepatitis B (CHB) without cirrhosis, treatment should be administered for at least 6-12 months after HBeAg seroconversion (HBeAg and HBV DNA loss with HBeAb detection) is confirmed, to limit the risk of virological relapse, or until HBsAg seroconversion or there is loss of efficacy (see section 4.4). Serum ALT and HBV DNA levels

should be followed regularly after treatment discontinuation to detect any late virological relapse.

- In patients with HBeAg negative CHB (pre-core mutant) without cirrhosis, treatment should be administered at least until HBs seroconversion or there is evidence of loss of efficacy. With prolonged treatment, regular reassessment is recommended to confirm that continuation of the selected therapy remains appropriate for the patient.
- In patients with decompensated liver disease or cirrhosis and in liver transplant recipients, treatment cessation is not recommended (see section 5.1).

If lamivudine is discontinued, patients should be periodically monitored for evidence of recurrent hepatitis (see section 4.4).

Clinical resistance

In patients with either HBeAg positive or HBeAg negative CHB, the development of YMDD (tyrosine-methionine-aspartate-aspartate) mutant HBV may result in a diminished therapeutic response to lamivudine, indicated by a rise in HBV DNA and ALT from previous on-treatment levels. In order to reduce the risk of resistance in patients receiving lamivudine monotherapy, a switch to or addition of an alternative agent without cross-resistance to lamivudine based on therapeutic guidelines should be considered if serum HBV DNA remains detectable at or beyond 24 weeks of treatment (see section 5.1).

Special populations

Renal impairment

Lamivudine serum concentrations (AUC) are increased in patients with moderate to severe renal impairment due to decreased renal clearance. The dosage should, therefore, be reduced for patients with a creatinine clearance of < 50 ml/minute. When doses below 100 mg are required Zeffix oral solution should be used (see Table 1 below).

Table 1: Dosage of Zeffix in patients with decreased renal clearance.

Creatinine clearance ml/min	First dose of Zeffix oral solution *	Maintenance dose once daily
30 to < 50	20 ml (100 mg)	10 ml (50 mg)
15 to < 30	20 ml (100 mg)	5 ml (25 mg)
5 to < 15	7 ml (35 mg)	3 ml (15 mg)
< 5	7 ml (35 mg)	2 ml (10 mg)

* Zeffix oral solution containing 5 mg/ml lamivudine.

Data available in patients undergoing intermittent haemodialysis (for less than or equal to 4 hrs dialysis 2-3 times weekly), indicate that following the initial dosage reduction of lamivudine to correct for the patient's creatinine clearance, no further dosage adjustments are required while undergoing dialysis.

Hepatic impairment

Data obtained in patients with hepatic impairment, including those with end-stage liver disease awaiting transplant, show that lamivudine pharmacokinetics are not significantly affected by hepatic dysfunction. Based on these data, no dose adjustment is necessary in patients with hepatic impairment unless accompanied by renal impairment.

HIV co-infection

For the treatment of patients who are co-infected with HIV and are currently receiving or plan to receive combined antiretroviral treatment including lamivudine, the dose of lamivudine prescribed for HIV infection (usually 150 mg/twice daily in combination with other antiretrovirals) should be used.

Elderly

In elderly patients, normal ageing with accompanying renal decline has no clinically significant effect on lamivudine exposure, except in patients with creatinine clearance of < 50 ml/min.

Paediatric population

The safety and efficacy of Zeffix in infants, children and adolescents aged below 18 years have not been established. Currently available data are described in sections 4.4 and 5.1 but no recommendation on a posology can be made.

Method of administration

Oral use.

Zeffix can be taken with or without food.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

Exacerbations of hepatitis

Exacerbations on treatment

Spontaneous exacerbations in chronic hepatitis B are relatively common and are characterised by transient increases in serum ALT. After initiating antiviral therapy, serum ALT may increase in some patients as serum HBV DNA levels decline. In patients with compensated liver disease, these increases in serum ALT were generally not accompanied by an increase in serum bilirubin concentrations or signs of hepatic decompensation.

HBV viral subpopulations with reduced susceptibility to lamivudine (YMDD mutant HBV) have been identified with extended therapy. In some patients the development of YMDD mutant HBV can lead to exacerbation of hepatitis, primarily detected by serum ALT elevations and re-emergence of HBV DNA (see section 4.2). In patients who have YMDD mutant HBV, a switch to or addition of an alternative agent without cross resistance to lamivudine based on therapeutic guidelines should be considered (see section 5.1).

Exacerbations after treatment discontinuation

Acute exacerbation of hepatitis has been observed in patients who have discontinued hepatitis B therapy and is usually detected by serum ALT elevations and re-emergence of HBV DNA. In the controlled Phase III trials with no-active-treatment follow-up, the incidence of post-treatment ALT elevations (more than 3 times baseline) was higher in lamivudine-treated patients (21%) compared with those receiving placebo (8%). However, the proportion of patients who had post-treatment elevations associated with bilirubin elevations was low and similar in both treatment arms (see Table 3 in section 5.1). For lamivudine-treated patients, the majority of post-treatment ALT elevations occurred between 8 and 12 weeks post-treatment. Most events have been self-limiting, however some fatalities have been observed. If Zeffix is discontinued, patients should be periodically monitored both clinically and by assessment of serum liver function tests (ALT and bilirubin levels), for at least four months, and then as clinically indicated.

Exacerbations in patients with decompensated cirrhosis

Transplantation recipients and patients with decompensated cirrhosis are at greater risk from active viral replication. Due to the marginal liver function in these patients, hepatitis reactivation at

discontinuation of lamivudine or loss of efficacy during treatment may induce severe and even fatal decompensation. These patients should be monitored for clinical, virological and serological parameters associated with hepatitis B, liver and renal function, and antiviral response during treatment (at least every month), and, if treatment is discontinued for any reason, for at least 6 months after treatment. Laboratory parameters to be monitored should include (as a minimum) serum ALT, bilirubin, albumin, blood urea nitrogen, creatinine, and virological status: HBV antigen/antibody, and serum HBV DNA concentrations when possible. Patients experiencing signs of hepatic insufficiency during or post-treatment should be monitored more frequently as appropriate.

For patients who develop evidence of recurrent hepatitis post-treatment, there are insufficient data on the benefits of re-initiation of lamivudine treatment.

Mitochondrial dysfunction

Nucleoside and nucleotide analogues have been demonstrated *in vitro* and *in vivo* to cause a variable degree of mitochondrial damage. There have been reports of mitochondrial dysfunction in infants exposed *in utero* and/or post-natally to nucleoside analogues. The main adverse events reported are haematological disorders (anaemia, neutropenia), metabolic disorders (hyperlipasemia). Some late-onset neurological disorders have been reported (hypertonia, convulsion, abnormal behaviour). The neurological disorders might be transient or permanent. Any child exposed *in utero* to nucleoside and nucleotide analogues, should have clinical and laboratory follow-up and should be fully investigated for possible mitochondrial dysfunction in cases which have relevant signs or symptoms.

Paediatric patients

Lamivudine has been administered to children (2 years and above) and adolescents with compensated chronic hepatitis B. However, due to limitations of the data, the administration of lamivudine to this patient population is not currently recommended (see section 5.1).

Delta hepatitis or hepatitis C

The efficacy of lamivudine in patients co-infected with Delta hepatitis or hepatitis C has not been established and caution is advised.

Immunosuppressive treatments

Data are limited on the use of lamivudine in HBeAg negative (pre-core mutant) patients and in those receiving concurrent immunosuppressive regimes, including cancer chemotherapy. Lamivudine should be used with caution in these patients.

Monitoring

During treatment with Zeffix patients should be monitored regularly. Serum ALT and HBV DNA levels should be monitored at 3 month intervals and in HBeAg positive patients HBeAg should be assessed every 6 months.

HIV co-infection

For the treatment of patients who are co-infected with HIV and are currently receiving or plan to receive treatment with an antiretroviral combination regimen including lamivudine, the dose of lamivudine prescribed for HIV infection (usually 150 mg/twice daily in combination with other antiretrovirals) should be used.

The 100 mg usual dose of lamivudine used for the treatment of HBV is not appropriate for patients who acquire HIV or are co-infected with HBV and HIV. If a patient with unrecognised or untreated HIV infection is prescribed the dose of lamivudine recommended for the treatment of HBV, rapid emergence of HIV resistance and a limitation of treatment options is likely to result because of the

subtherapeutic dose and the inappropriate use of monotherapy for HIV treatment. HIV counselling and testing should be offered to all patients before beginning treatment with lamivudine for HBV and periodically during treatment.

Transmission of hepatitis B

There is no information available on maternal-foetal transmission of hepatitis B virus in pregnant women receiving treatment with lamivudine. The standard recommended procedures for hepatitis B virus immunisation in infants should be followed.

Patients should be advised that therapy with lamivudine has not been proven to reduce the risk of transmission of hepatitis B virus to others and therefore, appropriate precautions should still be taken.

Interactions with other medicinal products

Zeffix should not be taken with any other medicinal products containing lamivudine or medicinal products containing emtricitabine (see section 4.5).

The combination of lamivudine with cladribine is not recommended (see section 4.5).

Excipients

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

4.5 Interaction with other medicinal products and other forms of interaction

Interaction studies have only been performed in adults.

The likelihood of metabolic interactions is low due to limited metabolism and plasma protein binding and almost complete renal elimination of unchanged substance.

Lamivudine is predominantly eliminated by active organic cationic secretion. The possibility of interactions with other medicinal products administered concurrently should be considered, particularly when their main route of elimination is active renal secretion via the organic cationic transport system e.g. trimethoprim. Other medicinal products (e.g. ranitidine, cimetidine) are eliminated only in part by this mechanism and were shown not to interact with lamivudine.

Substances shown to be predominately excreted either via the active organic anionic pathway, or by glomerular filtration are unlikely to yield clinically significant interactions with lamivudine. Administration of trimethoprim/sulphamethoxazole 160 mg/800 mg increased lamivudine exposure by about 40 %. Lamivudine had no effect on the pharmacokinetics of trimethoprim or sulphamethoxazole. However, unless the patient has renal impairment, no dosage adjustment of lamivudine is necessary.

A modest increase in C_{max} (28 %) was observed for zidovudine when administered with lamivudine, however overall exposure (AUC) was not significantly altered. Zidovudine had no effect on the pharmacokinetics of lamivudine (see section 5.2).

Lamivudine has no pharmacokinetic interaction with alpha-interferon when the two medicinal products are concurrently administered. There were no observed clinically significant adverse interactions in patients taking lamivudine concurrently with commonly used immunosuppressant medicinal products (e.g. cyclosporin A). However, formal interaction studies have not been performed.

Emtricitabine

Due to similarities, Zeffix should not be administered concomitantly with other cytidine analogues, such as emtricitabine. Moreover, Zeffix should not be taken with any other medicinal products containing lamivudine (see section 4.4).

Cladribine

In vitro lamivudine inhibits the intracellular phosphorylation of cladribine leading to a potential risk of cladribine loss of efficacy in case of combination in the clinical setting. Some clinical findings also support a possible interaction between lamivudine and cladribine. Therefore, the concomitant use of lamivudine with cladribine is not recommended (see section 4.4).

Sorbitol

Co-administration of sorbitol solution (3.2 g, 10.2 g, 13.4 g) with a single 300 mg dose (Adult HIV daily dose) of lamivudine oral solution resulted in dose-dependent decreases of 14%, 32%, and 36% in lamivudine exposure (AUC_{∞}) and 28%, 52%, and 55% in the C_{max} of lamivudine in adults. When possible, avoid chronic co-administration of Zeffix with medicinal products containing sorbitol or other osmotic acting poly-alcohols or monosaccharide alcohols (e.g. xylitol, mannitol, lactitol, maltitol). Consider more frequent monitoring of HBV viral load when chronic co-administration cannot be avoided.

4.6 Fertility, pregnancy and lactation

Pregnancy

Animal studies with lamivudine showed an increase in early embryonic deaths in rabbits but not in rats (see section 5.3). Placental transfer of lamivudine has been shown to occur in humans.

Available human data from the Antiretroviral Pregnancy Registry reporting more than 1000 outcomes from first trimester and more than 1000 outcomes from second and third trimester exposure in pregnant women indicate no malformative and foeto/neonatal effect. Less than 1% of these women have been treated for HBV, whereas the majority was treated for HIV at higher doses and with other concomitant medications. Zeffix can be used during pregnancy if clinically needed.

For patients who are being treated with lamivudine and subsequently become pregnant consideration should be given to the possibility of a recurrence of hepatitis on discontinuation of lamivudine.

Breast-feeding

Based on more than 200 mother/child pairs treated for HIV, serum concentrations of lamivudine in breastfed infants of mothers treated for HIV are very low (less than 4% of maternal serum concentrations) and progressively decrease to undetectable levels when breastfed infants reach 24 weeks of age. The total amount of lamivudine ingested by a breastfed infant is very low and is therefore likely to result in exposures exerting a sub-optimal antiviral effect. Maternal hepatitis B is not a contraindication to breast-feeding if the newborn is adequately managed for hepatitis B prevention at birth, and there is no evidence that the low concentration of lamivudine in human milk leads to adverse reactions in breastfed infants. Therefore, breastfeeding may be considered in breast-feeding mothers being treated with lamivudine for HBV taking into account the benefit of breast feeding for the child and the benefit of therapy for the woman. Where there is maternal transmission of HBV, despite adequate prophylaxis, consideration should be given to discontinuing breastfeeding to reduce the risk of the emergence of lamivudine resistant mutants in the infant.

Fertility

Reproductive studies in animals have shown no effect on male or female fertility (see section 5.3).

Mitochondrial dysfunction

Nucleoside and nucleotide analogues have been demonstrated *in vitro* and *in vivo* to cause a variable degree of mitochondrial damage. There have been reports of mitochondrial dysfunction in infants exposed in utero and/or post-natally to nucleoside analogues (see section 4.4).

4.7 Effects on ability to drive and use machines

Patients should be informed that malaise and fatigue have been reported during treatment with lamivudine. The clinical status of the patient and the adverse reaction profile of lamivudine should be borne in mind when considering the patient's ability to drive or operate machinery.

4.8 Undesirable effects

Summary of the safety profile

The incidence of adverse reactions and laboratory abnormalities (with the exception of elevations of ALT and CPK, see below) were similar between placebo and lamivudine treated patients. The most common adverse reactions reported were malaise and fatigue, respiratory tract infections, throat and tonsil discomfort, headache, abdominal discomfort and pain, nausea, vomiting and diarrhoea.

Tabulated list of adverse reactions

Adverse reactions are listed below by system organ class and frequency. Frequency categories are only assigned to those adverse reactions considered to be at least possibly causally related to lamivudine. Frequencies are defined as: very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1000$), very rare ($< 1/10,000$) and not known (cannot be estimated from the available data).

The frequency categories assigned to the adverse reactions are mainly based on experience from clinical trials including a total of 1,171 patients with chronic hepatitis B receiving lamivudine at 100 mg.

Blood and lymphatic system disorders	
Not known	Thrombocytopenia
Metabolism and nutrition disorders	
Very rare	Lactic acidosis
Immune system disorders:	
Rare	Angioedema
Hepatobiliary disorders	
Very common	ALT elevations (see section 4.4)
Exacerbations of hepatitis, primarily detected by serum ALT elevations, have been reported 'on-treatment' and following lamivudine withdrawal. Most events have been self-limited, however fatalities have been observed very rarely (see section 4.4).	
Skin and subcutaneous tissue disorders	
Common	Rash, pruritus
Musculoskeletal and connective tissue disorders	
Common	Elevations of CPK
Common	Muscle disorders, including myalgia and cramps*
Not known	Rhabdomyolysis

* In Phase III studies frequency observed in the lamivudine treatment group was not greater than observed in the placebo group

Paediatric population

Based on limited data in children aged 2 to 17 years, there were no new safety issues identified compared to adults.

Other special populations

In patients with HIV infection, cases of pancreatitis and peripheral neuropathy (or paresthesia) have been reported. In patients with chronic hepatitis B there was no observed difference in incidence of these events between placebo and lamivudine treated patients.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via [the national reporting system listed in Appendix V](#).

4.9 Overdose

No specific signs or symptoms have been identified following acute overdose with lamivudine, apart from those listed as adverse reactions.

If overdose occurs the patient should be monitored and standard supportive treatment applied as required. Since lamivudine is dialysable, continuous haemodialysis could be used in the treatment of overdose, although this has not been studied.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group - Antivirals for systemic use, nucleoside and nucleotide reverse transcriptase inhibitors, ATC Code: J05AF05.

Mechanism of action

Lamivudine is an antiviral agent which is active against hepatitis B virus in all cell lines tested and in experimentally infected animals.

Lamivudine is metabolised by both infected and uninfected cells to the triphosphate (TP) derivative which is the active form of the parent compound. The intracellular half-life of the triphosphate in hepatocytes is 17-19 hours *in vitro*. Lamivudine-TP acts as a substrate for the HBV viral polymerase.

The formation of further viral DNA is blocked by incorporation of lamivudine-TP into the chain and subsequent chain termination.

Lamivudine-TP does not interfere with normal cellular deoxynucleotide metabolism. It is also only a weak inhibitor of mammalian DNA polymerases alpha and beta. Furthermore, lamivudine-TP has little effect on mammalian cell DNA content.

In assays relating to potential substance effects on mitochondrial structure and DNA content and function, lamivudine lacked appreciable toxic effects. It has a very low potential to decrease mitochondrial DNA content, is not permanently incorporated into mitochondrial DNA, and does not act as an inhibitor of mitochondrial DNA polymerase gamma.

Clinical efficacy and safety

Experience in patients with HBeAg positive CHB and compensated liver disease

In controlled studies, 1 year of lamivudine therapy significantly suppressed HBV DNA replication [34-57 % of patients were below the assay detection limits (Abbott Genostics solution hybridization assay, LLOD < 1.6pg/ml)], normalised ALT level (40-72 % of patients), induced HBeAg seroconversion (HBeAg loss and HBeAb detection with HBV DNA loss [by conventional assay], 16-18 % of patients), improved histology (38-52 % of patients had a ≥ 2 point decrease in the Knodell Histologic Activity Index [HAI]) and reduced progression of fibrosis (in 3-17 % of patients) and progression to cirrhosis.

Continued lamivudine treatment for an additional 2 years in patients who had failed to achieve HBeAg seroconversion in the initial 1 year controlled studies resulted in further improvement in bridging fibrosis. In patients with YMDD mutant HBV, 41/82 (50 %) patients had improvement in liver inflammation and 40/56 (71 %) patients without YMDD mutant HBV had improvement. Improvement in bridging fibrosis occurred in 19/30 (63 %) patients without YMDD mutant and 22/44 (50 %) patients with the mutant. Five percent (3/56) of patients without the YMDD mutant and 13 % (11/82) of patients with YMDD mutant showed worsening in liver inflammation compared to pre-treatment. Progression to cirrhosis occurred in 4/68 (6 %) patients with the YMDD mutant, whereas no patients without the mutant progressed to cirrhosis.

In an extended treatment study in Asian patients (NUCB3018) the HBeAg seroconversion rate and ALT normalisation rate at the end of the 5 year treatment period was 48 % (28/58) and 47 % (15/32), respectively. HBeAg seroconversion was increased in patients with elevated ALT levels; 77 % (20/26) of patients with pre-treatment ALT > 2 x ULN seroconverted. At the end of 5 years, all patients had HBV DNA levels that were undetectable or lower than pre-treatment levels.

Further results from the trial by YMDD mutant status are summarised in Table 2.

Table 2: Efficacy results 5 years by YMDD status (Asian Study) NUCB3018	Subjects, % (no.)	
	YMDD¹	Non-YMDD¹
<u>YMDD mutant HBV status</u>		
<u>HBeAg seroconversion</u>		
- All patients	38 (15/40)	72 (13/18)
- Baseline ALT $\leq 1 \times$ ULN ²	9 (1/11)	33 (2/6)
- Baseline ALT > 2 x ULN	60 (9/15)	100 (11/11)
<u>Undetectable HBV DNA</u>		
- Baseline ³	5 (2/40)	6 (1/18)
- Week 260 ⁴		
negative	8 (2/25)	0
positive < baseline	92 (23/25)	100 (4/4)
positive > baseline	0	0
<u>ALT normalisation</u>		
- Baseline		
normal	28 (11/40)	33 (6/18)
above normal	73 (29/40)	67 (12/18)
- Week 260		

normal	46 (13/28)	50 (2/4)
above normal < baseline	21 (6/28)	0
above normal > baseline	32 (9/28)	50 (2/4)

- 1 Patients designated as YMDD mutant were those with $\geq 5\%$ YMDD mutant HBV at any annual time-point during the 5-year period. Patients categorised as non-YMDD mutant were those with $> 95\%$ wild-type HBV at all annual time-points during the 5-year study period
- 2 Upper limit of normal
- 3 Abbott Genostics solution hybridisation assay (LLOD < 1.6 pg/ml)
- 4 Chiron Quantiplex assay (LLOD 0.7 Meq/ml)

Comparative data according to YMDD status were also available for histological assessment but only up to three years. In patients with YMDD mutant HBV, 18/39 (46 %) had improvements in necroinflammatory activity and 9/39 (23 %) had worsening. In patients without the mutant, 20/27 (74 %) had improvements in necroinflammatory activity and 2/27 (7 %) had worsening.

Following HBeAg seroconversion, serologic response and clinical remission are generally durable after stopping lamivudine. However, relapse following seroconversion can occur. In a long-term follow-up study of patients who had previously seroconverted and discontinued lamivudine, late virological relapse occurred in 39 % of the subjects. Therefore, following HBeAg seroconversion, patients should be periodically monitored to determine that serologic and clinical responses are being maintained. In patients who do not maintain a sustained serological response, consideration should be given to retreatment with either lamivudine or an alternative antiviral agent for resumption of clinical control of HBV.

In patients followed for up to 16 weeks after discontinuation of treatment at one year, post-treatment ALT elevations were observed more frequently in patients who had received lamivudine than in patients who had received placebo. A comparison of post-treatment ALT elevations between weeks 52 and 68 in patients who discontinued lamivudine at week 52 and patients in the same studies who received placebo throughout the treatment course is shown in Table 3. The proportion of patients who had post-treatment ALT elevations in association with an increase in bilirubin levels was low and similar in patients receiving either lamivudine or placebo.

Table 3: Post-treatment ALT Elevations in 2 Placebo-Controlled Studies in Adults

Abnormal Value	Patients with ALT Elevation/ Patients with Observations*	
	Lamivudine	Placebo
ALT ≥ 2 x baseline value	37/137 (27 %)	22/116 (19 %)
ALT ≥ 3 x baseline value [†]	29/137 (21 %)	9/116 (8 %)
ALT ≥ 2 x baseline value and absolute ALT > 500 IU/l	21/137 (15 %)	8/116 (7 %)
ALT ≥ 2 x baseline value; and bilirubin > 2 x ULN and ≥ 2 x baseline value	1/137 (0.7 %)	1/116 (0.9 %)

*Each patient may be represented in one or more category.

[†]Comparable to a Grade 3 toxicity in accordance with modified WHO criteria.

ULN = Upper limit of normal.

Experience in patients with HBeAg negative CHB

Initial data indicate the efficacy of lamivudine in patients with HBeAg negative CHB is similar to patients with HBeAg positive CHB, with 71 % of patients having HBV DNA suppressed below the detection limit of the assay, 67 % ALT normalisation and 38 % with improvement in HAI after one year of treatment. When lamivudine was discontinued, the majority of patients (70 %) had a return of viral replication. Data is available from an extended treatment study in HBeAg negative patients (NUCAB3017) treated with lamivudine. After two years of treatment in this study, ALT normalisation and undetectable HBV DNA occurred in 30/69 (43 %) and 32/68 (47 %) patients respectively and improvement in necroinflammatory score in 18/49 (37 %) patients. In patients without YMDD mutant HBV, 14/22 (64 %) showed improvement in necroinflammatory score and 1/22 (5 %) patients

worsened compared to pre-treatment. In patients with the mutant, 4/26 (15 %) patients showed improvement in necroinflammatory score and 8/26 (31 %) patients worsened compared to pre-treatment. No patients in either group progressed to cirrhosis.

Frequency of emergence of YMDD mutant HBV and impact on the treatment response

Lamivudine monotherapy results in the selection of YMDD mutant HBV in approximately 24 % of patients following one year of therapy, increasing to 69 % following 5 years of therapy. Development of YMDD mutant HBV is associated with reduced treatment response in some patients, as evidenced by increased HBV DNA levels and ALT elevations from previous on-therapy levels, progression of signs and symptoms of hepatitis disease and/or worsening of hepatic necroinflammatory findings. Given the risk of YMDD mutant HBV, maintenance of lamivudine monotherapy is not appropriate in patients with detectable serum HBV DNA at or beyond 24 weeks of treatment (see section 4.4).

In a double-blind study in CHB patients with YMDD mutant HBV and compensated liver disease (NUC20904), with a reduced virological and biochemical response to lamivudine (n=95), the addition of adefovir dipivoxil 10 mg once daily to ongoing lamivudine 100 mg for 52 weeks resulted in a median decrease in HBV DNA of 4.6 log₁₀ copies/ml compared to a median increase of 0.3 log₁₀ copies/ml in those patients receiving lamivudine monotherapy. Normalisation of ALT levels occurred in 31 % (14/45) of patients receiving combined therapy versus 6 % (3/47) receiving lamivudine alone. Viral suppression was maintained (follow-on study NUC20917) with combined therapy during the second year of treatment to week 104 with patients having continued improvement in virologic and biochemical responses.

In a retrospective study to determine the factors associated with HBV DNA breakthrough, 159 Asian HBeAg-positive patients were treated with lamivudine and followed up for a median period of almost 30 months. Those with HBV DNA levels greater than 200 copies/mL at 6 months (24 weeks) of lamivudine therapy had a 60 % chance of developing the YMDD mutant compared with 8 % of those with HBV DNA levels less than 200 copies/mL at 24 weeks of lamivudine therapy. The risk for developing YMDD mutant was 63% versus 13% with a cut off of 1000 copies/ml (NUCB3009 and NUCB3018).

Experience in patients with decompensated liver disease

Placebo controlled studies have been regarded as inappropriate in patients with decompensated liver disease, and have not been undertaken. In non-controlled studies, where lamivudine was administered prior to and during transplantation, effective HBV DNA suppression and ALT normalisation was demonstrated. When lamivudine therapy was continued post transplantation there was reduced graft re-infection by HBV, increased HBsAg loss and on one-year survival rate of 76 – 100 %.

As anticipated due to the concomitant immunosuppression, the rate of emergence of YMDD mutant HBV after 52 weeks treatment was higher (36 % - 64 %) in the liver transplant population than in the immunocompetent CHB patients (14 % - 32 %).

Forty patients (HBeAg negative or HBeAg positive) with either decompensated liver disease or recurrent HBV following liver transplantation and YMDD mutant were enrolled into an open label arm of study NUC20904. Addition of 10 mg adefovir dipivoxil once daily to ongoing lamivudine 100 mg for 52 weeks resulted in a median decrease in HBV DNA of 4.6 log₁₀ copies/ml. Improvement in liver function was also seen after one year of therapy. This degree of viral suppression was maintained (follow-on study NUC20917) with combined therapy during the second year of treatment to week 104 and most patients had improved markers of liver function and continued to derive clinical benefit.

Experience in CHB patients with advanced fibrosis or cirrhosis

In a placebo-controlled study in 651 patients with clinically compensated chronic hepatitis B and histologically confirmed fibrosis or cirrhosis, lamivudine treatment (median duration 32 months) significantly reduced the rate of overall disease progression (34/436, 7.8 % for lamivudine versus 38/215, 17.7 % for placebo, p=0.001), demonstrated by a significant reduction in the proportion of patients having increased Child-Pugh scores (15/436, 3.4 % versus 19/215, 8.8 %, p=0.023) or

developing hepatocellular carcinoma (17/436, 3.9 % versus 16/215, 7.4 %, $p=0.047$). The rate of overall disease progression in the lamivudine group was higher for subjects with detectable YMDD mutant HBV DNA (23/209, 11 %) compared to those without detectable YMDD mutant HBV (11/221, 5 %). However, disease progression in YMDD subjects in the lamivudine group was lower than the disease progression in the placebo group (23/209, 11 % versus 38/214, 18 % respectively). Confirmed HBeAg seroconversion occurred in 47 % (118/252) of subjects treated with lamivudine and 93 % (320/345) of subjects receiving lamivudine became HBV DNA negative (VERSANT [version 1], bDNA assay, LLOD < 0.7 MEq/ml) during the study.

Experience in children and adolescents

Lamivudine has been administered to children and adolescents with compensated CHB in a placebo controlled study of 286 patients aged 2 to 17 years. This population primarily consisted of children with minimal hepatitis B. A dose of 3 mg/kg once daily (up to a maximum of 100 mg daily) was used in children aged 2 to 11 years and a dose of 100 mg once daily in adolescents aged 12 years and above. This dose needs to be further substantiated. The difference in the HBeAg seroconversion rates (HBeAg and HBV DNA loss with HBeAb detection) between placebo and lamivudine was not statistically significant in this population (rates after one year were 13 % (12/95) for placebo versus 22 % (42/191) for lamivudine; $p=0.057$). The incidence of YMDD mutant HBV was similar to that observed in adults, ranging from 19 % at week 52 up to 45 % in patients treated continuously for 24 months.

5.2 Pharmacokinetic properties

Absorption

Lamivudine is well absorbed from the gastrointestinal tract, and the bioavailability of oral lamivudine in adults is normally between 80 and 85 %. Following oral administration, the mean time (T_{max}) to maximal serum concentrations (C_{max}) is about an hour. At therapeutic dose levels i.e. 100 mg once daily, C_{max} is in the order of 1.1-1.5 $\mu\text{g/ml}$ and trough levels were 0.015-0.020 $\mu\text{g/ml}$.

Co-administration of lamivudine with food resulted in a delay of T_{max} and a lower C_{max} (decreased by up to 47 %). However, the extent (based on the AUC) of lamivudine absorbed was not influenced, therefore lamivudine can be administered with or without food.

Distribution

From intravenous studies the mean volume of distribution is 1.3 l/kg. Lamivudine exhibits linear pharmacokinetics over the therapeutic dose range and displays low plasma protein binding to albumin. Limited data shows lamivudine penetrates the central nervous system and reaches the cerebro-spinal fluid (CSF). The mean lamivudine CSF/serum concentration ratio 2-4 hours after oral administration was approximately 0.12.

Biotransformation

Lamivudine is predominately cleared by renal excretion of unchanged substance. The likelihood of metabolic substance interactions with lamivudine is low due to the small (5-10 %) extent of hepatic metabolism and the low plasma protein binding.

Elimination

The mean systemic clearance of lamivudine is approximately 0.3 l/h/kg. The observed half-life of elimination is 18 to 19 hours. The majority of lamivudine is excreted unchanged in the urine via glomerular filtration and active secretion (organic cationic transport system). Renal clearance accounts for about 70 % of lamivudine elimination.

Special populations

Studies in patients with renal impairment show lamivudine elimination is affected by renal dysfunction. Dose reduction in patients with a creatinine clearance of < 50 ml/min is necessary (see section 4.2).

The pharmacokinetics of lamivudine are unaffected by hepatic impairment. Limited data in patients undergoing liver transplantation show that impairment of hepatic function does not impact significantly on the pharmacokinetics of lamivudine unless accompanied by renal dysfunction.

In elderly patients the pharmacokinetic profile of lamivudine suggests that normal ageing with accompanying renal decline has no clinically significant effect on lamivudine exposure, except in patients with creatinine clearance of < 50 ml/min (see section 4.2).

5.3 Preclinical safety data

Administration of lamivudine in animal toxicity studies at high doses was not associated with any major organ toxicity. At the highest dosage levels, minor effects on indicators of liver and kidney function were seen together with occasional reduction in liver weights. Reduction of erythrocytes and neutrophil counts were identified as the effects most likely to be of clinical relevance. These events were seen infrequently in clinical studies.

Lamivudine was not mutagenic in bacterial tests but, like many nucleoside analogues showed activity in an *in vitro* cytogenetic assay and the mouse lymphoma assay. Lamivudine was not genotoxic *in vivo* at doses that gave plasma concentrations around 60-70 times higher than the anticipated clinical plasma levels. As the *in vitro* mutagenic activity of lamivudine could not be confirmed by *in vivo* tests, it is concluded that lamivudine should not represent a genotoxic hazard to patients undergoing treatment.

Reproductive studies in animals have not shown evidence of teratogenicity and showed no effect on male or female fertility. Lamivudine induces early embryoletality when administered to pregnant rabbits at exposure levels comparable to those achieved in man, but not in the rat even at very high systemic exposures.

The results of long term carcinogenicity studies with lamivudine in rats and mice did not shown any carcinogenic potential.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet core

Microcrystalline cellulose
Sodium starch glycolate
Magnesium stearate

Tablet film coat

Hypromellose
Titanium dioxide
Macrogol 400
Polysorbate 80
Synthetic yellow and red iron oxides

6.2 Incompatibilities

Not applicable

6.3 Shelf life

3 years

6.4 Special precautions for storage

Do not store above 30°C.

6.5 Nature and contents of container

Double foil blisters, laminated with polyvinyl chloride.

Pack sizes of 28 or 84 film-coated tablets.

Not all pack-sizes may be marketed.

6.6 Special precautions for disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

GlaxoSmithKline Trading Services Limited
12 Riverwalk
Citywest Business Campus
Dublin 24
Ireland
D24 YK11

8. MARKETING AUTHORISATION NUMBERS

EU/1/99/114/001
EU/1/99/114/002

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 29 July 1999
Date of latest renewal: 23 June 2014

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>

1. NAME OF THE MEDICINAL PRODUCT

Zeffix 5 mg/ml oral solution

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ml of the oral solution contains 5 mg lamivudine

Excipients with known effect

Each 20 ml dose contains:

- 4 g of sucrose
- 30 mg of methyl parahydroxybenzoate (E218)
- 3.6 mg of propyl parahydroxybenzoate (E216)
- 400 mg of propylene glycol (E1520)
- 58.8 mg of sodium

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Oral solution

Clear, colourless to pale yellow in colour.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Zeffix is indicated for the treatment of chronic hepatitis B in adults with:

- compensated liver disease with evidence of active viral replication, persistently elevated serum alanine aminotransferase (ALT) levels and histological evidence of active liver inflammation and/or fibrosis. Initiation of lamivudine treatment should only be considered when the use of an alternative antiviral agent with a higher genetic barrier is not available or appropriate (see section 5.1).
- decompensated liver disease in combination with a second agent, without cross-resistance to lamivudine (see section 4.2).

4.2 Posology and method of administration

Therapy with Zeffix should be initiated by a physician experienced in the management of chronic hepatitis B.

Posology

Adults

The recommended dosage of Zeffix is 100 mg once daily.

In patients with decompensated liver disease, lamivudine should always be used in combination with a second agent, without cross-resistance to lamivudine, to reduce the risk of resistance and to achieve rapid viral suppression.

Duration of treatment

he optimal duration of treatment is unknown.

- In patients with HBeAg positive chronic hepatitis B (CHB) without cirrhosis, treatment should be administered for at least 6-12 months after HBeAg seroconversion (HBeAg and HBV DNA loss with HBeAb detection) is confirmed, to limit the risk of virological relapse, or until HBsAg seroconversion or there is loss of efficacy (see section 4.4). Serum ALT and HBV DNA levels should be followed regularly after treatment discontinuation to detect any late virological relapse.
- In patients with HBeAg negative CHB (pre-core mutant), without cirrhosis, treatment should be administered at least until HBs seroconversion or there is evidence of loss of efficacy. With prolonged treatment, regular reassessment is recommended to confirm that continuation of the selected therapy remains appropriate for the patient.
- In patients with decompensated liver disease or cirrhosis and in liver transplant recipients, treatment cessation is not recommended (see section 5.1).

If lamivudine is discontinued, patients should be periodically monitored for evidence of recurrent hepatitis (see section 4.4).

Clinical resistance

In patients with either HBeAg positive or HBeAg negative CHB, the development of YMDD (tyrosine-methionine-aspartate-aspartate) mutant HBV may result in a diminished therapeutic response to lamivudine, indicated by a rise in HBV DNA and ALT from previous on-treatment levels. In order to reduce the risk of resistance in patients receiving lamivudine monotherapy, a switch to or addition of an alternative agent without cross-resistance to lamivudine based on therapeutic guidelines should be considered if serum HBV DNA remains detectable at or beyond 24 weeks of treatment (see section 5.1).

Special populations

Renal impairment

Lamivudine serum concentrations (AUC) are increased in patients with moderate to severe renal impairment due to decreased renal clearance. The dosage should, therefore, be reduced for patients with a creatinine clearance of < 50 ml/minute. When doses below 100 mg are required Zeffix oral solution should be used (see Table 1 below).

Table 1: Dosage of Zeffix in patients with decreased renal clearance.

Creatinine clearance ml/min	First dose of Zeffix oral solution	Maintenance dose once daily
30 to < 50	20 ml (100 mg)	10 ml (50 mg)
15 to < 30	20 ml (100 mg)	5 ml (25 mg)
5 to < 15	7 ml (35 mg)	3 ml (15 mg)
< 5	7 ml (35 mg)	2 ml (10 mg)

Data available in patients undergoing intermittent haemodialysis (for less than or equal to 4 hrs dialysis 2-3 times weekly), indicate that following the initial dosage reduction of lamivudine to correct for the patient's creatinine clearance, no further dosage adjustments are required while undergoing dialysis.

Hepatic impairment

Data obtained in patients with hepatic impairment, including those with end-stage liver disease awaiting transplant, show that lamivudine pharmacokinetics are not significantly affected by hepatic dysfunction. Based on these data, no dose adjustment is necessary in patients with hepatic impairment unless accompanied by renal impairment.

HIV co-infection

For the treatment of patients who are co-infected with HIV and are currently receiving or plan to receive combined antiretroviral treatment including lamivudine, the dose of lamivudine prescribed for HIV infection (usually 150 mg/twice daily in combination with other antiretrovirals) should be used.

Elderly

In elderly patients, normal ageing with accompanying renal decline has no clinically significant effect on lamivudine exposure, except in patients with creatinine clearance of < 50 ml/min.

Paediatric population

The safety and efficacy of Zeffix in infants, children and adolescents aged below 18 years have not been established. Currently available data are described in sections 4.4 and 5.1 but no recommendation on a posology can be made.

Method of administration

Oral use.

Zeffix can be taken with or without food.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

Exacerbations of hepatitis

Exacerbations on treatment

Spontaneous exacerbations in chronic hepatitis B are relatively common and are characterised by transient increases in serum ALT. After initiating antiviral therapy, serum ALT may increase in some patients as serum HBV DNA levels decline. In patients with compensated liver disease, these increases in serum ALT were generally not accompanied by an increase in serum bilirubin concentrations or signs of hepatic decompensation.

HBV viral subpopulations with reduced susceptibility to lamivudine (YMDD mutant HBV) have been identified with extended therapy. In some patients the development of YMDD mutant HBV can lead to exacerbation of hepatitis, primarily detected by serum ALT elevations and re-emergence of HBV DNA (see section 4.2). In patients who have YMDD mutant HBV, a switch to or addition of an alternative agent without cross-resistance to lamivudine based on therapeutic guideline should be considered (see section 5.1).

Exacerbations after treatment discontinuation

Acute exacerbation of hepatitis has been observed in patients who have discontinued hepatitis B therapy and is usually detected by serum ALT elevations and re-emergence of HBV DNA. In the controlled Phase III trials with no-active-treatment follow-up, the incidence of post-treatment ALT elevations (more than 3 times baseline) was higher in lamivudine-treated patients (21%) compared with those receiving placebo (8%). However, the proportion of patients who had post-treatment elevations associated with bilirubin elevations was low and similar in both treatment arms (see Table 3 in section 5.1). For lamivudine-treated patients, the majority of post-treatment ALT elevations occurred between 8 and 12 weeks post-treatment. Most events have been self-limiting, however some fatalities have been observed. If Zeffix is discontinued, patients should be periodically monitored both

clinically and by assessment of serum liver function tests (ALT and bilirubin levels), for at least four months, and then as clinically indicated.

Exacerbations in patients with decompensated cirrhosis

Transplantation recipients and patients with decompensated cirrhosis are at greater risk from active viral replication. Due to the marginal liver function in these patients, hepatitis reactivation at discontinuation of lamivudine or loss of efficacy during treatment may induce severe and even fatal decompensation. These patients should be monitored for clinical, virological and serological parameters associated with hepatitis B, liver and renal function, and antiviral response during treatment (at least every month), and, if treatment is discontinued for any reason, for at least 6 months after treatment. Laboratory parameters to be monitored should include (as a minimum) serum ALT, bilirubin, albumin, blood urea nitrogen, creatinine, and virological status: HBV antigen/antibody, and serum HBV DNA concentrations when possible. Patients experiencing signs of hepatic insufficiency during or post-treatment should be monitored more frequently as appropriate.

For patients who develop evidence of recurrent hepatitis post-treatment, there are insufficient data on the benefits of re-initiation of lamivudine treatment.

Mitochondrial dysfunction

Nucleoside and nucleotide analogues have been demonstrated *in vitro* and *in vivo* to cause a variable degree of mitochondrial damage. There have been reports of mitochondrial dysfunction in infants exposed *in utero* and/or post-natally to nucleoside analogues. The main adverse events reported are haematological disorders (anaemia, neutropenia), metabolic disorders (hyperlipasemia). Some late-onset neurological disorders have been reported (hypertonia, convulsion, abnormal behaviour). The neurological disorders might be transient or permanent. Any child exposed *in utero* to nucleoside and nucleotide analogues, should have clinical and laboratory follow-up and should be fully investigated for possible mitochondrial dysfunction in cases which have relevant signs or symptoms.

Paediatric patients

Lamivudine has been administered to children (2 years and above) and adolescents with compensated chronic hepatitis B. However, due to limitations of the data, the administration of lamivudine to this patient population is not currently recommended (see section 5.1).

Delta hepatitis or hepatitis C

The efficacy of lamivudine in patients co-infected with Delta hepatitis or hepatitis C has not been established and caution is advised.

Immunosuppressive treatments

Data are limited on the use of lamivudine in HBeAg negative (pre-core mutant) patients and in those receiving concurrent immunosuppressive regimes, including cancer chemotherapy. Lamivudine should be used with caution in these patients.

Monitoring

During treatment with Zeffix patients should be monitored regularly. Serum ALT and HBV DNA levels should be monitored at 3 month intervals and in HBeAg positive patients HBeAg should be assessed every 6 months.

HIV co-infection

For the treatment of patients who are co-infected with HIV and are currently receiving or plan to receive treatment with an antiretroviral combination regimen including lamivudine, the dose of lamivudine prescribed for HIV infection (usually 150 mg/twice daily in combination with other

antiretrovirals) should be used.

The 100 mg usual dose of lamivudine used for the treatment of HBV is not appropriate for patients who acquire HIV or are co-infected with HBV and HIV. If a patient with unrecognised or untreated HIV infection is prescribed the dose of lamivudine recommended for the treatment of HBV, rapid emergence of HIV resistance and a limitation of treatment options is likely to result because of the subtherapeutic dose and the inappropriate use of monotherapy for HIV treatment. HIV counselling and testing should be offered to all patients before beginning treatment with lamivudine for HBV and periodically during treatment.

Transmission of hepatitis B

There is no information available on maternal-foetal transmission of hepatitis B virus in pregnant women receiving treatment with lamivudine. The standard recommended procedures for hepatitis B virus immunisation in infants should be followed.

Patients should be advised that therapy with lamivudine has not been proven to reduce the risk of transmission of hepatitis B virus to others and therefore, appropriate precautions should still be taken.

Interactions with other medicinal products

Zeffix should not be taken with any other medicinal products containing lamivudine or medicinal products containing emtricitabine (see section 4.5).

The combination of lamivudine with cladribine is not recommended (see section 4.5).

Excipients

Sucrose

Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine. Diabetic patients should be advised that each dose of oral solution (100 mg = 20 ml) contains 4 g of sucrose.

Propyl parahydroxybenzoate and methyl parahydroxybenzoate

The oral solution contains propyl and methyl parahydroxybenzoate. These products may cause allergic reactions (possibly delayed).

Sodium

This medicinal product contains 58.8 mg sodium per 20 ml, equivalent to 2.9% of the WHO recommended maximum daily intake of 2 g sodium for an adult.

4.5 Interaction with other medicinal products and other forms of interaction

Interaction studies have only been performed in adults.

The likelihood of metabolic interactions is low due to limited metabolism and plasma protein binding and almost complete renal elimination of unchanged substance.

Lamivudine is predominantly eliminated by active organic cationic secretion. The possibility of interactions with other medicinal products administered concurrently should be considered, particularly when their main route of elimination is active renal secretion via the organic cationic transport system e.g. trimethoprim. Other medicinal products (e.g. ranitidine, cimetidine) are eliminated only in part by this mechanism and were shown not to interact with lamivudine.

Substances shown to be predominately excreted either via the active organic anionic pathway, or by glomerular filtration are unlikely to yield clinically significant interactions with lamivudine.

Administration of trimethoprim/sulphamethoxazole 160 mg/800 mg increased lamivudine exposure by about 40 %. Lamivudine had no effect on the pharmacokinetics of trimethoprim or sulphamethoxazole. However, unless the patient has renal impairment, no dosage adjustment of lamivudine is necessary.

A modest increase in C_{max} (28 %) was observed for zidovudine when administered with lamivudine, however overall exposure (AUC) was not significantly altered. Zidovudine had no effect on the pharmacokinetics of lamivudine (see section 5.2).

Lamivudine has no pharmacokinetic interaction with alpha-interferon when the two medicinal products are concurrently administered. There were no observed clinically significant adverse interactions in patients taking lamivudine concurrently with commonly used immunosuppressant medicinal products (e.g. cyclosporin A). However, formal interaction studies have not been performed.

Emtricitabine

Due to similarities, Zeffix should not be administered concomitantly with other cytidine analogues, such as emtricitabine. Moreover, Zeffix should not be taken with any other medicinal products containing lamivudine (see section 4.4).

Cladribine

In vitro lamivudine inhibits the intracellular phosphorylation of cladribine leading to a potential risk of cladribine loss of efficacy in case of combination in the clinical setting. Some clinical findings also support a possible interaction between lamivudine and cladribine. Therefore, the concomitant use of lamivudine with cladribine is not recommended (see section 4.4).

Sorbitol

Co-administration of sorbitol solution (3.2 g, 10.2 g, 13.4 g) with a single 300 mg dose (Adult HIV daily dose) of lamivudine oral solution resulted in dose-dependent decreases of 14%, 32%, and 36% in lamivudine exposure (AUC_{∞}) and 28%, 52%, and 55% in the C_{max} of lamivudine in adults. When possible, avoid chronic co-administration of Zeffix with medicinal products containing sorbitol or other osmotic acting poly-alcohols or monosaccharide alcohols (e.g. xylitol, mannitol, lactitol, maltitol). Consider more frequent monitoring of HBV viral load when chronic co-administration cannot be avoided.

4.6 Fertility, pregnancy and lactation

Pregnancy

Animal studies with lamivudine showed an increase in early embryonic deaths in rabbits but not in rats (see section 5.3). Placental transfer of lamivudine has been shown to occur in humans.

Available human data from the Antiretroviral Pregnancy Registry reporting more than 1000 outcomes from first trimester and more than 1000 outcomes from second and third trimester exposure in pregnant women indicate no malformative and foeto/neonatal effect. Less than 1% of these women have been treated for HBV, whereas the majority was treated for HIV at higher doses and with other concomitant medications. Zeffix can be used during pregnancy if clinically needed.

For patients who are being treated with lamivudine and subsequently become pregnant consideration should be given to the possibility of a recurrence of hepatitis on discontinuation of lamivudine.

Breast-feeding

Based on more than 200 mother/child pairs treated for HIV, serum concentrations of lamivudine in breastfed infants of mothers treated for HIV are very low (less than 4% of maternal serum concentrations) and progressively decrease to undetectable levels when breastfed infants reach 24 weeks of age. The total amount of lamivudine ingested by a breastfed infant is very low and is therefore likely to result in exposures exerting a sub-optimal antiviral effect. Maternal hepatitis B is not a contraindication to breast-feeding if the newborn is adequately managed for hepatitis B prevention at birth, and there is no evidence that the low concentration of lamivudine in human milk leads to adverse reactions in breastfed infants. Therefore breastfeeding may be considered in breast-feeding mothers being treated with lamivudine for HBV taking into account the benefit of breast feeding for the child and the benefit of therapy for the woman. Where there is maternal transmission of HBV, despite adequate prophylaxis, consideration should be given to discontinuing breastfeeding to reduce the risk of the emergence of lamivudine resistant mutants in the infant.

Fertility

Reproductive studies in animals have shown no effect on male or female fertility (see section 5.3).

Mitochondrial dysfunction

Nucleoside and nucleotide analogues have been demonstrated *in vitro* and *in vivo* to cause a variable degree of mitochondrial damage. There have been reports of mitochondrial dysfunction in infants exposed in utero and/or post-natally to nucleoside analogues (see section 4.4).

4.7 Effects on ability to drive and use machines

Patients should be informed that malaise and fatigue have been reported during treatment with lamivudine. The clinical status of the patient and the adverse reaction profile of lamivudine should be borne in mind when considering the patient's ability to drive or operate machinery.

4.8 Undesirable effects

Summary of the safety profile

The incidence of adverse reactions and laboratory abnormalities (with the exception of elevations of ALT and CPK, see below) were similar between placebo and lamivudine treated patients. The most common adverse reactions reported were malaise and fatigue, respiratory tract infections, throat and tonsil discomfort, headache, abdominal discomfort and pain, nausea, vomiting and diarrhoea.

Tabulated list of adverse reactions

Adverse reactions are listed below by system organ class and frequency. Frequency categories are only assigned to those adverse reactions considered to be at least possibly causally related to lamivudine. Frequencies are defined as: very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1000$), very rare ($< 1/10,000$) and not known (cannot be estimated from the available data).

The frequency categories assigned to the adverse reactions are mainly based on experience from clinical trials including a total of 1,171 patients with chronic hepatitis B receiving lamivudine at 100 mg.

Blood and lymphatic system disorders	
Not known	Thrombocytopenia
Metabolism and nutrition disorders	
Very rare	Lactic acidosis

Immune system disorders:	
Rare	Angioedema
Hepatobiliary disorders	
Very common	ALT elevations (see section 4.4)
Exacerbations of hepatitis, primarily detected by serum ALT elevations, have been reported 'on-treatment' and following lamivudine withdrawal. Most events have been self-limited, however fatalities have been observed very rarely (see section 4.4).	
Skin and subcutaneous tissue disorders	
Common	Rash, pruritus
Musculoskeletal and connective tissue disorders	
Common	Elevations of CPK
Common	Muscle disorders, including myalgia and cramps*
Not known	Rhabdomyolysis

* In Phase III studies frequency observed in the lamivudine treatment group was not greater than observed in the placebo group

Paediatric population

Based on limited data in children aged 2 to 17 years, there were no new safety issues identified compared to adults.

Other special populations

In patients with HIV infection, cases of pancreatitis and peripheral neuropathy (or paresthesia) have been reported. In patients with chronic hepatitis B there was no observed difference in incidence of these events between placebo and lamivudine treated patients.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

4.9 Overdose

No specific signs or symptoms have been identified following acute overdose with lamivudine, apart from those listed as adverse reactions.

If overdose occurs the patient should be monitored, and standard supportive treatment applied as required. Since lamivudine is dialysable, continuous haemodialysis could be used in the treatment of overdose, although this has not been studied.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group - Antivirals for systemic use, nucleoside and nucleotide reverse transcriptase inhibitors, ATC Code: J05AF05.

Mechanism of action

Lamivudine is an antiviral agent which is active against hepatitis B virus in all cell lines tested and in experimentally infected animals.

Lamivudine is metabolised by both infected and uninfected cells to the triphosphate (TP) derivative which is the active form of the parent compound. The intracellular half-life of the triphosphate in hepatocytes is 17-19 hours *in vitro*. Lamivudine-TP acts as a substrate for the HBV viral polymerase.

The formation of further viral DNA is blocked by incorporation of lamivudine-TP into the chain and subsequent chain termination.

Lamivudine-TP does not interfere with normal cellular deoxynucleotide metabolism. It is also only a weak inhibitor of mammalian DNA polymerases alpha and beta. Furthermore, lamivudine-TP has little effect on mammalian cell DNA content.

In assays relating to potential substance effects on mitochondrial structure and DNA content and function, lamivudine lacked appreciable toxic effects. It has a very low potential to decrease mitochondrial DNA content, is not permanently incorporated into mitochondrial DNA, and does not act as an inhibitor of mitochondrial DNA polymerase gamma.

Clinical efficacy and safety

Experience in patients with HBeAg positive CHB and compensated liver disease

In controlled studies, 1 year of lamivudine therapy significantly suppressed HBV DNA replication (34-57 % of patients were below the assay detection limits (Abbott Genostics solution hybridization assay, LLOD < 1.6pg/ml)], normalised ALT level (40-72 % of patients), induced HBeAg seroconversion (HBeAg loss and HBeAb detection with HBV DNA loss [by conventional assay], 16-18 % of patients), improved histology (38-52 % of patients had a ≥ 2 point decrease in the Knodell Histologic Activity Index [HAI]) and reduced progression of fibrosis (in 3-17 % of patients) and progression to cirrhosis.

Continued lamivudine treatment for an additional 2 years in patients who had failed to achieve HBeAg seroconversion in the initial 1 year controlled studies resulted in further improvement in bridging fibrosis. In patients with YMDD mutant HBV, 41/82 (50 %) patients had improvement in liver inflammation and 40/56 (71 %) patients without YMDD mutant HBV had improvement. Improvement in bridging fibrosis occurred in 19/30 (63 %) patients without YMDD mutant and 22/44 (50 %) patients with the mutant. Five percent (3/56) of patients without the YMDD mutant and 13 % (11/82) of patients with YMDD mutant showed worsening in liver inflammation compared to pre-treatment. Progression to cirrhosis occurred in 4/68 (6 %) patients with the YMDD mutant, whereas no patients without the mutant progressed to cirrhosis.

In an extended treatment study in Asian patients (NUCB3018) the HBeAg seroconversion rate and ALT normalisation rate at the end of the 5 year treatment period was 48 % (28/58) and 47 % (15/32), respectively. HBeAg seroconversion was increased in patients with elevated ALT levels; 77 % (20/26) of patients with pre-treatment ALT > 2 x ULN seroconverted. At the end of 5 years, all patients had HBV DNA levels that were undetectable or lower than pre-treatment levels.

Further results from the trial by YMDD mutant status are summarised in Table 2.

Table 2: Efficacy results 5 years by YMDD status (Asian Study) NUCB3018

<i>YMDD mutant HBV status</i>	Subjects, % (no.)	
	YMDD ¹	Non-YMDD ¹
<u>HBeAg seroconversion</u>		
- All patients	38 (15/40)	72 (13/18)
- Baseline ALT ≤ 1 x ULN ²	9 (1/11)	33 (2/6)
- Baseline ALT > 2 x ULN	60 (9/15)	100 (11/11)
<u>Undetectable HBV DNA</u>		
- Baseline ³	5 (2/40)	6 (1/18)
- Week 260 ⁴		
negative	8 (2/25)	0
positive < baseline	92 (23/25)	100 (4/4)
positive > baseline	0	0
<u>ALT normalisation</u>		
- Baseline		
normal	28 (11/40)	33 (6/18)
above normal	73 (29/40)	67 (12/18)
- Week 260		
normal	46 (13/28)	50 (2/4)
above normal < baseline	21 (6/28)	0
above normal > baseline	32 (9/28)	50 (2/4)

1 Patients designated as YMDD mutant were those with ≥5% YMDD mutant HBV at any annual time-point during the 5-year period. Patients categorised as non-YMDD mutant were those with > 95% wild-type HBV at all annual time-points during the 5-year study period

2 Upper limit of normal

3 Abbott Genostics solution hybridisation assay (LLOD < 1.6 pg/ml)

4 Chiron Quantiplex assay (LLOD 0.7 Meq/ml)

Comparative data according to YMDD status were also available for histological assessment but only up to three years. In patients with YMDD mutant HBV, 18/39 (46 %) had improvements in necroinflammatory activity and 9/39 (23 %) had worsening. In patients without the mutant, 20/27 (74 %) had improvements in necroinflammatory activity and 2/27 (7 %) had worsening.

Following HBeAg seroconversion, serologic response and clinical remission are generally durable after stopping lamivudine. However, relapse following seroconversion can occur. In a long-term follow-up study of patients who had previously seroconverted and discontinued lamivudine, late virological relapse occurred in 39 % of the subjects. Therefore, following HBeAg seroconversion, patients should be periodically monitored to determine that serologic and clinical responses are being maintained. In patients who do not maintain a sustained serological response, consideration should be given to retreatment with either lamivudine or an alternative antiviral agent for resumption of clinical control of HBV.

In patients followed for up to 16 weeks after discontinuation of treatment at one year, post-treatment ALT elevations were observed more frequently in patients who had received lamivudine than in patients who had received placebo. A comparison of post-treatment ALT elevations between weeks 52 and 68 in patients who discontinued lamivudine at week 52 and patients in the same studies who received placebo throughout the treatment course is shown in Table 3. The proportion of patients who had post-treatment ALT elevations in association with an increase in bilirubin levels was low and similar in patients receiving either lamivudine or placebo.

Table 3: Post-treatment ALT Elevations in 2 Placebo-Controlled Studies in Adults

Abnormal Value	Patients with ALT Elevation/ Patients with Observations*	
	Lamivudine	Placebo
ALT \geq 2 x baseline value	37/137 (27 %)	22/116 (19 %)
ALT \geq 3 x baseline value [†]	29/137 (21 %)	9/116 (8 %)
ALT \geq 2 x baseline value and absolute ALT > 500 IU/l	21/137 (15 %)	8/116 (7 %)
ALT \geq 2 x baseline value; and bilirubin > 2 x ULN and \geq 2 x baseline value	1/137 (0.7 %)	1/116 (0.9 %)

*Each patient may be represented in one or more category.

[†]Comparable to a Grade 3 toxicity in accordance with modified WHO criteria.

ULN = Upper limit of normal.

Experience in patients with HBeAg negative CHB

Initial data indicate the efficacy of lamivudine in patients with HBeAg negative CHB is similar to patients with HBeAg positive CHB, with 71 % of patients having HBV DNA suppressed below the detection limit of the assay, 67 % ALT normalisation and 38 % with improvement in HAI after one year of treatment. When lamivudine was discontinued, the majority of patients (70 %) had a return of viral replication. Data is available from an extended treatment study in HBeAg negative patients (NUCAB3017) treated with lamivudine. After two years of treatment in this study, ALT normalisation and undetectable HBV DNA occurred in 30/69 (43 %) and 32/68 (47 %) patients respectively and improvement in necroinflammatory score in 18/49 (37 %) patients. In patients without YMDD mutant HBV, 14/22 (64 %) showed improvement in necroinflammatory score and 1/22 (5 %) patients worsened compared to pre-treatment. In patients with the mutant, 4/26 (15%) patients showed improvement in necroinflammatory score and 8/26 (31 %) patients worsened compared to pre-treatment. No patients in either group progressed to cirrhosis.

Frequency of emergence of YMDD mutant HBV and impact on the treatment response

Lamivudine monotherapy results in the selection of YMDD mutant HBV in approximately 24 % of patients following one year of therapy, increasing to 69 % following 5 years of therapy. Development of YMDD mutant HBV is associated with reduced treatment response in some patients, as evidenced by increased HBV DNA levels and ALT elevations from previous on-therapy levels, progression of signs and symptoms of hepatitis disease and/or worsening of hepatic necroinflammatory findings. Given the risk of YMDD mutant HBV, maintenance of lamivudine monotherapy is not appropriate in patients with detectable serum HBV DNA at or beyond 24 weeks of treatment (see section 4.4).

In a double-blind study in CHB patients with YMDD mutant HBV and compensated liver disease (NUC20904), with a reduced virological and biochemical response to lamivudine (n=95), the addition of adefovir dipivoxil 10 mg once daily to ongoing lamivudine 100 mg for 52 weeks resulted in a median decrease in HBV DNA of 4.6 log₁₀ copies/ml compared to a median increase of 0.3 log₁₀ copies/ml in those patients receiving lamivudine monotherapy. Normalisation of ALT levels occurred in 31 % (14/45) of patients receiving combined therapy versus 6 % (3/47) receiving lamivudine alone. Viral suppression was maintained (follow-on study NUC20917) with combined therapy during the second year of treatment to week 104 with patients having continued improvement in virologic and biochemical responses.

In a retrospective study to determine the factors associated with HBV DNA breakthrough, 159 Asian HBeAg-positive patients were treated with lamivudine and followed up for a median period of almost 30 months. Those with HBV DNA levels greater than 200 copies/mL at 6 months (24 weeks) of lamivudine therapy had a 60 % chance of developing the YMDD mutant compared with 8 % of those with HBV DNA levels less than 200 copies/mL at 24 weeks of lamivudine therapy. The risk for developing YMDD mutant was 63% versus 13% with a cut off of 1000 copies/ml (NUCB3009 and NUCB3018).

Experience in patients with decompensated liver disease

Placebo controlled studies have been regarded as inappropriate in patients with decompensated liver disease, and have not been undertaken. In non-controlled studies, where lamivudine was administered prior to and during transplantation, effective HBV DNA suppression and ALT normalisation was demonstrated. When lamivudine therapy was continued post transplantation there was reduced graft re-infection by HBV, increased HBsAg loss and on one-year survival rate of 76 – 100 %.

As anticipated due to the concomitant immunosuppression, the rate of emergence of YMDD mutant HBV after 52 weeks treatment was higher (36 % - 64 %) in the liver transplant population than in the immunocompetent CHB patients (14 % - 32 %).

Forty patients (HBeAg negative or HBeAg positive) with either decompensated liver disease or recurrent HBV following liver transplantation and YMDD mutant were enrolled into an open label arm of study NUC20904. Addition of 10 mg adefovir dipivoxil once daily to ongoing lamivudine 100 mg for 52 weeks resulted in a median decrease in HBV DNA of 4.6 log₁₀ copies/ml. Improvement in liver function was also seen after one year of therapy. This degree of viral suppression was maintained (follow-on study NUC20917) with combined therapy during the second year of treatment to week 104 and most patients had improved markers of liver function and continued to derive clinical benefit.

Experience in CHB patients with advanced fibrosis or cirrhosis

In a placebo-controlled study in 651 patients with clinically compensated chronic hepatitis B and histologically confirmed fibrosis or cirrhosis, lamivudine treatment (median duration 32 months) significantly reduced the rate of overall disease progression (34/436, 7.8 % for lamivudine versus 38/215, 17.7 % for placebo, p=0.001), demonstrated by a significant reduction in the proportion of patients having increased Child-Pugh scores (15/436, 3.4 % versus 19/215, 8.8 %, p=0.023) or developing hepatocellular carcinoma (17/436, 3.9 % versus 16/215, 7.4 %, p=0.047). The rate of overall disease progression in the lamivudine group was higher for subjects with detectable YMDD mutant HBV DNA (23/209, 11 %) compared to those without detectable YMDD mutant HBV (11/221, 5 %). However, disease progression in YMDD subjects in the lamivudine group was lower than the disease progression in the placebo group (23/209, 11 % versus 38/214, 18 % respectively). Confirmed HBeAg seroconversion occurred in 47 % (118/252) of subjects treated with lamivudine and 93 % (320/345) of subjects receiving lamivudine became HBV DNA negative (VERSANT [version 1], bDNA assay, LLOD < 0.7 MEq/ml) during the study.

Experience in children and adolescents

Lamivudine has been administered to children and adolescents with compensated CHB in a placebo controlled study of 286 patients aged 2 to 17 years. This population primarily consisted of children with minimal hepatitis B. A dose of 3 mg/kg once daily (up to a maximum of 100 mg daily) was used in children aged 2 to 11 years and a dose of 100 mg once daily in adolescents aged 12 years and above. This dose needs to be further substantiated. The difference in the HBeAg seroconversion rates (HBeAg and HBV DNA loss with HBeAb detection) between placebo and lamivudine was not statistically significant in this population (rates after one year were 13 % (12/95) for placebo versus 22 % (42/191) for lamivudine; p=0.057). The incidence of YMDD mutant HBV was similar to that observed in adults, ranging from 19 % at week 52 up to 45 % in patients treated continuously for 24 months.

5.2 Pharmacokinetic properties

Absorption

Lamivudine is well absorbed from the gastrointestinal tract, and the bioavailability of oral lamivudine in adults is normally between 80 and 85 %. Following oral administration, the mean time (T_{max}) to maximal serum concentrations (C_{max}) is about an hour. At therapeutic dose levels i.e., 100 mg once daily, C_{max} is in the order of 1.1-1.5 µg/ml and trough levels were 0.015-0.020 µg/ml.

Co-administration of lamivudine with food resulted in a delay of T_{max} and a lower C_{max} (decreased by up to 47 %). However, the extent (based on the AUC) of lamivudine absorbed was not influenced, therefore lamivudine can be administered with or without food.

Distribution

From intravenous studies the mean volume of distribution is 1.3 l/kg. Lamivudine exhibits linear pharmacokinetics over the therapeutic dose range and displays low plasma protein binding to albumin.

Limited data shows lamivudine penetrates the central nervous system and reaches the cerebro-spinal fluid (CSF). The mean lamivudine CSF/serum concentration ratio 2-4 hours after oral administration was approximately 0.12.

Biotransformation

Lamivudine is predominately cleared by renal excretion of unchanged substance. The likelihood of metabolic substance interactions with lamivudine is low due to the small (5-10 %) extent of hepatic metabolism and the low plasma protein binding.

Elimination

The mean systemic clearance of lamivudine is approximately 0.3 l/h/kg. The observed half-life of elimination is 18 to 19 hours. The majority of lamivudine is excreted unchanged in the urine via glomerular filtration and active secretion (organic cationic transport system). Renal clearance accounts for about 70 % of lamivudine elimination.

Special populations

Studies in patients with renal impairment show lamivudine elimination is affected by renal dysfunction. Dose reduction in patients with a creatinine clearance of < 50 ml/min is necessary (see section 4.2).

The pharmacokinetics of lamivudine are unaffected by hepatic impairment. Limited data in patients undergoing liver transplantation, show that impairment of hepatic function does not impact significantly on the pharmacokinetics of lamivudine unless accompanied by renal dysfunction.

In elderly patients the pharmacokinetic profile of lamivudine suggests that normal ageing with accompanying renal decline has no clinically significant effect on lamivudine exposure, except in patients with creatinine clearance of < 50 ml/min (see section 4.2).

5.3 Preclinical safety data

Administration of lamivudine in animal toxicity studies at high doses was not associated with any major organ toxicity. At the highest dosage levels, minor effects on indicators of liver and kidney function were seen together with occasional reduction in liver weights. Reduction of erythrocytes and neutrophil counts were identified as the effects most likely to be of clinical relevance. These events were seen infrequently in clinical studies.

Lamivudine was not mutagenic in bacterial tests but, like many nucleoside analogues showed activity in an *in vitro* cytogenetic assay and the mouse lymphoma assay. Lamivudine was not genotoxic *in vivo* at doses that gave plasma concentrations around 60-70 times higher than the anticipated clinical plasma levels. As the *in vitro* mutagenic activity of lamivudine could not be confirmed by *in vivo* tests, it is concluded that lamivudine should not represent a genotoxic hazard to patients undergoing treatment.

Reproductive studies in animals have not shown evidence of teratogenicity and showed no effect on male or female fertility. Lamivudine induces early embryoletality when administered to pregnant

rabbits at exposure levels comparable to those achieved in man, but not in the rat even at very high systemic exposures.

The results of long term carcinogenicity studies with lamivudine in rats and mice did not shown any carcinogenic potential.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sucrose (20 % w/v)
Methyl parahydroxybenzoate (E218)
Propyl parahydroxybenzoate (E216)
Citric acid (anhydrous)
Propylene glycol (E1520)
Sodium citrate
Artificial strawberry flavour
Artificial banana flavour
Purified water

6.2 Incompatibilities

Not applicable

6.3 Shelf life

2 years

After first opening: 1 month

6.4 Special precautions for storage

Do not store above 25 °C.

6.5 Nature and contents of container

Opaque, white, high-density polyethylene (HDPE) bottle with a polypropylene child-resistant closure. The pack also includes a polyethylene applicator-adapter, and a 10 ml oral applicator comprised of a polypropylene barrel (with ml graduations) and a polyethylene plunger. The oral applicator is provided for accurate measurement of the prescribed dose of oral solution.

Pack size of 240 ml lamivudine oral solution.

6.6 Special precautions for disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

GlaxoSmithKline Trading Services Limited
12 Riverwalk
Citywest Business Campus
Dublin 24
Ireland
D24 YK11

8. MARKETING AUTHORISATION NUMBER

EU/1/99/114/003

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 29 July 1999

Date of latest renewal: 23 June 2014

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>

ANNEX II

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturers responsible for batch release

Film-coated tablet:

Delpharm Poznań Spółka Akcyjna
ul. Grunwaldzka 189
60-322 Poznan
Poland

Oral solution:

GlaxoSmithKline Trading Services Limited
12 Riverwalk
Citywest Business Campus
Dublin 24
Ireland
D24 YK11

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to restricted medical prescription (See Annex I: Summary of Product Characteristics, section 4.2).

C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal products are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

- **Risk management plan (RMP)**

Not applicable.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

OUTER CARTON FOR TABLETS: 28 tablet box, 84 tablet box

1. NAME OF THE MEDICINAL PRODUCT

Zeffix 100 mg film-coated tablets
lamivudine

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each film-coated tablet contains 100 mg lamivudine

3. LIST OF EXCIPIENTS

4. PHARMACEUTICAL FORM AND CONTENTS

28 film-coated tablets
84 film-coated tablets

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use

Oral use

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 30°C

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

GlaxoSmithKline Trading Services Limited
12 Riverwalk
Citywest Business Campus
Dublin 24
Ireland
D24 YK11

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/99/114/001 28 tablets
EU/1/99/114/002 84 tablets

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

zeffix 100 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC:
SN:
NN:

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

28 tablet box, 84 tablet box

1. NAME OF THE MEDICINAL PRODUCT

Zeffix 100 mg tablets
lamivudine

2. NAME OF THE MARKETING AUTHORISATION HOLDER

GlaxoSmithKline Trading Services Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

OUTER CARTON FOR ORAL SOLUTION

1. NAME OF THE MEDICINAL PRODUCT

Zeffix 5 mg/ml oral solution
lamivudine

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each ml of the oral solution contains 5 mg lamivudine.

3. LIST OF EXCIPIENTS

Contains amongst others:
sugar (sucrose), sodium, propylene glycol (E1520), preservatives: methyl parahydroxybenzoate (E218) and propyl parahydroxybenzoate (E216)

4. PHARMACEUTICAL FORM AND CONTENTS

Each bottle contains 240 ml oral solution
The pack contains an oral applicator.

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use

Oral use

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

Discard one month after first opening

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

GlaxoSmithKline Trading Services Limited
12 Riverwalk
Citywest Business Campus
Dublin 24
Ireland
D24 YK11

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/99/114/003

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

zeffix 5 mg/ml

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC:
SN:
NN:

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING

BOTTLE LABEL FOR ORAL SOLUTION

1. NAME OF THE MEDICINAL PRODUCT

Zeffix 5 mg/ml oral solution
lamivudine

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each ml of the oral solution contains 5 mg lamivudine.

3. LIST OF EXCIPIENTS

Contains amongst others:
sugar (sucrose), sodium, propylene glycol (E1520), preservatives: methyl parahydroxybenzoate (E218) and propyl parahydroxybenzoate (E216)

4. PHARMACEUTICAL FORM AND CONTENTS

Each bottle contains 240 ml oral solution

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use

Oral use

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

Discard one month after first opening

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

GlaxoSmithKline Trading Services Limited
12 Riverwalk
Citywest Business Campus
Dublin 24
Ireland
D24 YK11

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/99/114/003

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

17. UNIQUE IDENTIFIER – 2D BARCODE

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

B. PACKAGE LEAFLET

Package leaflet: Information for the patient

Zeffix 100 mg film-coated tablets lamivudine

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Zeffix is and what it is used for
2. What you need to know before you take Zeffix
3. How to take Zeffix
4. Possible side effects
5. How to store Zeffix
6. Contents of the pack and other information

1. What Zeffix is and what it is used for

The active ingredient in Zeffix is lamivudine.

Zeffix is used to treat long term (chronic) hepatitis B infection in adults.

Zeffix is an antiviral medicine that suppresses the hepatitis B virus and belongs to a group of medicines called *nucleoside analogue reverse transcriptase inhibitors (NRTIs)*.

Hepatitis B is a virus which infects the liver, causes long term (chronic) infection, and can lead to liver damage. Zeffix can be used in people whose liver is damaged, but still functions normally (compensated liver disease) and in combination with other medicines in people whose liver is damaged and does not function normally (decompensated liver disease).

Treatment with Zeffix can reduce the amount of hepatitis B virus in your body. This should lead to a reduction in liver damage and an improvement in your liver function. Not everyone responds to treatment with Zeffix in the same way. Your doctor will monitor the effectiveness of your treatment with regular blood tests.

2. What you need to know before you take Zeffix

Your healthcare provider should offer you counselling and testing for HIV infection before you start treatment with lamivudine for hepatitis B infection and during treatment. If you have or get HIV infection, see section 3.

Do not take Zeffix

- if you're **allergic** to lamivudine or to any of the other ingredients of this medicine (*listed in Section 6*).

→ **Check with your doctor** if you think this applies to you.

Warnings and precautions

Some people taking Zeffix or other similar medicines are more at risk of serious side effects. You need to be aware of the extra risks:

- if you have ever had other types of **liver disease**, such as hepatitis C
- if you're seriously **overweight** (especially if you're a woman).

➔ **Talk to your doctor if any of these apply to you.** You may need extra check-ups, including blood tests, while you're taking your medication. **See Section 4** for more information about the risks.

Don't stop taking Zeffix without your doctor's advice, as there is a risk of your hepatitis getting worse. When you stop taking Zeffix your doctor will monitor you for at least four months to check for any problems. This will mean taking blood samples to check for any raised liver enzyme levels, which may indicate liver damage. See section 3 for more information about how to take Zeffix.

Protect other people

Hepatitis B infection is spread by sexual contact with someone who has the infection, or by transfer of infected blood (for example, by sharing injection needles). Zeffix will not stop you passing hepatitis B infection on to other people. To protect other people from becoming infected with hepatitis B:

- **Use a condom** when you have oral or penetrative sex.
- **Don't risk blood transfer** — for example, don't share needles.

Other medicines and Zeffix

Tell your doctor or pharmacist if you're taking, have recently taken or might take any other medicines, including herbal medicines or other medicines you bought without a prescription.

Remember to tell your doctor or pharmacist if you begin taking a new medicine while you're taking Zeffix.

These medicines should not be used with Zeffix:

- medicines (usually liquids) containing sorbitol and other sugar alcohols (such as xylitol, mannitol, lactitol or maltitol), if taken regularly
- other medicines containing lamivudine, used to treat **HIV infection** (sometimes called the AIDS virus)
- emtricitabine used to treat **HIV** or **hepatitis B infection**
- cladribine, used to treat **hairy cell leukaemia**

➔ **Tell your doctor** if you're being treated with any of these.

Pregnancy

If you are pregnant, think you may be pregnant or are planning to have a baby:

➔ **Talk to your doctor** about the risks and benefits of taking Zeffix during your pregnancy.

Do not stop treatment with Zeffix without your doctor's advice.

Breast-feeding

Zeffix can pass into breast-milk. If you are breast-feeding, or thinking about breast-feeding:

➔ **Talk to your doctor** before you take Zeffix.

Driving and using machines

Zeffix may make you feel tired, which could affect your ability to drive or use machines.

➔ Don't drive or use machines unless you are sure you're not affected.

Zeffix contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take Zeffix

Always take this medicine exactly as your doctor has told you to. Check with your doctor or pharmacist if you're not sure.

Stay in regular contact with your doctor

Zeffix helps to control your hepatitis B infection. You need to keep taking it every day to control your infection and stop your illness getting worse.

→ **Keep in touch with your doctor, and don't stop taking Zeffix** without your doctor's advice.

How much to take

The usual dose of Zeffix is one tablet (100 mg lamivudine) **once a day.**

Your doctor may prescribe a lower dose if you have problems with your kidneys. An oral solution of Zeffix is available for people who need a lower than usual dose, or who can't take tablets.

→ **Talk to your doctor** if this applies to you.

Patients who also have or may get HIV infection

If you have or get HIV that is not being treated with medicines while taking lamivudine for the treatment of hepatitis B infection, the HIV virus may develop resistance to certain HIV medicines and become harder to treat. Lamivudine can also be used to treat HIV infection. Talk to your doctor if you have HIV infection. Your doctor may treat you with another medicine that contains a higher dose of lamivudine, usually 150 mg twice a day, as the lower dose of 100 mg lamivudine is not enough to treat HIV infection. If you are planning to change your HIV treatment, discuss this change with your doctor first.

→ **Talk to your doctor** if this applies to you.

Swallow the tablet whole, with some water. Zeffix can be taken with or without food.

If you take more Zeffix than you should

If you accidentally take too much Zeffix, tell your doctor or your pharmacist, or contact your nearest hospital emergency department for further advice. If possible, show them the Zeffix pack.

If you forget to take Zeffix

If you forget to take a dose, take it as soon as you remember. Then continue your treatment as before. Don't take a double dose to make up for a missed dose.

Don't stop taking Zeffix

You must not stop taking Zeffix without consulting your doctor. There is a risk of your hepatitis getting worse (*see section 2*). When you stop taking Zeffix your doctor will monitor you for at least four months to check for any problems. This will mean taking blood samples to check for any raised liver enzyme levels, which may indicate liver damage.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Side effects that were commonly reported in Zeffix clinical trials were tiredness, respiratory tract infections, throat discomfort, headache, stomach discomfort and pain, nausea, vomiting and diarrhoea, increases in liver enzymes and enzymes produced in the muscles (*see below*).

Allergic reaction

These are rare (may affect up to 1 in 1,000 people). Signs include:

- swelling of eyelids, face or lips
- difficulty swallowing or breathing

➔ **Contact a doctor immediately** if you get these symptoms. **Stop taking Zeffix.**

Side effects thought to be caused by Zeffix

A very common side effect (these may affect **more than 1 in 10** people) which may show up in blood tests is:

- an increase in the level of some liver enzymes (*transaminases*), which may be a sign of inflammation or damage in the liver.

A common side effect (these may affect **up to 1 in 10** people) is:

- cramps and muscle pains
- skin rash or 'hives' anywhere on the body

A common side effect which may show up in blood tests is:

- an increase in the level of an enzyme produced in the muscles (*creatine phosphokinase*) which may be a sign that body tissue is damaged.

A very rare side effect (these may affect up to 1 in 10,000 people) is:

- lactic acidosis (excess lactic acid in the blood).

Other side effects

Other side effects have occurred in a very small number of people but their exact frequency is unknown

- breakdown of muscle tissue
- a worsening of liver disease after Zeffix is stopped or during treatment if the hepatitis B virus becomes resistant to Zeffix. This can be fatal in some people.

A side effect which may show up in blood tests is:

- a decrease in the number of cells involved in blood clotting (*thrombocytopenia*).

If you get any side effects

➔ **Talk to your doctor or pharmacist.** This includes any possible side effects not listed in this leaflet.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Zeffix

Keep this medicine out of the sight and reach of children.

Don't take this medicine after the expiry date shown on the carton and blister.

Don't store above 30 °C.

Do not throw away any medicines in your wastewater or household rubbish. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Zeffix contains

The active substance is lamivudine. Each film-coated tablet contains 100 mg of lamivudine.

The other ingredients are: microcrystalline cellulose, sodium starch glycolate, magnesium stearate, hypromellose, titanium dioxide, macrogol 400, polysorbate 80, synthetic yellow and red iron oxide.

What Zeffix looks like and contents of the pack

Zeffix film-coated tablets are supplied in tamper evident foil blisters containing 28 or 84 tablets. The tablets are butterscotch coloured, capsule shaped, biconvex and engraved "GX CG5" on one side.

Not all pack-sizes may be available in your country.

Manufacturer

Delpharm Poznań Spółka Akcyjna
ul. Grunwaldzka 189
60-322 Poznan
Poland

Marketing Authorisation Holder

GlaxoSmithKline Trading Services
Limited
12 Riverwalk
Citywest Business Campus
Dublin 24
Ireland
D24 YK11

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder:

België/Belgique/Belgien

GlaxoSmithKline Pharmaceuticals s.a./n.v.
Tél/Tel: + 32 (0)10 85 52 00

Lietuva

GlaxoSmithKline Trading Services Limited
Tel: + 370 80000334

България

GlaxoSmithKline Trading Services Limited
Тел.: + 359 80018205

Luxembourg/Luxemburg

GlaxoSmithKline Pharmaceuticals s.a./n.v.
Belgique/Belgien
Tél/Tel: + 32 (0)10 85 52 00

Česká republika

GlaxoSmithKline s.r.o.
Tel: + 420 222 001 111
cz.info@gsk.com

Magyarország

GlaxoSmithKline Trading Services Limited
Tel.: + 36 80088309

Danmark

GlaxoSmithKline Pharma A/S
Tlf.: + 45 36 35 91 00
dk-info@gsk.com

Malta

GlaxoSmithKline Trading Services Limited
Tel: + 356 80065004

Deutschland

GlaxoSmithKline GmbH & Co. KG
Tel.: + 49 (0)89 36044 8701
produkt.info@gsk.com

Nederland

GlaxoSmithKline BV
Tel: + 31 (0)33 2081100

Eesti

GlaxoSmithKline Trading Services Limited
Tel: + 372 8002640

Norge

GlaxoSmithKline AS
Tlf: + 47 22 70 20 00

Ελλάδα

GlaxoSmithKline Μονοπρόσωπη Α.Ε.Β.Ε.
Τηλ: + 30 210 68 82 100

España

GlaxoSmithKline, S.A.
Tel: + 34 900 202 700
Es-ci@gsk.com

France

Laboratoire GlaxoSmithKline
Tél.: + 33 (0)1 39 17 84 44
diam@gsk.com

Hrvatska

GlaxoSmithKline Trading Services Limited
Tel: + 385 800787089

Ireland

GlaxoSmithKline Trading Services Limited
Tel: + 353 (0)1 4955000

Ísland

Vistor hf.
Sími: + 354 535 7000

Italia

GlaxoSmithKline S.p.A.
Tel: + 39 (0)45 7741 111

Κύπρος

GlaxoSmithKline Trading Services Limited
Τηλ: + 357 80070017

Latvija

GlaxoSmithKline Trading Services Limited
Tel: + 371 80205045

Österreich

GlaxoSmithKline Pharma GmbH
Tel: + 43 (0)1 97075 0
at.info@gsk.com

Polska

GSK Services Sp. z o.o.
Tel.: + 48 (0)22 576 9000

Portugal

Glaxo Wellcome Farmacêutica, Lda.
Tel: + 351 21 412 95 00
FI.PT@gsk.com

România

GlaxoSmithKline Trading Services Limited
Tel: + 40 800672524

Slovenija

GlaxoSmithKline Trading Services Limited
Tel: + 386 80688869

Slovenská republika

GlaxoSmithKline Trading Services Limited
Tel: + 421 800500589

Suomi/Finland

GlaxoSmithKline Oy
Puh/Tel: + 358 (0)10 30 30 30

Sverige

GlaxoSmithKline AB
Tel: + 46 (0)8 638 93 00
info.produkt@gsk.com

This leaflet was last revised in

Detailed information on this medicine is available on the European Medicines Agency web site:
<http://www.ema.europa.eu>.

Package leaflet: Information for the patient

Zeffix 5 mg/ml oral solution lamivudine

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Zeffix is and what it is used for
2. What you need to know before you take Zeffix
3. How to take Zeffix
4. Possible side effects
5. How to store Zeffix
6. Contents of the pack and other information

1. What Zeffix is and what it is used for

The active ingredient in Zeffix is lamivudine.

Zeffix is used to treat long term (chronic) hepatitis B infection in adults.

Zeffix is an antiviral medicine that suppresses the hepatitis B virus and belongs to a group of medicines called *nucleoside analogue reverse transcriptase inhibitors (NRTIs)*.

Hepatitis B is a virus which infects the liver, causes long term (chronic) infection, and can lead to liver damage. Zeffix can be used in people whose liver is damaged, but still functions normally (compensated liver disease) and in combination with other medicines in people whose liver is damaged and does not function normally (decompensated liver disease).

Treatment with Zeffix can reduce the amount of hepatitis B virus in your body. This should lead to a reduction in liver damage and an improvement in your liver function. Not everyone responds to treatment with Zeffix in the same way. Your doctor will monitor the effectiveness of your treatment with regular blood tests.

2. What you need to know before you take Zeffix

Your healthcare provider should offer you counselling and testing for HIV infection before you start treatment with lamivudine for hepatitis B infection and during treatment. If you have or get HIV infection, see section 3.

Do not take Zeffix

- if you're **allergic** to lamivudine or to any of the other ingredients of this medicine (*listed in Section 6*).

→ **Check with your doctor** if you think this applies to you.

Warnings and precautions

Some people taking Zeffix or other similar medicines are more at risk of serious side effects. You need to be aware of the extra risks:

- if you have ever had other types of **liver disease**, such as hepatitis C
- if you're seriously **overweight** (especially if you're a woman).

→ **Talk to your doctor if any of these apply to you.** You may need extra check-ups, including blood tests, while you're taking your medication. See **Section 4** for more information about the risks.

Don't stop taking Zeffix without your doctor's advice, as there is a risk of your hepatitis getting worse. When you stop taking Zeffix your doctor will monitor you for at least four months to check for any problems. This will mean taking blood samples to check for any raised liver enzyme levels, which may indicate liver damage. See section 3 for more information about how to take Zeffix.

Protect other people

Hepatitis B infection is spread by sexual contact with someone who has the infection, or by transfer of infected blood (for example, by sharing injection needles). Zeffix will not stop you passing hepatitis B infection on to other people. To protect other people from becoming infected with hepatitis B:

- **Use a condom** when you have oral or penetrative sex.
- **Don't risk blood transfer** — for example, don't share needles.

Other medicines and Zeffix

Tell your doctor or pharmacist if you're taking, have recently taken or might take any other medicines, including herbal medicines or other medicines you bought without a prescription.

Remember to tell your doctor or pharmacist if you begin taking a new medicine while you're taking Zeffix.

These medicines should not be used with Zeffix:

- medicines (usually liquids) containing sorbitol and other sugar alcohols (such as xylitol, mannitol, lactitol or maltitol), if taken regularly
- other medicines containing lamivudine, used to treat **HIV infection** (sometimes called the AIDS virus)
- emtricitabine, used to treat **HIV** or **hepatitis B infection**
- cladribine, used to treat **hairy cell leukaemia**

→ **Tell your doctor** if you're being treated with any of these.

Pregnancy

If you are pregnant, think you may be pregnant or are planning to have a baby:

→ **Talk to your doctor** about the risks and benefits of taking Zeffix during your pregnancy.

Do not stop treatment with Zeffix without your doctor's advice.

Breast-feeding

Zeffix can pass into breast-milk. If you are breast-feeding, or thinking about breast-feeding:

→ **Talk to your doctor** before you take Zeffix.

Driving and using machines

Zeffix may make you feel tired, which could affect your ability to drive or use machines.

→ Don't drive or use machines unless you are sure you're not affected.

Zeffix contains sugar, preservatives, propylene glycol and sodium

If you are diabetic, please note that each dose of Zeffix (100 mg = 20 ml) contains 4 g of sucrose.

Zeffix contains sucrose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking Zeffix. Sucrose may be harmful to the teeth.

Zeffix contains preservatives (*parahydroxybenzoates*) which may cause allergic reactions (possibly delayed).

This medicine contains 400 mg propylene glycol in each 20 ml.

This medicine contains 58.8 mg sodium (main component of cooking/table salt) in each 20 ml. This is equivalent to 2.9% of the recommended maximum daily dietary intake of sodium for an adult.

3. How to take Zeffix

Always take this medicine exactly as your doctor has told you to. Check with your doctor or pharmacist if you're not sure.

Stay in regular contact with your doctor

Zeffix helps to control your hepatitis B infection. You need to keep taking it every day to control your infection and stop your illness getting worse.

→ **Keep in touch with your doctor, and don't stop taking Zeffix** without your doctor's advice.

How much to take

The usual dose of Zeffix is 20 ml (100 mg lamivudine) once a day.

Your doctor may prescribe a lower dose if you have problems with your kidneys.

→ **Talk to your doctor** if this applies to you.

Patients who also have or may get HIV infection

If you have or get HIV that is not being treated with medicines while taking lamivudine for the treatment of hepatitis B infection, the HIV virus may develop resistance to certain HIV medicines and become harder to treat. Lamivudine can also be used to treat HIV infection. Talk to your doctor if you have HIV infection. Your doctor may treat you with another medicine that contains a higher dose of lamivudine, usually 150 mg twice a day, as the lower dose of 100 mg lamivudine is not enough to treat HIV infection. If you are planning to change your HIV treatment, discuss this change with your doctor first.

→ **Talk to your doctor** if this applies to you.

Zeffix can be taken with or without food.

See the diagram and instructions after section 6 of this leaflet for how to measure and take a dose of medicine.

If you take more Zeffix than you should

If you accidentally take too much Zeffix, tell your doctor or your pharmacist, or contact your nearest hospital emergency department for further advice. If possible, show them the Zeffix pack.

If you forget to take Zeffix

If you forget to take a dose, take it as soon as you remember. Then continue your treatment as before. Don't take a double dose to make up for a missed dose.

Don't stop taking Zeffix

You must not stop taking Zeffix without consulting your doctor. There is a risk of your hepatitis getting worse (*see section 2*). When you stop taking Zeffix your doctor will monitor you for at least four months to check for any problems. This will mean taking blood samples to check for any raised liver enzyme levels, which may indicate liver damage.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Side effects that were commonly reported in Zeffix clinical trials were tiredness, respiratory tract infections, throat discomfort, headache, stomach discomfort and pain, nausea, vomiting and diarrhoea, increases in liver enzymes and enzymes produced in the muscles (*see below*).

Allergic reaction

These are rare (may affect up to 1 in 1,000 people). Signs include:

- swelling of eyelids, face or lips
- difficulty swallowing or breathing

→ **Contact a doctor immediately** if you get these symptoms. **Stop taking Zeffix.**

Side effects thought to be caused by Zeffix

A very common side effect (these may affect **more than 1 in 10** people) which may show up in blood tests is:

- an increase in the level of some liver enzymes (*transaminases*), which may be a sign of inflammation or damage in the liver.

A common side effect (these may affect **up to 1 in 10** people) is:

- cramps and muscle pains
- skin rash or ‘hives’ anywhere on the body

A common side effect which may show up in blood tests is:

- an increase in the level of an enzyme produced in the muscles (*creatine phosphokinase*) which may be a sign that body tissue is damaged.

A very rare side effect (these may affect up to 1 in 10,000 people) is:

- lactic acidosis (excess lactic acid in the blood).

Other side effects

Other side effects have occurred in a very small number of people but their exact frequency is unknown

- breakdown of muscle tissue
- a worsening of liver disease after Zeffix is stopped or during treatment if the hepatitis B virus becomes resistant to Zeffix. This can be fatal in some people.

A side effect which may show up in blood tests is:

- a decrease in the number of cells involved in blood clotting (*thrombocytopenia*).

If you get any side effects

→ **Talk to your doctor or pharmacist.** This includes any possible side effects not listed in this leaflet.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via **the national reporting system** listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Zeffix

Keep this medicine out of the sight and reach of children.

Don't take this medicine after the expiry date shown on the bottle and the carton.

Don't store above 25 °C.

Discard one month after first opening.

Do not throw away any medicines in your wastewater or household rubbish. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Zeffix contains

The active substance is lamivudine. Each ml of oral solution contains 5 mg of lamivudine.

The other ingredients are: sucrose, methyl parahydroxybenzoate (E218), propyl parahydroxybenzoate (E216), citric acid, propylene glycol (E1520), sodium citrate, artificial strawberry flavour, artificial banana flavour, purified water.

What Zeffix looks like and contents of the pack

Zeffix oral solution is supplied in cartons containing a white polyethylene bottle, with a child resistant cap. The solution is clear, colourless to pale yellow in colour with strawberry/banana flavouring. The bottle contains 240 ml of lamivudine solution (5 mg/ml). The pack includes an oral applicator with ml graduations and an applicator-adaptor, which should be fitted to the bottle before use.

Manufacturer

GlaxoSmithKline Trading Services Limited
12 Riverwalk
Citywest Business Campus
Dublin 24
Ireland
D24 YK11

Marketing Authorisation Holder

GlaxoSmithKline Trading Services
Limited
12 Riverwalk
Citywest Business Campus
Dublin 24
Ireland
D24 YK11

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder:

België/Belgique/Belgien

GlaxoSmithKline Pharmaceuticals s.a./n.v.
Tél/Tel: + 32 (0)10 85 52 00

Lietuva

GlaxoSmithKline Trading Services Limited
Tel: + 370 80000334

България

GlaxoSmithKline Trading Services Limited
Тел.: + 359 80018205

Luxembourg/Luxemburg

GlaxoSmithKline Pharmaceuticals s.a./n.v.
Belgique/Belgien
Tél/Tel: + 32 (0)10 85 52 00

Česká republika

GlaxoSmithKline s.r.o.
Tel: + 420 222 001 111

Magyarország

GlaxoSmithKline Trading Services Limited
Tel.: + 36 80088309

cz.info@gsk.com

Danmark

GlaxoSmithKline Pharma A/S
Tlf.: + 45 36 35 91 00
dk-info@gsk.com

Deutschland

GlaxoSmithKline GmbH & Co. KG
Tel.: + 49 (0)89 36044 8701
produkt.info@gsk.com

Eesti

GlaxoSmithKline Trading Services Limited
Tel: + 372 8002640

Ελλάδα

GlaxoSmithKline Μονοπρόσωπη Α.Ε.Β.Ε.
Τηλ: + 30 210 68 82 100

España

GlaxoSmithKline, S.A.
Tel: + 34 900 202 700
Es-ci@gsk.com

France

Laboratoire GlaxoSmithKline
Tél.: + 33 (0)1 39 17 84 44
diam@gsk.com

Hrvatska

GlaxoSmithKline Trading Services Limited
Tel: + 385 800787089

Ireland

GlaxoSmithKline Trading Services Limited
Tel: + 353 (0)1 4955000

Ísland

Vistor hf.
Sími: + 354 535 7000

Italia

GlaxoSmithKline S.p.A.
Tel: + 39 (0)45 7741 111

Κύπρος

GlaxoSmithKline Trading Services Limited
Τηλ: + 357 80070017

Latvija

GlaxoSmithKline Trading Services Limited
Tel: + 371 80205045

Malta

GlaxoSmithKline Trading Services Limited
Tel: + 356 80065004

Nederland

GlaxoSmithKline BV
Tel: + 31 (0)33 2081100

Norge

GlaxoSmithKline AS
Tlf: + 47 22 70 20 00

Österreich

GlaxoSmithKline Pharma GmbH
Tel: + 43 (0)1 97075 0
at.info@gsk.com

Polska

GSK Services Sp. z o.o.
Tel.: + 48 (0)22 576 9000

Portugal

Glaxo Wellcome Farmacêutica, Lda.
Tel: + 351 21 412 95 00
FI.PT@gsk.com

România

GlaxoSmithKline Trading Services Limited
Tel: + 40 800672524

Slovenija

GlaxoSmithKline Trading Services Limited
Tel: + 386 80688869

Slovenská republika

GlaxoSmithKline Trading Services Limited
Tel: + 421 800500589

Suomi/Finland

GlaxoSmithKline Oy
Puh/Tel: + 358 (0)10 30 30 30


Sverige

GlaxoSmithKline AB
Tel: + 46 (0)8 638 93 00
info.produkt@gsk.com

This leaflet was last revised in

Detailed information on this medicine is available on the European Medicines Agency web site:
<http://www.ema.europa.eu>.

How to measure the dose and take the medicine



Use the oral applicator supplied with the pack to measure your dose accurately (*see also Section 3*). When full, the applicator contains **10 ml of solution**.

- 1. Remove the plastic wrap** from the applicator/adapter.
- 2. Remove the adapter** from the applicator.
- 3. Remove the child-proof bottle cap (A)** and keep it safely.
- Hold the bottle. **Push the adapter (B) firmly into the neck of the bottle, as far as it will go.**
- Insert the applicator (C)** firmly into the adapter.
- Turn the bottle upside down.
- Pull out the applicator plunger (D)** until the applicator contains the first part of your full dose.
- Turn the bottle the right way up. **Remove the applicator from the adapter.**
- Put the applicator into your mouth**, placing the tip of the applicator against the inside of your cheek. **Slowly push the plunger in**, allowing time to swallow. **Don't push too hard** and squirt the liquid into the back of your throat, or you may choke.
- Repeat steps 5 to 9** in the same way until you have taken your whole dose. For example, if your dose is 20 ml, you need to take 2 applicator-fulls of medicine.
- Take the applicator out of the bottle** and **wash** it thoroughly in clean water. Let it dry completely before you use it again. Leave the adapter in the bottle.
- Close the bottle tightly** with the cap.

Discard oral solution one month after first opening.