

ANNEX I

SUMMARY OF PRODUCT CHARACTERISTICS

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

1. NAME OF THE MEDICINAL PRODUCT

ZEPZELCA 2 mg powder for concentrate for solution for infusion
ZEPZELCA 4 mg powder for concentrate for solution for infusion

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

ZEPZELCA 2 mg powder for concentrate for solution for infusion

Each vial of powder contains 2 mg of lurbinectedin.

ZEPZELCA 4 mg powder for concentrate for solution for infusion

Each vial of powder contains 4 mg of lurbinectedin.

One mL of reconstituted solution contains 0.5 mg of lurbinectedin.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Powder for concentrate for solution for infusion (powder for concentrate).

White to off-white powder

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

ZEPZELCA, in combination with atezolizumab, is indicated for the maintenance treatment of adult patients with extensive-stage small cell lung cancer (ES-SCLC) whose disease has not progressed after first-line induction therapy with atezolizumab, carboplatin and etoposide.

4.2 Posology and method of administration

ZEPZELCA therapy should be initiated and supervised by health professionals experienced in the use of anticancer products.

Posology

The recommended dose of lurbinectedin is 3.2 mg/m² every 21 days until disease progression or unacceptable toxicity when it is administered in combination with atezolizumab.

When administering lurbinectedin on the same day, atezolizumab should be administered first (see section 5.1).

For the recommended intravenous or subcutaneous dose of atezolizumab, as well as for recommendations regarding dose modification due to toxicity, refer to their prescribing information.

Treatment with ZEPZELCA should be initiated only if absolute neutrophil count (ANC) is at least $1.5 \times 10^9/L$ and platelet count is at least $100 \times 10^9/L$.

Treatment continuation and treatment delays

Further treatment cycles (i.e., cycle 2 or subsequent) will be administered every 21 days if the patient fulfils all the treatment continuation criteria listed above (see also Table 2 for dose modifications criteria for ZEPZELCA adverse reactions).

If a patient does not meet the requirements for treatment continuation on Day 1 of any cycle after Cycle 1, treatment will be withheld until appropriate recovery, for a maximum of 21 days after the treatment due date. If there is no recovery after a 21-days delay, treatment must be stopped.

In case atezolizumab is discontinued due to an immune-related severe adverse reaction, treatment with lurbinectedin may be continued at its current dose as a single agent. If immune toxicity re-occurs despite discontinuation of atezolizumab, treatment with lurbinectedin should also be discontinued.

Pre-infusion medicinal products

The following pre-infusion medicinal products should be administered for antiemetic prophylaxis:

- Corticosteroids (intravenous dexamethasone 8 mg or equivalent)
- Serotonin antagonists (intravenous ondansetron 8 mg or equivalent)

Post-infusion medicinal products

Primary prophylaxis with granulocyte colony-stimulating factor (G-CSF) is recommended to reduce the risk of severe neutropenia/febrile neutropenia.

If needed, post-medication can include administration of extended antiemetic treatment for 2 days:

- Corticosteroids (oral dexamethasone 4 mg or equivalent), or
- Serotonin antagonists (oral ondansetron 8 mg or equivalent) or
- Metoclopramide (intravenous or oral 10 mg or equivalent every 8 hours)

Dose adjustment for adverse reactions

The recommended dose reductions for adverse reactions are listed in Table 1.

Table 1: Dose reduction for ZEPZELCA for adverse reactions

Recommended starting dose	1 st Dose reduction	2 nd Dose reduction	3 rd Dose reduction
3.2 mg/m ²	2.6 mg/m ²	2.0 mg/m ²	Stop
1.6 mg/m ² *	1.3 mg/m ²	1.0 mg/m ²	Stop

*Dose reduction schedule applicable to the 50% reduced dose (i.e., 1.6 mg/m²) used in cases of moderated hepatic impairment or co-administration with strong or moderate CYP3A inhibitors.

The recommended dose modifications for adverse reactions are presented in Table 2.

Table 2: Dose modifications criteria for ZEPZELCA for adverse reactions

Adverse reaction	Severity ^a	Dose modification
Neutropenia ^b (see section 4.4)	Grade 4 OR any grade febrile neutropenia OR	• Withhold ZEPZELCA until Grade ≤ 1 and resolution of any associated fever/infection/sepsis, AND • Resume ZEPZELCA at a reduced dose^b

	associated with infection/sepsis at any grade	
Thrombocytopenia (see section 4.4)	Grade 3 with bleeding OR Grade 4	• Withhold ZEPZELCA until platelet $\geq 100 \times 10^9/L$, AND • Resume ZEPZELCA at reduced dose
Hepatotoxicity (see section 4.4) and other adverse reactions	Grade 2	• Withhold ZEPZELCA until Grade ≤ 1 (for AST and ALT until ≤ 3 ULN), AND • Resume ZEPZELCA at same dose
	Grade ≥ 3	• Withhold ZEPZELCA until Grade ≤ 1 (for AST and ALT until ≤ 3 ULN). AND • Resume ZEPZELCA at reduced dose
Rhabdomyolysis	Grade 2	• Withhold ZEPZELCA until Grade ≤ 1, AND • Resume ZEPZELCA at same dose
	Grade ≥ 3	• Permanently discontinue ZEPZELCA
Non-haematological toxicity	Grade 2	• Withhold ZEPZELCA until Grade ≤ 1, AND • Resume ZEPZELCA at same dose
	Grade ≥ 3	• Withhold ZEPZELCA until Grade ≤ 1, AND • Resume ZEPZELCA at reduced dose
Tumour Lysis Syndrome	Grade 2	• Withhold ZEPZELCA until Grade ≤ 1, AND • Resume ZEPZELCA at same dose
	Grade ≥ 3	• Permanently discontinue ZEPZELCA
Any adverse reaction that requires frequent or prolonged (>2 weeks) dose delays	-	• Reduce the dose of ZEPZELCA or discontinue

^a National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE) version 5.0.

^b Patients with isolated Grade 4 neutropenia (neutrophil count less than 500 cells/mm^3) and who had not received G-CSF as primary prophylaxis, may receive G-CSF prophylaxis rather than undergo lurbinectedin dose reduction.

Dose adjustment for co-administration with strong or moderate CYP3A inhibitors

Co-administration of lurbinectedin with strong or moderate CYP3A inhibitors should be avoided. If co-administration cannot be avoided, dose of lurbinectedin should be reduced by 50% of the approved dose (see section 4.5). In case of adverse reactions with the reduced initial dose, up to two subsequent dose reductions by 20% each are allowed (see Table 1 in section 4.2).

Special population

Elderly

No dose adjustment is needed in patients aged ≥ 65 years.

Renal impairment

No dose adjustment is recommended in patients with mild (CrCL 60-89 mL/min) or moderate (CrCL of 30-59 mL/min) renal impairment.

Lurbinectedin has not been evaluated in a sufficient number of patients with severe renal impairment (CrCL < 30 mL/min) or end-stage renal disease to estimate the risk; therefore, it should not be administered to these patients (see section 5.2).

Hepatic impairment

Treatment with lurbinectedin is not recommended in patients with elevated AST or ALT (AST or ALT > 3 × ULN), due to limited clinical experience.

No dose adjustment is recommended for patients with mild hepatic impairment (total bilirubin ≤ ULN and AST > ULN, or total bilirubin 1 to ≤ 1.5 × ULN and any AST).

In patients with moderate hepatic impairment (total bilirubin > 1.5 to ≤ 3 × ULN and any AST), the recommended dose of ZEPZELCA is 1.6 mg/m² by intravenous infusion over 60 minutes every 21 days until disease progression or unacceptable toxicity (see section 5.2). Patients with moderate hepatic impairment should be monitored for increased adverse reactions. In case of adverse reactions with the reduced initial dose, up to two subsequent dose reductions by 20% each are allowed (see Table 1 in section 4.2).

Administration of ZEPZELCA in patients with severe hepatic impairment (total bilirubin > 3 × ULN) should be avoided. If administration of ZEPZELCA cannot be avoided, the recommended dose is 1.6 mg/m² by intravenous infusion over 60 minutes every 21 days until disease progression or unacceptable toxicity (see section 5.2). Patients with severe hepatic impairment should be monitored for increased adverse reactions. In case of adverse reactions with the reduced initial dose, up to two subsequent dose reductions by 20% each are allowed (see Table 1 in section 4.2).

Paediatric population

There is no relevant use of ZEPZELCA in the paediatric population in the treatment of SCLC.

Method of administration

ZEPZELCA is for intravenous use only. It must be administered by intravenous infusion over a period of one hour.

Precautions to be taken before handling or administering the medicinal product

ZEPZELCA is to be reconstituted and then further diluted prior to administration.

The use of a central venous catheter should be considered to reduce the risk of extravasation (see section 4.4) and thrombophlebitis, particularly in patients with limited venous access.

For instructions on reconstitution and dilution of the medicinal product before administration, see section 6.6.

4.3 Contraindications

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- Breastfeeding (see section 4.6).

4.4 Special warnings and precautions for use

Myelosuppression

ZEPZELCA can cause severe and life-threatening myelosuppression including febrile neutropenia and sepsis.

ZEPZELCA should not be administered to patients with baseline neutrophil counts of less than $1.5 \times 10^9/L$ and platelet counts of less than $100 \times 10^9/L$.

Full blood counts including differential white blood cells and platelet count should be monitored at baseline and prior to each cycle. Dose modifications may be required (see Table 2 in section 4.2).

In case of neutrophil counts of less than $500/mm^3$ or any value less than lower limit of normal that is associated with infection/sepsis, the use of G-CSF is recommended.

Hepatotoxicity

ALT and AST increases have been reported with ZEPZELCA (see section 4.8).

Liver tests, including ALT, AST and bilirubin should be monitored prior to initiating ZEPZELCA and periodically during treatment as clinically indicated. Dose modifications may be required (see Table 1 in section 4.2).

Extravasation resulting in tissue necrosis

Extravasation of ZEPZELCA resulting in skin and soft tissue injury, including necrosis requiring debridement, may occur (see section 4.8).

The use of a central venous catheter should be considered to reduce the risk of extravasation, particularly in patients with limited venous access. Patients should be monitored for signs and symptoms of extravasation during the ZEPZELCA infusion.

If extravasation occurs, the infusion should be immediately discontinued, the infusion catheter should be removed, and the patient should be monitored for signs and symptoms of tissue necrosis. The time to onset of necrosis after extravasation may vary. Supportive care should be administered and an appropriate medical specialist should be consulted as needed for management of signs and symptoms of extravasation. Subsequent infusions should be administered at a site that was not affected by extravasation.

Rhabdomyolysis

Rhabdomyolysis has been reported in patients treated with ZEPZELCA (see section 4.8).

Creatine phosphokinase (CPK) should be monitored prior to initiating ZEPZELCA and periodically during treatment as clinically indicated.

If rhabdomyolysis occurs, supportive measures such as parenteral hydration, urine alkalinisation and dialysis should be promptly established, as indicated. Based on severity, ZEPZELCA treatment should be withheld or the dose should be reduced [see Table 2 in section 4.2].

Caution should be taken if medicinal products with known association with rhabdomyolysis (e.g., statins), are administered concomitantly with lurbinectedin, since the risk of rhabdomyolysis may be increased.

Tumour lysis syndrome (TLS)

Tumour lysis syndrome (TLS), which may be fatal, has been reported with ZEPZELCA therapy. Healthcare professionals are advised to closely monitor patients for TLS, especially those with a high tumour burden. Key precautions include preventing dehydration and managing electrolyte imbalances.

If TLS develops, it should be treated promptly, and the potential need for interruption or discontinuation of treatment should be considered (see section 4.2).

Co-administration with strong CYP3A inducers

Co-administration of strong CYP3A inducers should be avoided (see section 4.5).

Embryo-foetal toxicity

Lurbinectedin can cause fetal harm when administered to a pregnant woman. Pregnancy testing is recommended in women of childbearing potential prior to starting treatment.

Female patient of childbearing potential should use highly effective contraception during treatment and for 7 months after the last dose.

Male patients with female partners of childbearing potential should use condom during treatment and for 4 months after the last dose. Female partners of childbearing potential should use highly effective contraception for the same period (see section 4.6 and 5.3).

Disease-specific precautions - SCLC

Patients with ECOG performance status ≥ 2 ; central nervous system (CNS) metastases, a history of autoimmune disease, or administration of systemic immunosuppressive medicinal products within 1 week prior to enrolment were excluded in the pivotal study in SCLC (see section 5.1). In the absence of data, lurbinectedin in combination with atezolizumab should be used with caution in these populations after careful consideration of the potential benefit/risk on an individual basis.

Excipients

This medicinal product contains less than 1 mmol of sodium (23 mg) per vial, that is to say essentially “sodium-free”.

4.5 Interaction with other medicinal products and other forms of interaction

Effect of strong or moderate CYP3A inhibitors on lurbinectedin

In a dedicated drug-drug interaction study (n=8) with itraconazole, a strong CYP3A4 inhibitor, systemic exposure of total lurbinectedin was increased by approximately 2.7-fold ($AUC_{0-\infty}$) and total plasma clearance was reduced by 63%, when lurbinectedin was given concomitantly with itraconazole (total daily dose of 200 mg during 12 days, 4 days before up to 8 days after the lurbinectedin administration).

Co-administration of ZEPZELCA with strong or moderate CYP3A inhibitors should be avoided. If co-administration with strong CYP3A inhibitors (e.g., ketoconazole, itraconazole, posaconazole, voriconazole, clarithromycin, telithromycin, lopinavir, ritonavir, saquinavir, nelfinavir, atazanavir, indinavir, boceprevir, telaprevir) or moderate CYP3A inhibitors (e.g., aprepitant, ciprofloxacin, erythromycin, cyclosporine, fluconazole, diltiazem, verapamil) cannot be avoided, the dose of ZEPZELCA should be reduced by 50% of the approved dose (see section 4.2). In case of adverse reactions with the reduced initial dose, up to two subsequent dose reductions by 20% each are allowed (see Table 1 in section 4.2).

Effect of strong CYP3A inducers on lurbinectedin

In a dedicated drug-drug interaction study (n=8) with bosentan, a moderate CYP3A4 inducer, systemic exposure of total lurbinectedin was decreased by approximately 20% ($AUC_{0-\infty}$) and total plasma clearance was increased by 25% when lurbinectedin was given concomitantly with bosentan (125 mg twice daily during 5 days). Therefore, the magnitude of these changes precludes a clinically relevant

effect of co-administration of moderate CYP3A4 inducers (e.g., bosentan, cenobamate, dabrafenib, efavirenz, etravirine, lorlatinib, pexidartinib, phenobarbital, primidone, sotorasib) on lurbinectedin exposure and no dose adjustment is required.

Co-administration of strong CYP3A inducers (e.g., carbamazepine, phenobarbital, phenytoin, rifampicin, rifabutin, rifapentine, St. John's Wort (*Hypericum perforatum*)) with ZEPZELCA should be avoided. Consider alternative agents with less CYP3A induction (see section 4.4).

4.6 Fertility, pregnancy and lactation

Women of childbearing potential / Contraception in males and females

Pregnancy testing is recommended in women of childbearing potential prior to starting treatment with lurbinectedin.

Female patients of childbearing potential should use highly effective contraception during treatment and for 7 months after the last dose.

Male patients with female partners of childbearing potential should use condom during treatment and for 4 months after the last dose. Female partners of childbearing potential should use highly effective contraception for the same period (see sections 4.4 and 5.3).

Pregnancy

There are no or limited amount of data from the use of lurbinectedin in pregnant women.

Studies in animals have shown severe embryo-foetal development toxicity (see section 5.3).

Lurbinectedin should not be used during pregnancy unless the clinical condition of the woman requires treatment with lurbinectedin.

Pregnant or non-pregnant women of childbearing potential should be advised of the potential risk to a foetus. If ZEPZELCA is used during pregnancy, or if a patient becomes pregnant while receiving ZEPZELCA, the patient should be apprised of the potential risk to the foetus.

Breastfeeding

It is unknown whether lurbinectedin/metabolites are excreted in human milk.

A risk to the suckling child cannot be excluded.

Lurbinectedin is contraindicated during breastfeeding.

Fertility

Although no specific studies were conducted to assess fertility with lurbinectedin, and no clear signals of toxicity of reproductive organs were observed in toxicity studies, due to the nature of the compound (cytotoxic and mutagenic) it is likely to affect the reproductive capacity.

Advice on conservation of ovules or sperm should be sought prior to treatment because of the possibility of irreversible infertility due to therapy with lurbinectedin. Genetic counselling is also recommended for patients wishing to have children after therapy.

4.7 Effects on ability to drive and use machines

ZEPZELCA has moderate influence on the ability to drive and use machines. Patients experiencing fatigue, dizziness, vertigo and nausea should be advised not to drive and use machines until symptoms abate (see section 4.8)

4.8 Undesirable effects

Summary of the safety profile

The most common adverse reactions were nausea (37.6%), fatigue* (34.3%), anaemia (33.9%), thrombocytopenia (27.7%), and neutropenia (25.2%).

The most frequent grade 3/4 adverse reactions were neutropenia (12.4%), thrombocytopenia (11.2%), anaemia (9.5%) and fatigue* (5.0%).

Serious adverse reactions occurred in 34.3% of patients receiving ZEPZELCA with atezolizumab. The most frequent serious adverse reactions were thrombocytopenia (2.9%), pneumonia (3.7%), respiratory tract infection (2.5%) and dyspnoea (2.1%). Fatal adverse reactions occurred in 5% of patients receiving ZEPZELCA with atezolizumab, in most cases due to pneumonia and other lung infections.

Treatment with ZEPZELCA was permanently discontinued due to adverse reactions in 5.8% of patients who were receiving ZEPZELCA in combination with atezolizumab. The most frequent adverse reaction requiring permanent discontinuation of ZEPZELCA was neutropenia (1.7%).

Adverse reactions leading to interruption of ZEPZELCA in patients who received ZEPZELCA with atezolizumab occurred in 28.9% of patients; the most common adverse reactions leading to interruption were neutropenia (5.4%), anaemia (5.0%), fatigue* (4.6%) and thrombocytopenia (3.3%).

Dose reductions of ZEPZELCA due to an adverse reaction in patients who received ZEPZELCA with atezolizumab occurred in 16.1% of patients. The most frequent adverse reactions requiring dose reductions in patients who received ZEPZELCA with atezolizumab included thrombocytopenia (4.1%), fatigue* (3.3%), nausea (2.1%) and vomiting (2.1%).

* For Preferred Terms merged see footnote in Table 3.

Tabulated list of adverse reactions

Adverse reactions reported in IMforte clinical study are listed by MedDRA System Organ Class and by frequency in Table 3.

The frequencies of adverse reactions are based on all-cause adverse event frequencies identified in 242 patients exposed to lurbinectedin in combination with atezolizumab during a median treatment duration of 4.4 months in the clinical study IMforte (see section 5.1 for information on the main characteristics of participants in this clinical study). Additional adverse reactions were reported post-marketing.

Frequencies are defined as: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1\ 000$ to $< 1/100$); rare ($\geq 1/10\ 000$ to $< 1/1\ 000$); very rare ($< 1/10\ 000$); “not known (cannot be estimated from available data)”. Within each frequency grouping, adverse reactions are presented in the order of decreasing seriousness.

Table 3. Adverse reactions experienced by patients treated with ZEPZELCA in combination with atezolizumab

Frequency category (any grade)	Adverse reaction by system organ class	Any grade (%)	Grade ≥ 3 (%)
Infections and infestations			
Common	Pneumonia	5.4	3.3

Frequency category (any grade)	Adverse reaction by system organ class	Any grade (%)	Grade ≥ 3 (%)
Uncommon	Urinary tract infection ^a	5.4	0.4
	Infection	3.3	1.2
	Skin infection ^b	2.1	0.4
	Sepsis	0.4	0.4
Blood and lymphatic system disorders			
Very common	Anaemia	33.9	9.5
	Thrombocytopenia	27.7	11.2
	Neutropenia	25.2	12.4
	Leukopenia	12.4	2.9
Common	Lymphopenia	5.4	2.1
	Febrile neutropenia	1.7	1.7
Uncommon	Pancytopenia	0.4	0.4
Endocrine disorders			
Common	Hypothyroidism	7.9	0
Metabolism and nutrition disorders			
Very common	Decreased appetite	18.2	0.8
Common	Hypomagnesaemia	5.4	0.4
	Hypocalcaemia	4.5	0.8
Very rare	Tumour Lysis Syndrome ^c	frequency not known	-
Nervous system disorders			
Common	Neuropathy peripheral ^d	8.3	0.8
	Headache	6.6	0
	Dysgeusia	2.9	0
Vascular disorders			
Common	Phlebitis	7.0	0
	Thrombophlebitis	4.5	0.4
Respiratory, thoracic and mediastinal disorders			
Very common	Dyspnoea	10.7	2.5
Common	Cough	9.9	0
	Pneumonitis	4.5	0.8
	Productive cough	4.1	0
Gastrointestinal disorders			
Very common	Nausea	37.6	2.9
	Diarrhoea	15.7	0.4
	Vomiting	14.9	0.8
	Constipation	12.8	0
Common	Abdominal pain ^e	9.9	0.4
	Dyspesia	4.5	0
	Stomatitis	2.5	0
Skin and subcutaneous tissue disorders			
Common	Pruritus	7.9	0.4
	Rash	5.8	0
Musculoskeletal and connective tissue disorders			
Very common	Musculoskeletal pain ^f	15.7	0.8
Common	Arthralgia	8.3	1.2
Rare	Rhabdomyolysis ^c	frequency not known	-
General disorders and administration site conditions			
Very common	Fatigue ^g	34.3	5.0
Common	Oedema ^h	6.2	0.4

Frequency category (any grade)	Adverse reaction by system organ class	Any grade (%)	Grade ≥ 3 (%)
	Pyrexia	5.4	0
	Peripheral swelling	4.5	0.4
	Extravasation ⁱ	3.3	0
	Mucosal inflammation	2.5	0
Investigations			
Common	Transaminases increased ^j	9.1	2.9
	Blood creatinine increased	5.4	0
	Gamma-glutamyltransferase increased	3.3	0.8
	Blood creatine phosphokinase increased	2.1	0.4
	Weight decreased	3.3	0
^a including, Urinary tract infection, Cystitis ^b including Skin infection, Cellulitis ^c frequency not known (cannot be estimated from available data), reported in port-marketing setting (information related to grade not available). ^d including Hypoesthesia, Neuropathy peripheral, Paraesthesia, Peripheral sensory neuropathy. ^e including Abdominal discomfort, Abdominal distension, Abdominal pain, Abdominal pain upper. ^f including Back pain, Musculoskeletal chest pain, Musculoskeletal pain, Myalgia, Neck pain, Pain in extremity ^g including Asthenia, Fatigue. ^h including Oedema, Oedema peripheral ⁱ in few cases tissue necrosis was reported ^j including Alanine aminotransferase increased, Aspartate aminotransferase increased, Transaminases increased			

Description of selected adverse reaction

Neutropenia

In IMforte, 25.2% of patients experienced neutropenia (all grades), 12.4% experienced Grade 3/4 neutropenia, and 1.7% experienced febrile neutropenia and 0.4% sepsis. The median time to first onset of neutropenia* (all grade) was 10 (range: 7-29) days. The median duration was 11 (range: 1-196) days. Neutropenia* led to dose reduction or interruption in 1.7% or 5.4% of patients, respectively. Treatment was permanently discontinued in 1.7% of patients.

Hepatotoxicity

In IMforte, ALT increase was reported in 6.6% of patients (2.5% $\geq$ Grade 3), while AST increase was reported in 7.0% of patients (1.2% $\geq$ Grade 3). The median time to first onset of ALT increase (all grade) was 7 (range: 3-22) days. The median duration was 17 (range: 7-21) days. ALT increase led to dose reduction or interruption in 0.4% of patients each, respectively. The median time to first onset of AST increased (all grade) was 4 (range: 3-8) days. The median duration was 9 (range: 6-21) days. AST increased led to dose reduction in 0.8% of patients.

Rhabdomyolysis

Cases of rhabdomyolysis have been reported with post-marketing use of ZEPZELCA. No fatal cases have been reported.

Extravasation

Cases of extravasation with local irritation have been uncommonly reported with post-marketing use of ZEPZELCA. In a few cases, tissue necrosis requiring debridement was reported.

Tumour lysis syndrome

Cases of tumour lysis syndrome have been reported with post-marketing use of ZEPZELCA.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

4.9 Overdose

If an overdose is suspected, monitor the patient closely for myelosuppression and hepatic enzymes and institute supportive-care measures as appropriate.

There is no known antidote for overdose with lurbinectedin.

Haemodialysis is not expected to enhance the elimination of lurbinectedin because lurbinectedin is highly bound to plasma proteins (99%), and renal excretion is negligible.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antineoplastic agents, other antineoplastic agents, ATC code: L01XX69

Mechanism of action

Lurbinectedin inhibits the oncogenic transcription process through (i) its binding to CG-rich sequences of DNA, located within promoters of protein-coding genes; (ii) the eviction of oncogenic transcription factors from their binding sites; and (iii) the stalling of elongating RNA polymerase II and its specific degradation by the ubiquitin/proteasome machinery with all these processes leading to subsequent cell cycle arrest and tumour cell apoptosis.

Lurbinectedin suppresses the expression of inflammatory and motility-related genes at non-toxic nanomolar concentrations *in vitro*, while also inhibiting cell migration and adhesion. At higher concentrations, it induces apoptosis in monocytes and macrophages through caspase-8 activation. *In vivo* (murine models), antitumour dosing (0.18–0.20 mg/kg) restricts tumour growth, reduces specific immune cell populations, and decreases tumour vascularity.

Pharmacodynamic effects

Cardiac Electrophysiology

The potential for QTc prolongation with lurbinectedin was evaluated in 39 patients with advanced cancer. Large effects (>10 ms) on the QTc interval were not detected with lurbinectedin dosed at 3.2 mg/m² every 21 days.

Clinical efficacy and safety

Extensive-stage small cell lung cancer

The efficacy of maintenance treatment with ZEPZELCA in combination with atezolizumab was investigated in 483 patients with first-line ES-SCLC in IMforte, a randomised, multicentre, open-label

study. Participants were eligible for randomisation if they have achieved CR, PR, or SD by RECIST v1.1 based on radiographic assessment within 28 days prior to randomisation after completion of 4 cycles of induction treatment with atezolizumab, carboplatin and etoposide and if they had an ECOG performance status of 0 or 1. Eligible patients were randomised 1:1 to receive maintenance treatment with either lurbinectedin with atezolizumab or atezolizumab alone. Unless contraindicated, primary prophylaxis with G-CSF was given for patients assigned to the lurbinectedin with atezolizumab arm. The study excluded patients with CNS metastases, a history of autoimmune disease, or administration of systemic immunosuppressive medicines within 1 week prior to enrolment. Randomisation was stratified by ECOG performance status (0 vs. 1), lactate dehydrogenase (LDH) ($\leq$ ULN vs. $>$ ULN), presence of liver metastases at enrolment (yes vs. no), and prior receipt of prophylactic cranial irradiation (yes vs. no).

Patients were randomised to one of the following two treatment arms:

- ZEPZELCA 3.2 mg/m² intravenous with atezolizumab 1200 mg intravenous once every 3 weeks until disease progression or unacceptable toxicity, or
- Atezolizumab 1200 mg intravenous once every 3 weeks until disease progression or unacceptable toxicity.

Primary efficacy endpoints were overall survival (OS) and Independent Review Facility (IRF)-assessed progression-free survival (PFS) per the Response Evaluation Criteria in Solid Tumours (RECIST) v1.1 in the randomised population. (see Table 5).

A total of 483 patients were randomised: 242 to the ZEPZELCA with atezolizumab arm and 241 to the atezolizumab arm. The median age was 66 years (range 35 to 85 years, being 13% $\geq$ 75 years). The majority of patients were White (81.6%); 12.8% were Asian, 6.6% were Hispanic and $<$ 1% were Black or African American. Most patients were male (62.5%) and 97.5% were current or previous smokers. Baseline ECOG performance status was 0 (42.9%) or 1 (57.1%).

Efficacy results are presented in Table 5 and Figures 1 and 2.

Table 4: Efficacy results from IMforte

	lurbinectedin with atezolizumab N=242	atezolizumab N=241
Overall Survival¹		
Deaths (%)	113 (46.7%)	136 (56.4%)
Median, months (95% CI)	13.2 (11.9, 16.4)	10.6 (9.5, 12.2)
Hazard ratio ² (95% CI) p-value ^{3, 6}		0.73 (0.57, 0.95) 0.0174
Progression-Free Survival^{1, 4, 5}		
Number of events (%)	174 (71.9%)	202 (83.8%)
Median, months (95% CI)	5.4 (4.2, 5.8)	2.1 (1.6, 2.7)
Hazard ratio ² (95% CI) p-value ^{3, 7}		0.54 (0.43, 0.67) $<$ 0.0001
Cut-off: 29 July 2024		
¹ Measured from the time of randomisation		
² Stratified by ECOG performance status, LDH level, presence of liver metastases and prior receipt of prophylactic cranial irradiation		
³ Based on the stratified log-rank test		
⁴ As determined by IRF		
⁵ per RECIST v1.1		
⁶ Compared to the allocated alpha of 0.0313 (two-sided) for this interim OS analysis.		
⁷ Compared to the allocated alpha of 0.001 (two-sided) for this final PFS analysis.		
CI=confidence interval		

Figure 1: Kaplan-Meier plot of overall survival in IMforte





Figure 2: Kaplan-Meier plot of IRF-assessed progression free survival in IMforte



Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies with ZEPZELCA in all subsets of the paediatric population in the treatment of SCLC (see section 4.2 for information on paediatric use).

5.2 Pharmacokinetic properties

After a 3.2 mg/m² lurbinectedin dose administered as a 1-hour intravenous infusion, geometric means of total plasma C_{max} and AUC_∞, were 107 µg/L and 551 µg*h/L, respectively. No accumulation of lurbinectedin in plasma is observed upon repeated administrations every 21 days.

Distribution

Typical volume of distribution of lurbinectedin at steady state is 504 L. Binding to plasma proteins is approximately 99%, to both albumin and α-1-acid glycoprotein, with a calculated blood-to-plasma partitioning ratio of 0.68.

Biotransformation

In vitro studies

In vitro studies with human liver microsomes and supersomes indicate that CYP3A4 is the main CYP enzyme responsible for the hepatic metabolism of lurbinectedin.

Cytochrome P450 (CYP) Enzymes: Lurbinectedin is not an inhibitor of CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP2E1, or CYP3A4. Lurbinectedin is not an inducer of CYP1A2 or CYP3A4. The potential of lurbinectedin to induce CYP2B6 is unknown.

Transporter Systems: Lurbinectedin is a substrate of MDR1 (P-gp), but is not a substrate of OATB1P1, OATP1B3, OCT1, or MATE1. *In vitro*, lurbinectedin showed inhibitory potential towards MDR1, OATP1B1, OATP1B3, and OCT1 however, these findings are not considered clinically relevant. Lurbinectedin is not an inhibitor of BCRP, BSEP, MATE1, OAT1, OAT3, or OCT2.

Elimination

The terminal half-life of lurbinectedin is 51 hours. Total plasma clearance of lurbinectedin is 11 L/h.

The major route of lurbinectedin-related radioactivity excretion was via faeces (89% of dose), with only traces amounts of unchanged lurbinectedin detected in faeces (<0.2% of dose). Excretion in urine was the minor route (6% of dose), mainly as unchanged compound (1% of dose) and one metabolite (up to 1% of dose).

Linearity/non-linearity

Lurbinectedin pharmacokinetics is linear at the dose range of 0.02-6.9 mg/m².

Special populations

Population pharmacokinetics analyses showed that weight (range: 39-154 kg), age (range: 18-85 years), and gender do not have a clinically meaningful influence on the systemic exposure of lurbinectedin.

Hepatic impairment

A dedicated study was conducted to evaluate the influence of varying degrees of hepatic impairment (HI) on lurbinectedin in patients with advanced solid tumours. Patients were classified according to the National Cancer Institute Organ Dysfunction Working Group (NCI-ODWG) classification as having normal hepatic function or mild (total bilirubin ≤ ULN and AST > ULN, or total bilirubin > 1

to $\leq 1.5 \times \text{ULN}$ and $\text{AST} = \text{any}$), moderate (total bilirubin > 1.5 to $\leq 3 \times \text{ULN}$ and $\text{AST} = \text{any}$), or severe (total bilirubin $> 3 \times \text{ULN}$) HI. Patients with normal hepatic function and mild HI received lurbinectedin at 3.2 mg/m^2 and patients with moderate and severe HI received lurbinectedin at 1.6 mg/m^2 . No statistically significant differences were observed on total lurbinectedin pharmacokinetics among cohorts. A statistically significant higher dose-normalised M1 AUC metabolite/parent ratio (MPR) was observed in patients with severe HI (ratio: 5.95, 90% CI: 2.54–13.98) and moderate HI (ratio: 8.65, 90% CI: 3.94–19.01) compared to patients with mild HI. No statistically significant differences were observed in M4 MPR according to HI groups.

Based on population pharmacokinetic analysis, no apparent pharmacokinetic difference was observed in 125 patients with mild who received lurbinectedin 3.2 mg/m^2 every 21 days as compared to 625 patients with normal hepatic function.

Renal impairment

No dedicated studies of lurbinectedin have been conducted in patients with renal impairment. Based on population pharmacokinetic analyses, no apparent pharmacokinetic difference was observed in 165 patients with mild renal impairment (CrCL of 60-89 mL/min), 73 patients with moderate renal impairment (CrCL of 30-59 mL/min), and one patient with severe renal impairment (CrCL of 26 mL/min) who received lurbinectedin 3.2 mg/m^2 every 21 days as compared to 166 patients with normal renal function. The pharmacokinetic characteristics of lurbinectedin in patients with CrCL $< 30 \text{ mL/min}$ or patients on dialysis are unknown.

5.3 Preclinical safety data

Toxicology

The primary target for toxicity identified in nonclinical species (rat, dog and NHP) was characterised by severe, reversible and non-cumulative atrophy of bone marrow, which was associated with dose-related leukopenia, as well as thrombocytopenia and anaemia. In addition, lurbinectedin-treated animals experienced liver abnormalities (multiple dark areas or swollen liver, increased liver function markers, bile duct damage with necrosis and/or oedema, and hepatocellular degeneration/apoptosis and periportal hepatocytic hypertrophy). Additional findings were located in the gastro-intestinal tract (mucosal atrophy), kidneys (cortical tubular degeneration and vacuolation), heart (focal, slight to moderate myocardial degeneration and/or necrosis) and injection site (perivascular/vascular inflammatory reactions). A full recovery, after cessation of dosing, was noted for the majority of these alterations.

Genotoxicity

Positive genotoxicity results were obtained *in vitro* in mammalian cell lines showing dose related toxicity at all concentrations tested (range from 48 to 0.188 ng/mL). Positive genotoxicity findings are expected for lurbinectedin as a DNA-interacting antineoplastic agent (see section 4.6).

Carcinogenic potential

Carcinogenicity testing of lurbinectedin has not been performed.

Reproduction and development

Lurbinectedin induced maternal toxicity at the single dose MTD level of 0.6 mg/m^2 administered on Day 10 post-coitum and severe embryo-toxicity, leading to 100% embryo lethality (see section 4.4 and 4.6).

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sucrose
Lactic acid
Sodium hydroxide (for pH-adjustment)

6.2 Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

6.3 Shelf life

Unopened vial

ZEPZELCA 2 mg powder for concentrate for solution for infusion

18 months

ZEPZELCA 4 mg powder for concentrate for solution for infusion

5 years

Reconstituted and diluted solution

Chemical and physical in-use stability has been demonstrated for 24 hours at either 2 to 8°C or 25 °C.

From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 to 8°C, unless reconstitution / dilution has taken place in controlled and validated aseptic conditions. If reconstitution / dilution has taken place in controlled and validated aseptic conditions the prepared ready to administer product can be stored up to 24 hours at either +2°C to +8°C or +25 °C.

6.4 Special precautions for storage

Store in a refrigerator (2°C – 8°C).

For storage conditions after reconstitution and dilution of the medicinal product, see section 6.3.

6.5 Nature and contents of container

ZEPZELCA 2 mg powder for concentrate for solution for infusion

20 mL vial (clear type 1 glass) with a stopper (butyl rubber) and white coloured overseal (aluminium), containing 2 mg lurbinectedin.

Pack size of 1 vial.

ZEPZELCA 4 mg powder for concentrate for solution for infusion

30 mL vial (clear type 1 glass) with a stopper (butyl rubber) and blue coloured overseal (aluminium), containing 4 mg lurbinectedin.

Pack size of 1 vial.

6.6 Special precautions for disposal and other handling

Appropriate procedures for proper handling and disposal of cytotoxic medicinal products must be followed. You should have received training on the correct techniques to reconstitute and dilute ZEPZELCA and you should wear protective clothing including mask, goggles and gloves during the reconstitution and dilution. Accidental contact with the skin, eyes or mucous membranes must be treated immediately with copious amounts of water. You should not work with this medicine if you are pregnant.

Prepare the solution for infusion using aseptic technique as follows:

- Inject 8 mL (for 4 mg strength) or 4 mL (for 2 mg strength) of water for injections into the vial, yielding a solution containing 0.5 mg/mL lurbinectedin. Shake the vial until complete dissolution. The reconstituted solution is a clear, colourless or slightly yellowish solution, essentially free of visible particles. Visually inspect the solution for particulate matter and discoloration.
- Calculate the required volume of reconstituted solution as follows:
$$\text{Volume (mL)} = \frac{\text{Body Surface Area (m}^2\text{)} \times \text{Individual Dose (mg/m}^2\text{)}}{0.5 \text{ mg/mL}}$$
- For administration through a central venous line, withdraw the appropriate amount of reconstituted solution from the vial and add to an infusion container containing at least 100 mL of diluent sodium chloride 9 mg/mL (0.9%) solution for infusion or glucose 50 mg/mL (5%) solution for infusion.
- For administration through a peripheral venous line, withdraw the appropriate amount of reconstituted solution from the vial and add to an infusion container containing at least 250 mL of diluent sodium chloride 9 mg/mL (0.9%) solution for infusion or glucose 50 mg/mL (5%) solution for infusion.

The following materials are compatible with ZEPZELCA diluted solution:

- Polyolefin containers (polyethylene, polypropylene and mixtures).
- PVC (non-DEHP-containing), polyurethane and polyolefin infusion sets (polyethylene, polypropylene and polybutadiene).
- Polyether sulfone in-line filters with pore sizes of 0.22 micron.
- Implantable venous access systems with titanium and plastic resin ports and with polyurethane or silicone intravenous catheters.

ZEPZELCA can be administered with or without an in-line filter.

Infusion lines containing nylon membrane filters should not be used when the reconstituted ZEPZELCA solution is diluted with sodium chloride 9 mg/mL (0.9%) solution for infusion.

Lurbinectedin is a cytotoxic medicinal product. Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Pharma Mar, S.A.

Avda. de los Reyes 1
Polígono Industrial La Mina
28770 Colmenar Viejo (Madrid)
Spain
Tel: +34 91 846 60 00
Fax: +34 91 846 60 01

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/26/2032/001
EU/1/26/2032/002

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <https://www.ema.europa.eu>

ANNEX II

- A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturer responsible for batch release

Pharma Mar, S.A.
Avda. de los Reyes 1
Polígono Industrial La Mina
28770 Colmenar Viejo (Madrid)
Spain

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

The marketing authorisation holder (MAH) shall submit the first PSUR for this product within 6 months following authorisation.

D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

- **Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**CARTON BOX ZEPZELCA 2 mg****1. NAME OF THE MEDICINAL PRODUCT**

ZEPZELCA 2 mg powder for concentrate for solution for infusion
lurbinectedin

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each vial of powder contains 2 mg of lurbinectedin.
One mL of reconstituted solution contains 0.5 mg of lurbinectedin.

3. LIST OF EXCIPIENTS

Also contains: sucrose, lactic acid and sodium hydroxide

4. PHARMACEUTICAL FORM AND CONTENTS

Powder for concentrate for solution for infusion
1 vial

5. METHOD AND ROUTE(S) OF ADMINISTRATION

For intravenous use after reconstitution and further dilution.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

Cytotoxic: handle with caution.

8. EXPIRY DATE

EXP:

9. SPECIAL STORAGE CONDITIONS

Store in a refrigerator.
For storage conditions for the reconstituted and diluted solutions, see the package leaflet.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE
--

Discard any unused product or waste material in accordance with local requirements.

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Pharma Mar, S.A.
Avda. de los Reyes 1
Pol. Ind. La Mina
28770 Colmenar Viejo (Madrid)
Spain

12. MARKETING AUTHORISATION NUMBER(S)
--

EU/1/26/2032/001

13. BATCH NUMBER

Lot:

14. GENERAL CLASSIFICATION FOR SUPPLY
--

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA
--

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**VIAL ZEPZELCA 2 mg****1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

ZEPZELCA 2 mg powder for concentrate
lurbinectedin
IV use after reconstitution and dilution

2. METHOD OF ADMINISTRATION**3. EXPIRY DATE**

EXP:

4. BATCH NUMBER

Lot:

5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**6. OTHER**

Cytotoxic
PharmaMar, S.A. (logo)

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**CARTON BOX ZEPZELCA 4 mg****1. NAME OF THE MEDICINAL PRODUCT**

ZEPZELCA 4 mg powder for concentrate for solution for infusion
lurbinectedin

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each vial of powder contains 4 mg of lurbinectedin.
One mL of reconstituted solution contains 0.5 mg of lurbinectedin.

3. LIST OF EXCIPIENTS

Also contains: sucrose, lactic acid and sodium hydroxide

4. PHARMACEUTICAL FORM AND CONTENTS

Powder for concentrate for solution for infusion
1 vial

5. METHOD AND ROUTE(S) OF ADMINISTRATION

For intravenous use after reconstitution and further dilution.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

Cytotoxic: handle with caution.

8. EXPIRY DATE

EXP:

9. SPECIAL STORAGE CONDITIONS

Store in a refrigerator.
For storage conditions for the reconstituted and diluted solutions, see the package leaflet.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE
--

Discard any unused product or waste material in accordance with local requirements.

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Pharma Mar, S.A.
Avda. de los Reyes 1
Pol. Ind. La Mina
28770 Colmenar Viejo (Madrid)
Spain

12. MARKETING AUTHORISATION NUMBER(S)
--

EU/1/26/2032/002

13. BATCH NUMBER

Lot:

14. GENERAL CLASSIFICATION FOR SUPPLY
--

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA
--

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**VIAL ZEPZELCA 4 mg****1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

ZEPZELCA 4 mg powder for concentrate
lurbinectedin
IV use after reconstitution and dilution

2. METHOD OF ADMINISTRATION**3. EXPIRY DATE**

EXP:

4. BATCH NUMBER

Lot:

5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**6. OTHER**

Cytotoxic
PharmaMar, S.A. (logo)

B. PACKAGE LEAFLET

Package leaflet: Information for the user

ZEPZELCA 2 mg powder for concentrate for solution for infusion ZEPZELCA 4 mg powder for concentrate for solution for infusion

lurbinectedin

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor.
- If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What ZEPZELCA is and what it is used for
2. What you need to know before you use ZEPZELCA
3. How to use ZEPZELCA
4. Possible side effects
5. How to store ZEPZELCA
6. Contents of the pack and other information

1. What ZEPZELCA is and what it is used for

ZEPZELCA is a cancer medicine that contains the active substance lurbinectedin.

ZEPZELCA is used in adults for the treatment of a type of small cell lung cancer (SCLC) that has spread within the lung or to other parts of the body (extensive-stage SCLC). It is used together with atezolizumab as maintenance treatment in adults whose cancer has not progressed following treatment with atezolizumab, carboplatin and etoposide (other cancer medicines).

The active substance in ZEPZELCA, lurbinectedin, works by attaching to the genetic material (DNA) inside cancer cells. This damages the DNA and interferes with the cell growth and multiplication, leading to cancer cell death. It also reduces the activity of certain immune cells that help tumours grow.

Zepzelca may be given in combination with other anticancer medicine. It is important that you also read the package leaflets for the other anticancer medicine you may be receiving. If you have any questions about these medicines, ask your doctor.

2. What you need to know before you use ZEPZELCA

Do not use ZEPZELCA

- If you are allergic to lurbinectedin or any of the other ingredients of this medicine (listed in section 6).
- If you are breastfeeding.

Warnings and precautions

Talk to your doctor before using ZEPZELCA if you have liver problems.

If you are not sure, talk to your doctor before you are given ZEPZELCA.

Decreased blood cell counts

ZEPZELCA can cause severe and life-threatening myelosuppression (a condition in which the bone marrow cannot make enough blood cells). This can lead to febrile neutropenia (a decrease in neutrophils, white blood cells that fight infections, with fever) and a decrease in blood platelets, components that help the blood to clot. Your doctor will do blood tests to check your blood counts before you start treatment with ZEPZELCA and before each treatment cycle.

Tell your healthcare provider if you develop signs of infection such as:

- fever,
- chills,
- fatigue,
- body ache,
- cough;

or signs of low blood platelets such as:

- bruising,
- bleeding from gums or nose,
- blood in urine or stools
- cuts that bleed for a long time.

Liver problems

ZEPZELCA can cause an increase in liver enzymes which can be a sign of liver problems. Your doctor will do blood tests to check your liver function before and during treatment with ZEPZELCA.

Severe muscle problems (rhabdomyolysis).

ZEPZELCA may cause damage to your muscle which increase the level of an enzyme in your blood called creatine phosphokinase (CPK). You should seek medical attention immediately if you experience severe muscle pain or weakness.

Your doctor will do a blood test to check this enzyme levels before you start and periodically during treatment with ZEPZELCA.

Infusion leaking out of your vein (extravasation)

You should seek medical attention immediately if you notice that ZEPZELCA infusion leaks out of your vein while you are being given it, or if you notice any redness, swelling, itching or discomfort at the injection site at any time. It could lead to damage and death of your tissue cells around the injection site (tissue necrosis, see also section 4) which may require surgery.

Tumour Lysis Syndrome

ZEPZELCA can cause cancer cells to break down too quickly. When many cells break down at once, they release substances into the blood that can throw off the body's normal balance. This can harm the kidneys and may cause a life threatening problem such as abnormal heart rhythms and seizures.

Children and adolescents

Do not give this medicine to children and adolescents between the ages of 0 and 18 years because there is no relevant use of ZEPZELCA in the treatment of SCLC in this population.

Other medicines and ZEPZELCA

Tell your doctor if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription and herbal medicines.

The following medicines may reduce the effectiveness of ZEPZELCA by decreasing the level of ZEPZELCA in your body:

- phenytoin, phenobarbital or carbamazepine (for seizures);
- rifampicin, rifapentine and rifabutin (for tuberculosis);
- St. John's wort (*Hypericum perforatum*, a herbal remedy for depression and low mood).

The following medicines may increase the risk of side effects with ZEPZELCA since they increase the level of ZEPZELCA in your body:

- ketoconazole, itraconazole, posaconazole, fluconazole or voriconazole (for fungal infections);
- clarithromycin, erythromycin, telithromycin, ciprofloxacin (for bacterial infections);
- indinavir, lopinavir, nelfinavir, ritonavir, saquinavir, atazanavir, boceprevir, telaprevir (for HIV infection);
- aprepitant (a medicine used to prevent nausea and vomiting);
- ciclosporin (a medicine used to inhibit the defensive system of the body);
- verapamil, diltiazem (for hypertension);
- fluvoxamine (a medicine used to treat the depression).

The use of any of these medicines together with ZEPZELCA should be avoided, if possible.

Pregnancy, breastfeeding and fertility

ZEPZELCA should not be used during pregnancy.

This is because studies in animals have shown that ZEPZELCA may cause damage to the unborn baby. If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

You must stop breastfeeding before you start your treatment with ZEPZELCA and you must not begin breastfeeding again until your doctor has confirmed that it is safe to do so. This is because it is not known if ZEPZELCA pass into breast milk. Therefore a risk to newborns/infants cannot be ruled out.

If you are a woman who could become pregnant, you will need to have a pregnancy test before you start treatment with ZEPZELCA.

If you are a woman who could become pregnant, you must use effective contraception (birth control) during treatment with ZEPZELCA and for 7 months after your last dose.

If you are a man with a female partner who could become pregnant, you must use effective contraception during treatment with ZEPZELCA and for 4 months after your last dose.

Talk to your doctor or nurse about suitable methods of contraception for you and your partner.

ZEPZELCA may affect your ability to have a baby as it can cause damage to cells (cytotoxic) and may cause changes in genetic materials (mutagenic). Talk to your doctor for advice about options to preserve eggs or sperm before using ZEPZELCA.

Driving and using machines

During your treatment with ZEPZELCA you may experience tiredness, dizziness, vertigo and nausea. Do not drive or use any tools or machines if you are experiencing any of these side effects.

ZEPZELCA contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per dosage unit, that is to say essentially 'sodium-free'.

3. How to use ZEPZELCA

Treatment with ZEPZELCA should be started and supervised in a hospital or clinic by a healthcare professional experienced in the use of cancer medicines.

The dose of ZEPZELCA is based on your body surface area, which is calculated using your height and weight. The recommended dose is 3.2 mg/m² of body surface.

ZEPZELCA is given as an infusion (drip) into a vein over a period of one hour. To reduce the risk of infusion leaking out of your vein (extravasation) and inflammation of veins causing a blood clot (thrombophlebitis), this medicine should be given through a central venous line (a thin, flexible tube placed into a large vein, usually in the chest or neck, which allows medicines to be delivered directly into the bloodstream).

It is given once every 21 days. Before each dose your doctor will carefully test your blood counts, liver and kidney functions, and iron levels, based on the results of these tests your doctor may recommend dose delays to ensure that you receive the most appropriate dose of this medicine. Treatment can continue until the patient's cancer progresses or they experience serious side effects.

ZEPZELCA should be given after atezolizumab when both medicines are given on the same day.

If you develop serious side effects treatment may be withheld or permanently discontinued.

You will be given another medicine such as corticosteroids and serotonin antagonist before the treatment with ZEPZELCA to prevent feeling sick (nausea) and vomiting. If needed, the treatment with these medicines can be continued after the treatment with ZEPZELCA.

You will be given another medicine such as with granulocyte colony-stimulating factor (G-CSF) after the treatment with ZEPZELCA to prevent fever associated with lower-than-normal number of white blood cells (neutropenia).

If you stop receiving ZEPZELCA

Do not stop treatment with this medicine unless you have discussed this with your doctor. This is because stopping treatment may stop the effect of the medicine.

If you have any further questions on the use of this medicine, ask your doctor.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If you are not sure what the side effects below are, you should ask your doctor to explain them to you in more detail.

Tell your doctor immediately if you develop any of the following serious side effects:

Very common (may affect more than 1 in 10 people):

- difficulty breathing (dyspnoea).

Common (may affect up to 1 in 10 people):

- low levels of white blood cells with fever due to infection (febrile neutropenia).
- infusion leaking out of your vein (extravasation) while you are being given the medicine, causing redness, swelling, itchiness and discomfort at the infusion site.

It could cause tissue damage and death around the injection site (tissue necrosis), which may require surgery. Some of the symptoms or signs of extravasation may not be visible until several hours after it has occurred. There may be blistering, peeling and darkening of the skin over the site. It is possible for it to take a few days before the full extent of tissue damage is visible.

- lung infection (pneumonia) and cough.

Uncommon (may affect up to 1 in 100 people)

- blood poisoning (sepsis).

Frequency not known (cannot be estimated from the available data)

- breakdown of muscles often leading to kidney damage (rhabdomyolysis).
- rapid and massive destruction of cancer cells which release substances into the blood that can throw off the body's normal balance. This can harm the kidneys and may cause a life threatening problem such as abnormal heart rhythms and seizures (tumour lysis syndrome).

Talk to your doctor if you get any of the following other side effects

Very common (may affect more than 1 in 10 people):

- tiredness (fatigue),
- low levels of red blood cells which can cause tiredness and pale skin (anaemia),
- low levels of blood platelets which can lead to bleeding and bruising (thrombocytopenia),
- low levels of neutrophils, a type of white blood cell that fights infection (neutropenia),
- low levels of white blood cells (leukocytes),
- feeling sick (nausea),
- vomiting,
- diarrhoea,
- constipation,
- decreased appetite,
- pain in muscle and bones (musculoskeletal pain).

Common (may affect up to 1 in 10 people):

- high blood levels of creatinine, a sign of kidney problems,
- increased blood levels of liver enzymes (transaminases and gamma-glutamyl transferase), indicating problems with liver function.,
- increased level of creatine phosphokinase in the blood,
- infection of the parts of the body that collect and pass out urine (urinary tract infection),
- low levels of thyroid hormone (hypothyroidism) which can cause tiredness, weight gain, and skin and hair changes.
- joint pain (arthralgia),
- nerve damage in arms and legs causing pain or numbness, burning and tingling (peripheral neuropathy),
- low levels of a type of white blood cell called lymphocytes (lymphopenia),
- low blood levels of magnesium (hypomagnesaemia),
- low blood levels of calcium (hypocalcaemia),
- belly (abdominal) pain,
- headache,
- inflammation of a vein (phlebitis),
- inflammation of veins causing a blood clot to form and block one of your vein, leading to redness, swelling and pain in the area affected (thrombophlebitis),
- fever (pyrexia),
- swelling due to fluid retention (oedema),
- peripheral swelling,

- rash,
- itching (pruritus),
- infection,
- skin infection,
- taste disturbance (dysgeusia),
- pain or burning in the stomach, bloating, excessive belching or nausea (dyspepsia),
- inflammation of the lining of the mouth (stomatitis),
- cough or productive cough,
- inflammation in the lungs causing shortness of breath and cough (pneumonitis),
- mucosal inflammation, and
- weight decreased.

Uncommon (may affect up to 1 in 100 people)

- low levels of all types of blood cells (pancytopenia).

Reporting of side effects

If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store ZEPZELCA

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and the vial label after EXP. The expiry date refers to the last day of that month.

Store in a refrigerator (2°C – 8°C).

Information on storage of the reconstituted and diluted solutions is included in the section for healthcare professionals.

Do not use this medicine if you notice visible particles after the reconstitution or dilution of the medicine.

Any unused medicine or waste material should be disposed of in accordance with local requirements for cytotoxic medicines.

6. Contents of the pack and other information

What ZEPZELCA contains

ZEPZELCA 2 mg powder for concentrate for solution for infusion

- The active substance is lurbinectedin. Each vial contains 2 mg of lurbinectedin.
- The other excipients are sucrose, lactic acid and sodium hydroxide (see section 2).
-

ZEPZELCA 4 mg powder for concentrate for solution for infusion

- The active substance is lurbinectedin. Each vial contains 4 mg of lurbinectedin.
- The other excipients are sucrose, lactic acid and sodium hydroxide (see section 2).

What ZEPZELCA looks like and contents of the pack

ZEPZELCA is a powder for concentrate for solution for infusion.

ZEPZELCA 2 mg powder for concentrate for solution for infusion

The powder has a white to off-white colour and comes in a 20 mL glass vial with white coloured aluminium overseal.

Each carton contains one vial.

ZEPZELCA 4 mg powder for concentrate for solution for infusion

The powder has a white to off-white colour and comes in a 30 mL glass vial with blue coloured aluminium overseal.

Each carton contains one vial.

Marketing Authorisation Holder and Manufacturer

Pharma Mar, S.A.
Avda. de los Reyes 1
Polígono Industrial La Mina
28770 Colmenar Viejo (Madrid)
Spain
Tel: +34 91 846 60 00
Fax: +34 91 846 60 01

This leaflet was last revised in <{MM/YYYY}>.

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency web site: <https://www.ema.europa.eu>. There are also links to other websites about rare diseases and treatments.

The following information is intended for healthcare professionals only:

Instructions for use, preparation, handling and disposal

Appropriate procedures for proper handling and disposal of cytotoxic medicines must be followed.

You should have received training on the correct techniques to reconstitute and dilute ZEPZELCA and you should wear protective clothing including mask, goggles and gloves during the reconstitution and dilution. Accidental contact with the skin, eyes or mucous membranes must be treated immediately with copious amounts of water. You should not work with this medicine if you are pregnant.

Preparation for intravenous infusion

ZEPZELCA must be reconstituted and further diluted prior to infusion (see also section 3).
Appropriate aseptic techniques must be used.

ZEPZELCA must not be administered as a mixture with other medicines in the same infusion apart from the diluent. No incompatibilities have been observed between ZEPZELCA and polyolefin containers (polyethylene, polypropylene and mixtures), PVC (non-DEHP-containing), polyurethane and polyolefin infusion sets (polyethylene, polypropylene and polybutadiene), polyethersulfone in-line filters with pore sizes of 0.22 micron, and implantable venous access systems with titanium and plastic resin ports and with polyurethane or silicone intravenous catheters.

Infusion lines containing nylon membrane filters should not be used when the reconstituted ZEPZELCA solution is diluted with sodium chloride 9 mg/mL (0.9%) solution for infusion.

Instructions for reconstitution

Inject 8 mL (for 4 mg strength) or 4 mL (for 2 mg strength) of water for injections into the vial.

A syringe is used to inject the correct amount of water for injections into the vial. Shake the vial until complete dissolution. The reconstituted solution results in a clear, colourless or slightly yellowish solution, essentially free of visible particles.

This reconstituted solution contains 0.5 mg/mL of lurbinectedin which requires further dilution. It is for single use only.

Instructions for dilution

Calculate the required volume as follows:

$$\text{Volume (mL)} = \frac{\text{Body Surface Area (m}^2\text{)} \times \text{Individual Dose (mg/m}^2\text{)}}{0.5 \text{ mg/mL}}$$

Withdraw the appropriate amount of reconstituted solution from the vial.

If intravenous administration is to be made via a central venous line, add the reconstituted solution to an infusion bag containing at least 100 mL of diluent (sodium chloride 9 mg/mL (0.9%) solution for infusion or glucose 50 mg/mL (5%) solution for infusion)

If central venous access is not feasible and a peripheral venous line has to be used, add the reconstituted solution to an infusion bag containing at least 250 mL of diluent (sodium chloride 9 mg/mL (0.9%) solution for infusion or glucose 50 mg/mL (5%) solution for infusion).

Inspect the parenteral solution visually for particles prior to intravenous administration. Once the infusion is prepared, it should be administered immediately.

In-use stability of the solutions

Chemical and physical in-use stability has been demonstrated for 24 hours at either 2°C to 8°C or 25 °C.

From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C to 8°C, unless reconstitution / dilution has taken place in controlled and validated aseptic conditions. If reconstitution / dilution has taken place in controlled and validated aseptic conditions the prepared ready to administer product can be stored up to 24 hours at either +2°C to +8°C or +25 °C.

Disposal

Any unused medicine or waste material should be disposed of in accordance with local requirements for cytotoxic medicines.