

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

## **1. NAME OF THE MEDICINAL PRODUCT**

Zinforo 600 mg powder for concentrate for solution for infusion

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each vial contains ceftaroline fosamil acetic acid solvate monohydrate equivalent to 600 mg ceftaroline fosamil.

After reconstitution, 1 mL of the solution contains 30 mg of ceftaroline fosamil.

For the full list of excipients, see section 6.1.

## **3. PHARMACEUTICAL FORM**

Powder for concentrate for solution for infusion (powder for concentrate).

A pale yellowish-white to light yellow powder.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Zinforo is indicated for the treatment of the following infections in adults and children from the age of 2 months (see sections 4.4 and 5.1):

- Complicated skin and soft tissue infections (cSSTI)
- Community-acquired pneumonia (CAP)

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

### **4.2 Posology and method of administration**

#### Posology

The recommended durations of treatment are 5-14 days for cSSTI and 5-7 days for CAP.

**Table 1 Dosage in adults with normal renal function, creatinine clearance (CrCL) > 50 mL/min**

Indications	Posology (mg/infusion)	Infusion time (minutes)/Frequency
<u>Standard dose<sup>a</sup></u> Complicated skin and soft tissue infections (cSSTI) Community-acquired pneumonia (CAP)	600 mg	5 – 60 <sup>b</sup> /every 12 hours
<u>High dose<sup>b</sup></u> cSSTI confirmed or suspected to be caused by <i>S. aureus</i> with an MIC = 2 mg/L or 4 mg/L to ceftaroline <sup>c</sup>		120/every 8 hours

<sup>a</sup> For patients with supranormal renal clearance receiving the standard dose, an infusion time of 60 minutes may be preferable.

<sup>b</sup> Infusion times of less than 60 minutes and high dose recommendations are based on pharmacokinetic and pharmacodynamic analyses only. See sections 4.4 and 5.1.

<sup>c</sup> For treatment of *S. aureus* for which the ceftaroline MIC is  $\leq 1$  mg/L, the standard dose is recommended.

**Table 2 Dosage in paediatric patients with normal renal function, creatinine clearance (CrCL) > 50 mL/min**

Indications	Age group	Posology (mg/infusion)	Infusion time (minutes)/Frequency
<u>Standard dose<sup>a</sup></u> Complicated skin and soft tissue infections (cSSTI) Community-acquired pneumonia (CAP)	Adolescents aged from 12 to < 18 years with bodyweight $\geq 33$ kg	600 mg	5–60 <sup>b</sup> /every 12 hours
	Adolescents aged from 12 years to < 18 years bodyweight < 33 kg <b>and</b> children $\geq 2$ years to < 12 years	12 mg/kg to a maximum of 400 mg	5–60 <sup>b</sup> /every 8 hours
	Infants $\geq 2$ months to < 2 years	8 mg/kg	5–60 <sup>b</sup> /every 8 hours
<u>High dose<sup>b</sup></u> cSSTI confirmed or suspected to be caused by <i>S. aureus</i> with an MIC = 2 mg/L or 4 mg/L to ceftaroline <sup>c</sup>	Adolescents aged from 12 to < 18 years with bodyweight $\geq 33$ kg	600 mg	120/every 8 hours

<sup>a</sup> For patients with supranormal renal clearance receiving the standard dose, an infusion time of 60 minutes may be preferable.

<sup>b</sup> Infusion times of less than 60 minutes and high dose recommendations are based on pharmacokinetic and pharmacodynamic analyses only. See sections 4.4 and 5.1.

<sup>c</sup> For treatment of *S. aureus* for which the ceftaroline MIC is  $\leq 1$  mg/L, the standard dose is recommended.

### Special populations

#### *Elderly*

No dosage adjustment is required for the elderly with creatinine clearance values > 50 mL/min (see section 5.2).

#### *Renal impairment*

The dose should be adjusted when creatinine clearance (CrCL) is  $\leq 50$  mL/min, as shown in Tables 3 and 4 (see section 5.2). The recommended durations of treatment are 5-14 days for cSSTI and 5-7 days for CAP.

**Table 3 Dosage in adults with impaired renal function, creatinine clearance (CrCL) ≤ 50 mL/min**

Indications	Creatinine clearance (mL/min) <sup>a</sup>	Posology (mg/infusion)	Infusion time (minutes)/Frequency
<u>Standard dose</u> Complicated skin and soft tissue infections (cSSTI) Community-acquired pneumonia (CAP)	> 30 to ≤ 50	400 mg	5–60 <sup>c</sup> /every 12 hours
	≥ 15 to ≤ 30	300 mg	
	ESRD, including haemodialysis <sup>b</sup>	200 mg	
<u>High dose<sup>c</sup></u> cSSTI confirmed or suspected to be caused by <i>S. aureus</i> with an MIC = 2 mg/L or 4 mg/L to ceftaroline <sup>d</sup>	> 30 to ≤ 50	400 mg	120/every 8 hours
	≥ 15 to ≤ 30	300 mg	
	ESRD, including haemodialysis <sup>b</sup>	200 mg	

<sup>a</sup> Calculated using the Cockcroft-Gault formula for adults. Dose is based on CrCL. CrCL should be closely monitored and the dose adjusted according to changing renal function.

<sup>b</sup> Ceftaroline is haemodialyzable; thus Zinforo should be administered after haemodialysis on haemodialysis days.

<sup>c</sup> Infusion times of less than 60 minutes and high dose recommendations are based on pharmacokinetic and pharmacodynamic analyses only. See sections 4.4 and 5.1.

<sup>d</sup> For treatment of *S. aureus* for which the ceftaroline MIC is ≤ 1 mg/L, the standard dose is recommended.

Dose recommendations for children and adolescents are based on pharmacokinetic (PK) modelling.

There is insufficient information to recommend dosage adjustments in adolescents aged from 12 to < 18 years with bodyweight < 33 kg and in children aged from 2 to 12 years with End-stage renal disease (ESRD).

There is insufficient information to recommend dosage adjustments in children aged from 2 months to < 2 years with moderate or severe renal impairment or ESRD.

**Table 4 Dosage in paediatric patients with impaired renal function, creatinine clearance (CrCL) ≤ 50 mL/min**

Indications	Age group	Creatinine clearance (mL/min) <sup>a</sup>	Posology (mg/infusion)	Infusion time (minutes)/Frequency
<u>Standard dose</u> Complicated skin and soft tissue infections (cSSTI)	Adolescents aged from 12 to < 18 years with bodyweight ≥ 33 kg	> 30 to ≤ 50	400 mg	5–60 <sup>c</sup> /every 12 hours
		≥ 15 to ≤ 30	300 mg	
		ESRD, including haemodialysis <sup>b</sup>	200 mg	
Community-acquired pneumonia (CAP)	Adolescents aged from 12 years to < 18 years bodyweight < 33 kg <b>and</b> children ≥ 2 years to < 12 years	> 30 to ≤ 50	8 mg/kg to a maximum of 300 mg	5–60 <sup>c</sup> /every 8 hours
		≥ 15 to ≤ 30	6 mg/kg to a maximum of 200 mg	
<u>High dose<sup>c</sup></u>	Adolescents aged from 12 to	> 30 to ≤ 50	400 mg	120/every 8 hours
		≥ 15 to ≤ 30	300 mg	

Indications	Age group	Creatinine clearance (mL/min) <sup>a</sup>	Posology (mg/infusion)	Infusion time (minutes)/Frequency
cSSTI confirmed or suspected to be caused by <i>S. aureus</i> with an MIC = 2 mg/L or 4 mg/L to ceftaroline <sup>d</sup>	< 18 years with bodyweight ≥ 33 kg	ESRD, including haemodialysis <sup>b</sup>	200 mg	

<sup>a</sup> Calculated using the Cockcroft-Gault formula for adolescents aged from 12 to < 18 years with bodyweight ≥ 33 kg and Schwartz formula for children and adolescents with bodyweight less than 33 kg. Dose is based on CrCL. CrCL should be closely monitored and the dose adjusted according to changing renal function.

<sup>b</sup> Ceftaroline is haemodialyzable; thus Zinfofo should be administered after haemodialysis on haemodialysis days.

<sup>c</sup> Infusion times of less than 60 minutes and high dose recommendations are based on pharmacokinetic and pharmacodynamic analyses only. See sections 4.4 and 5.1.

<sup>d</sup> For treatment of *S. aureus* for which the ceftaroline MIC is ≤ 1 mg/L, the standard dose is recommended.

#### *Hepatic impairment*

No dosage adjustment is considered necessary in patients with hepatic impairment (see section 5.2).

#### *Paediatric population*

The safety and efficacy of Zinfofo in children below the age of 2 months have not yet been established. No data are available.

#### Method of administration

Intravenous use. Zinfofo is administered by intravenous infusion over 5 to 60 minutes for standard dose or 120 minutes for high dose (for cSSTI caused by *S. aureus* with MIC of 2 or 4 mg/L to ceftaroline) in infusion volumes of 50 mL, 100 mL or 250 mL (see section 6.6). Infusion related reactions (such as phlebitis) can be managed by prolonging the infusion duration.

Infusion volumes for paediatric patients will vary according to the weight of the child. The infusion solution concentration during preparation and administration should not exceed 12 mg/mL ceftaroline fosamil.

For instructions on reconstitution and dilution of the medicinal product before administration, see section 6.6.

### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Hypersensitivity to the cephalosporin class of antibacterials.

Immediate and severe hypersensitivity (e.g. anaphylactic reaction) to any other type of beta-lactam antibacterial agent (e.g. penicillins or carbapenems).

### **4.4 Special warnings and precautions for use**

#### Hypersensitivity reactions

Serious and occasionally fatal hypersensitivity reactions are possible (see sections 4.3 and 4.8).

Patients who have a history of hypersensitivity to cephalosporins, penicillins or other beta-lactam antibacterials may also be hypersensitive to ceftaroline fosamil. Ceftaroline should be used with caution in patients with a history of non-severe hypersensitivity reactions to any other beta-lactam antibiotics (e.g. penicillins or carbapenems). If a severe allergic reaction occurs during treatment with Zinfofo, the medicinal product should be discontinued and appropriate measures taken.

#### Clostridium difficile-associated diarrhoea

Antibacterial-associated colitis and pseudomembranous colitis have been reported with ceftaroline fosamil and may range in severity from mild to life threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhoea during or subsequent to the administration of ceftaroline fosamil (see section 4.8). In such circumstance, the discontinuation of therapy with ceftaroline fosamil and the use of supportive measures together with the administration of specific treatment for *Clostridium difficile* should be considered.

#### Non-susceptible organisms

Superinfections may occur during or following treatment with Zinfofo.

#### Patients with pre-existing seizure disorder

Seizures have occurred in toxicology studies at 7-25 times human ceftaroline C<sub>max</sub> levels (see section 5.3). Clinical study experience with ceftaroline fosamil in patients with pre-existing seizure disorders is very limited. Therefore, Zinfofo should be used with caution in this patient population.

#### Direct antiglobulin test (Coombs test) seroconversion and potential risk of haemolytic anaemia

The development of a positive direct antiglobulin test (DAGT) may occur during treatment with cephalosporins. The incidence of DAGT seroconversion in patients receiving ceftaroline fosamil was 11.2% in the five pooled pivotal studies with administration every 12 hours (600 mg administered over 60 minutes every 12 hours) and 32.3% in a study in patients receiving ceftaroline fosamil every 8 hours (600 mg administered over 120 minutes every 8 hours), (see section 4.8). In clinical studies there was no evidence of haemolysis in patients who developed a positive DAGT on treatment. However, the possibility that haemolytic anaemia may occur in association with cephalosporins including Zinfofo treatment cannot be ruled out. Patients experiencing anaemia during or after treatment with Zinfofo should be investigated for this possibility.

#### Limitations of the clinical data

There is no experience with ceftaroline in the treatment of CAP in the following patient groups: the immunocompromised, patients with severe sepsis/septic shock, severe underlying lung disease, those with PORT Risk Class V, and/or CAP requiring ventilation at presentation, CAP due to methicillin-resistant *S. aureus* or patients requiring intensive care. Caution is advised when treating such patients.

There is no experience with ceftaroline in the treatment of cSSTI in the following patient groups: the immunocompromised, patients with severe sepsis/septic shock, necrotizing fasciitis, perirectal abscess and patients with third degree and extensive burns. There is limited experience in treating patients with diabetic foot infections. Caution is advised when treating such patients.

There are limited clinical trial data on the use of ceftaroline to treat cSSTI caused by *S. aureus* with an MIC of > 1 mg/L. The recommended dosages of Zinfofo shown in Tables 1 to 4 for the treatment of cSSTI caused by *S. aureus* with ceftaroline MIC of 2 or 4 mg/L are based on pharmacokinetic-pharmacodynamic modelling and simulation (see sections 4.2 and 5.1). Zinfofo should not be used to treat cSSTI due to *S. aureus* for which the ceftaroline MIC is > 4 mg/L.

Infusion times of less than 60 minutes are based on pharmacokinetic and pharmacodynamic analyses only.

### **4.5 Interaction with other medicinal products and other forms of interaction**

No clinical drug-drug interaction studies have been conducted with ceftaroline fosamil.

The interaction potential of ceftaroline or ceftaroline fosamil on medicinal products metabolised by CYP450 enzymes is expected to be low since they are not inhibitors nor inducers of CYP450 enzymes *in vitro*. Ceftaroline or ceftaroline fosamil are not metabolised by CYP450 enzymes *in vitro*, therefore

co-administered CYP450 inducers or inhibitors are unlikely to influence the pharmacokinetics of ceftaroline.

Ceftaroline is neither a substrate, nor an inhibitor of renal uptake transporters (OCT2, OAT1, and OAT3) *in vitro*. Therefore, interactions of ceftaroline with medicinal products that are substrates or inhibitors (e.g. probenecid) of these transporters would not be expected.

#### Paediatric population

As with adults, the interaction potential is expected to be low in paediatrics.

### **4.6 Fertility, pregnancy and lactation**

#### Pregnancy

There are no or limited amount of data from the use of ceftaroline fosamil in pregnant women. Animal studies conducted in rat and rabbit do not indicate harmful effects with respect to reproductive toxicity at exposures similar to therapeutic concentrations. Following administration throughout pregnancy and lactation in the rat, there was no effect on pup birth weight or growth, although minor changes in foetal weight and delayed ossification of the interparietal bone were observed when ceftaroline fosamil was administered during organogenesis (see section 5.3).

As a precautionary measure, it is preferable to avoid the use of Zinforo during pregnancy unless the clinical condition of the woman requires treatment with an antibiotic with Zinforo's antibacterial profile.

#### Breast-feeding

It is unknown whether ceftaroline fosamil or ceftaroline is excreted in human milk. A risk to the newborns/infants cannot be excluded. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from Zinforo therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

#### Fertility

The effects of ceftaroline fosamil on fertility on humans have not been studied. Animal studies with ceftaroline fosamil do not indicate harmful effects with respect to fertility (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

Undesirable effects e.g. dizziness may occur and this may have an effect on the ability to drive and use of machines (see section 4.8).

### **4.8 Undesirable effects**

#### Summary of the safety profile

The most common adverse reactions occurring in  $\geq 3\%$  of approximately 3242 patients treated with Zinforo in clinical studies were diarrhoea, headache, nausea, and pruritus, and were generally mild or moderate in severity. CDAD and severe hypersensitivity reactions may also occur.

A greater incidence of rash in Asian patients (see below) and a greater incidence of DAGT seroconversion (see section 4.4) were observed in a study of adult patients with cSSTI conducted with Zinforo 600 mg administered over 120 minutes every 8 hours.

#### Tabulated list of adverse reactions

The following adverse reactions have been identified during clinical trials and post-marketing experience with Zinforo. Adverse reactions are classified according to System Organ Class and frequency. Frequency categories are derived according to the following conventions: very common

( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1,000$ ), not known (cannot be estimated from the available data).

**Table 5 Frequency of adverse reactions by system organ class from clinical trials and post-marketing experience**

System organ class	Very common	Common	Uncommon	Rare
Infections and infestations			<i>Clostridium difficile</i> colitis (see section 4.4)	
Blood and lymphatic system disorders			Anaemia, leucopenia, neutropenia, thrombocytopenia, prothrombin time (PT) prolonged, activated partial thromboplastin time (aPTT) prolonged, international normalized ratio (INR) increased	Agranulocytosis, eosinophilia
Immune system disorders		Rash, pruritus	Anaphylaxis (see sections 4.3 and 4.4), hypersensitivity (e.g. urticaria, lip and face swelling) (see sections 4.3 and 4.4)	
Nervous system disorders		Headache, dizziness		
Vascular disorders		Phlebitis		
Gastrointestinal disorders		Diarrhoea, nausea, vomiting, abdominal pain		
Hepatobiliary disorders		Increased transaminases		
Renal and urinary disorders			Blood creatinine increased	

System organ class	Very common	Common	Uncommon	Rare
General disorders and administration site conditions		Pyrexia, infusion site reactions (erythema, phlebitis, pain)		
Investigations	Coombs Direct Test Positive (see section 4.4)			

### Description of selected adverse reactions

#### *Rash*

Rash was observed at a common frequency in both the pooled Phase III studies in cSSTI with administration of Zinforo every 12 hours (600 mg administered over 60 minutes every 12 hours) and the study in cSSTI with administration every 8 hours (600 mg administered over 120 minutes every 8 hours). However, the frequency of rash in the subgroup of Asian patients receiving Zinforo every 8 hours was very common (18.5%).

#### Paediatric population

The safety assessment in children is based on the safety data from 2 trials in which 227 paediatric patients aged from 2 months to 17 years with cSSTI or CAP received Zinforo. Overall, the safety profile in these 227 children was similar to that observed in the adult population.

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

## **4.9 Overdose**

Limited data in patients receiving higher than recommended Zinforo dosages show similar adverse reactions as observed in the patients receiving recommended dosages. Relative overdosing could occur in patients with moderate renal impairment. Treatment of overdose should follow standard medical practice;

Ceftaroline can be removed by haemodialysis; over a 4 hour dialysis period, approximately 74% of a given dose was recovered in the dialysate.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Antibacterials for systemic use, other cephalosporins and penems, ATC code: J01DI02

The active moiety after Zinforo administration is ceftaroline.

### Mechanism of action

*In vitro* studies have shown that ceftaroline is bactericidal and able to inhibit bacterial cell wall synthesis in methicillin-resistant *Staphylococcus aureus* (MRSA) and penicillin non-susceptible *Streptococcus pneumoniae* (PNSP) due to its affinity for the altered penicillin-binding proteins (PBPs) found in these organisms. As a result, minimum inhibitory concentrations (MICs) of ceftaroline against a proportion of these organisms tested fall into the susceptible range (see Resistance section below).

### Resistance

Ceftaroline is not active against strains of *Enterobacteriaceae* producing extended-spectrum beta-lactamases (ESBLs) from the TEM, SHV or CTX-M families, serine carbapenemases (such as KPC), class B metallo-beta-lactamases or class C (AmpC) cephalosporinases. Organisms that express these enzymes and which are therefore resistant to ceftaroline occur at very variable rates between countries and between healthcare facilities within countries. If ceftaroline is commenced before susceptibility test results are available then local information on the risk of encountering organisms that express these enzymes should be taken into consideration. Resistance may also be mediated by bacterial impermeability or drug efflux pump mechanisms. One or more of these mechanisms may co-exist in a single bacterial isolate.

### Interaction with other antibacterial agents

*In vitro* studies have not demonstrated any antagonism between ceftaroline in combination with other commonly used antibacterial agents (e.g. amikacin, azithromycin, aztreonam, daptomycin, levofloxacin, linezolid, meropenem, tigecycline, and vancomycin).

### Susceptibility testing breakpoints

The European Committee on Antimicrobial Susceptibility Testing (EUCAST) breakpoints for susceptibility testing are presented below.

Organisms	MIC breakpoints (mg/L)	
	Susceptible ( $\leq$ S)	Resistant (R $>$ )
<i>Staphylococcus aureus</i>	1 <sup>1</sup>	$>2^2$
<i>Streptococcus pneumoniae</i>	0.25	0.25
<i>Streptococcus</i> Groups A, B, C, G	Note <sup>3</sup>	Note <sup>3</sup>
<i>Haemophilus influenzae</i>	0.03	0.03
<i>Enterobacteriaceae</i>	0.5	0.5

1. Refers to dosing of adults or adolescents (from 12 years and 33 kg) with ceftaroline every 12 hours using 1-hour infusions (see section 4.2). Note that: There are no clinical trial data regarding the use of ceftaroline to treat CAP due to *S. aureus* with ceftaroline MICs  $> 1$  mg/L
2. Refers to dosing of adults or adolescents (from 12 years and 33 kg) with ceftaroline every 8 hours using 2-hour infusions to treat cSSTI (see section 4.2). *S. aureus* with ceftaroline MICs  $\geq 4$  mg/L are rare. PK-PD analyses suggest that dosing of adults or adolescents (from 12 years and 33 kg) with ceftaroline every 8 hours using 2-hour infusions may treat cSSTI due to *S. aureus* for which the ceftaroline MIC is 4 mg/L.
3. Infer susceptibility from susceptibility to benzylpenicillin.

### Pharmacokinetic/pharmacodynamic relationship

As with other beta-lactam antimicrobial agents, the percent time above the minimum inhibitory concentration (MIC) of the infecting organism over the dosing interval (%T  $>$  MIC) has been shown to be the parameter that best correlates with the efficacy of ceftaroline.

### Clinical efficacy against specific pathogens

Efficacy has been demonstrated in clinical studies against the pathogens listed under each indication that were susceptible to ceftaroline *in vitro*.

#### *Complicated skin and soft tissue infections*

##### Gram-positive micro-organisms

- *Staphylococcus aureus* (including methicillin-resistant strains)
- *Streptococcus pyogenes*
- *Streptococcus agalactiae*
- *Streptococcus anginosus* group (includes *S. anginosus*, *S. intermedius*, and *S. constellatus*)
- *Streptococcus dysgalactiae*

##### Gram-negative micro-organisms

- *Escherichia coli*
- *Klebsiella pneumoniae*
- *Klebsiella oxytoca*
- *Morganella morganii*

#### *Community-acquired pneumonia*

No cases of CAP due to MRSA were enrolled into the studies. The available clinical data cannot substantiate efficacy against penicillin non-susceptible strains of *S. pneumoniae*.

##### Gram-positive micro-organisms

- *Streptococcus pneumoniae*
- *Staphylococcus aureus* (methicillin-susceptible strains only)

##### Gram-negative micro-organisms

- *Escherichia coli*
- *Haemophilus influenzae*
- *Haemophilus parainfluenzae*
- *Klebsiella pneumoniae*

#### Antibacterial activity against other relevant pathogens

Clinical efficacy has not been established against the following pathogens although *in vitro* studies suggest that they would be susceptible to ceftaroline in the absence of acquired mechanisms of resistance:

#### *Anaerobic micro-organisms*

##### Gram-positive micro-organisms

- *Peptostreptococcus* spp.

##### Gram-negative micro-organisms

- *Fusobacterium* spp.

*In vitro* data indicate that the following species are not susceptible to ceftaroline:

- *Chlamydophila* spp.
- *Legionella* spp.
- *Mycoplasma* spp.
- *Proteus* spp.
- *Pseudomonas aeruginosa*

#### Paediatric population

The European Medicines Agency has deferred the obligation to submit the results of studies with Zinforo in the paediatric population aged birth to < 2 months (see section 4.2 for information on paediatric use).

## 5.2 Pharmacokinetic properties

The  $C_{max}$  and AUC of ceftaroline increase approximately in proportion to dose within the single dose range of 50 to 1000 mg. No appreciable accumulation of ceftaroline is observed following multiple intravenous infusions of 600 mg every 8 or 12 hours in healthy adults with CrCL > 50 mL/min.

### Distribution

The plasma protein binding of ceftaroline is low (approximately 20%) and ceftaroline is not distributed into erythrocytes. The median steady-state volume of distribution of ceftaroline in healthy adult males following a single 600 mg intravenous dose of radiolabelled ceftaroline fosamil was 20.3 l, similar to the volume of extracellular fluid.

### Biotransformation

Ceftaroline fosamil (prodrug) is converted into the active ceftaroline in plasma by phosphatase enzymes and concentrations of the prodrug are measurable in plasma primarily during intravenous infusion. Hydrolysis of the beta-lactam ring of ceftaroline occurs to form the microbiologically inactive, open-ring metabolite, ceftaroline M-1. The mean plasma ceftaroline M-1 to ceftaroline AUC ratio following a single 600 mg intravenous infusion of ceftaroline fosamil in healthy subjects is approximately 20-30%.

In pooled human liver microsomes, metabolic turnover was low for ceftaroline, indicating that ceftaroline is not metabolised by hepatic CYP450 enzymes.

### Elimination

Ceftaroline is primarily eliminated by the kidneys. Renal clearance of ceftaroline is approximately equal, or slightly lower than the glomerular filtration rate in the kidney, and *in vitro* transporter studies indicate that active secretion does not contribute to the renal elimination of ceftaroline.

The mean terminal elimination half-life of ceftaroline in healthy adults is approximately 2.5 hours.

Following the administration of a single 600 mg intravenous dose of radiolabelled ceftaroline fosamil to healthy male adults, approximately 88% of radioactivity was recovered in urine and 6% in faeces.

### Special populations

#### *Renal impairment*

Dosage adjustments are required in adults, adolescents and children with CrCL  $\leq$  50 mL/min (see section 4.2).

There is insufficient information to recommend dosage adjustments in adolescents with ESRD aged from 12 to < 18 years and with bodyweight < 33 kg and in children with ESRD aged from 2 to < 12 years. There is insufficient information to recommend dosage adjustments in children aged < 2 years with moderate or severe renal impairment or ESRD.

#### *Hepatic impairment*

The pharmacokinetics of ceftaroline in patients with hepatic impairment has not been established. As ceftaroline does not appear to undergo significant hepatic metabolism, the systemic clearance of ceftaroline is not expected to be significantly affected by hepatic impairment. Therefore, no dosage adjustment is recommended for patients with hepatic impairment.

#### *Elderly*

Following administration of a single 600 mg intravenous dose of ceftaroline fosamil, the pharmacokinetics of ceftaroline were similar between healthy elderly subjects ( $\geq 65$  years of age), and healthy young adult subjects (18-45 years of age). There was a 33% increase in  $AUC_{0-\infty}$  in the elderly that was mainly attributable to age-related changes in renal function. Zinforo dose adjustment is not required in elderly patients with creatinine clearance above 50 mL/min.

#### *Paediatric population*

Dose adjustments are required for children aged from 2 months to  $< 12$  years and for adolescents aged 12 to  $< 18$  years with bodyweight  $< 33$  kg (see section 4.2). The safety and efficacy of Zinforo in children aged birth to  $< 2$  months have not been established.

### **5.3 Preclinical safety data**

The kidney was the primary target organ of toxicity in both the monkey and rat. Histopathologic findings included pigment deposition and inflammation of the tubular epithelium. Renal changes were not reversible but were reduced in severity following a 4 week recovery period.

Convulsions have been observed at relatively high exposures during single and multi-dose studies in both the rat and monkey ( $\geq 7$  times to the estimated ceftaroline  $C_{max}$  level of a 600 mg twice a day).

Other important toxicologic findings noted in the rat and monkey included histopathologic changes in the bladder and spleen.

#### Genetic toxicology

Ceftaroline fosamil and ceftaroline were clastogenic in an *in vitro* chromosomal aberration assay, however there was no evidence of mutagenic activity in an Ames test, mouse lymphoma and unscheduled DNA synthesis assay. Furthermore, *in vivo* micronucleus assays in rat and mouse were negative. Carcinogenicity studies have not been conducted.

#### Reproductive toxicology

Overall, no adverse effects on fertility or post-natal development were observed in the rat at up to 5 times the observed clinical exposure. When ceftaroline was administered during organogenesis, minor changes in foetal weight and delayed ossification of the interparietal bone were observed in the rat at exposures below that observed clinically. However, when ceftaroline was administered throughout pregnancy and lactation, there was no effect on pup weight or growth. Ceftaroline administration to pregnant rabbits resulted in an increased foetal incidence of angulated hyoid alae, a common skeletal variation in rabbit fetuses, at exposures similar to those observed clinically.

#### Juvenile toxicity

Intravenous bolus dosing of ceftaroline fosamil to suckling rats from post-natal day 7 to 20 was well tolerated at plasma exposures approximately 2-fold higher than those for paediatric patients. Renal cortical cysts were observed in all groups, including controls, on PND50. The cysts involved a small portion of the kidney and occurred in the absence of significant changes in either renal function or urinary parameters. Therefore, these findings were not considered to be adverse.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Arginine

### **6.2 Incompatibilities**

This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

### **6.3 Shelf life**

Dry powder: 3 years

#### After reconstitution:

The reconstituted vial should be diluted immediately.

#### After dilution:

Once the intravenous solution is prepared with diluents listed in section 6.6 it should be administered within 6 hours of preparation. The chemical and physical in-use stability has been demonstrated for up to 24 hours at 2-8°C. Once removed from refrigeration to room temperature, the diluted product must be used within 6 hours.

From a microbiological point of view, the medicinal product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 to 8°C, unless reconstitution/dilution has taken place in controlled and validated aseptic conditions.

### **6.4 Special precautions for storage**

Store below 30°C.

Store in the original package in order to protect from light.

For storage conditions after reconstitution of the medicinal product, see section 6.3.

### **6.5 Nature and contents of container**

20 mL glass vial (Type 1) closed with a rubber (halobutyl) stopper and aluminium seal with flip-off cap.

The medicinal product is supplied in packs of 10 vials.

### **6.6 Special precautions for disposal and other handling**

The powder must be reconstituted with water for injections and the resulting concentrate must then be immediately diluted prior to use. The reconstituted solution is a pale yellow solution that is free of any particles.

Standard aseptic techniques should be used for solution preparation and administration.

Zinforo powder should be reconstituted with 20 mL of sterile water for injections. The resulting solution should be shaken prior to being transferred to an infusion bag or bottle containing either sodium chloride 9 mg/mL (0.9%) solution for injection, dextrose 50 mg/mL (5%) solution for injection, sodium chloride 4.5 mg/mL and dextrose 25 mg/mL solution for injection (0.45% sodium chloride and 2.5% dextrose) or Lactated Ringer's solution. A 250 mL, 100 mL or 50 mL infusion bag can be used to prepare the infusion, based on the patient's volume requirements. The total time interval between starting reconstitution and completing preparation of the intravenous infusion should not exceed 30 minutes.

Infusion volumes for paediatric patients will vary according to the weight of the child. The infusion solution concentration during preparation and administration should not exceed 12 mg/mL ceftaroline fosamil.

Each vial is for single use only.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

**7. MARKETING AUTHORISATION HOLDER**

Pfizer Ireland Pharmaceuticals  
Operations Support Group  
Ringaskiddy, County Cork  
Ireland

**8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/12/785/001

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 23 August 2012

Date of the latest renewal: 24 April 2017

**10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

## **ANNEX II**

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## **A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer(s) responsible for batch release

ACS Dobfar S.p.A.  
Nucleo Industriale S. Atto  
64100 Teramo  
Italy

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to medical prescription.

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

### **• Periodic Safety Update Reports**

The requirements for submission of periodic safety update reports for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

### **• Risk Management Plan (RMP)**

The MAH shall perform the pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2. of the Marketing Authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as a result of an important (pharmacovigilance or risk minimisation) milestone being reached.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Zinfo 600 mg powder for concentrate for solution for infusion  
ceftaroline fosamil

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each vial contains ceftaroline fosamil acetic acid solvate monohydrate equivalent to 600 mg  
ceftaroline fosamil.

**3. LIST OF EXCIPIENTS**

Arginine

**4. PHARMACEUTICAL FORM AND CONTENTS**

Powder for concentrate for solution for infusion.  
10 vials

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
Intravenous use.  
For single use only.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store below 30°C.  
Store in the original package in order to protect from light.  
See leaflet for shelf life after reconstitution.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Pfizer Ireland Pharmaceuticals  
Operations Support Group  
Ringaskiddy, County Cork  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/12/785/001

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Justification for not including Braille accepted.

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC:  
SN:  
NN:

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**VIAL LABEL**

**1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

Zinfo 600 mg powder for concentrate  
ceftaroline fosamil  
IV

**2. METHOD OF ADMINISTRATION**

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

**6. OTHER**

**B. PACKAGE LEAFLET**

## **Package leaflet: Information for the user**

### **Zinforo 600 mg powder for concentrate for solution for infusion** Ceftaroline fosamil

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or nurse.
- If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet:**

1. What Zinforo is and what it is used for
2. What you need to know before you use Zinforo
3. How to use Zinforo
4. Possible side effects
5. How to store Zinforo
6. Contents of the pack and other information

#### **1. What Zinforo is and what it is used for**

##### **What Zinforo is**

Zinforo is an antibiotic medicine that contains the active substance ceftaroline fosamil. It belongs to a group of medicines called ‘cephalosporin antibiotics.’

##### **What Zinforo is used for**

Zinforo is used to treat children from the age of 2 months and adults with:

- infections of the skin and the tissues below the skin
- an infection of the lungs called ‘pneumonia’

##### **How Zinforo works**

Zinforo works by killing certain bacteria, which can cause serious infections.

#### **2. What you need to know before you use Zinforo**

##### **Do not use Zinforo:**

- If you are allergic to ceftaroline fosamil or any of the other ingredients of this medicine (listed in section 6).
- If you are allergic to other cephalosporin antibiotics
- If you have had previous severe allergic reactions to other antibiotics like penicillin or carbapenem.

Do not use Zinforo if any of the above applies to you. If you are not sure, talk to your doctor or nurse before using Zinforo.

##### **Warnings and precautions**

Talk to your doctor or nurse before using Zinforo:

- If you have kidney problems (your doctor may have to prescribe a lower dose)
- If you have ever had fits (seizures or convulsions)
- If you have ever had any non-severe allergic reactions to other antibiotics like penicillin or carbapenem

- If you have had severe diarrhoea whilst taking antibiotics in the past

You may get another infection caused by another bacteria during or following treatment with Zinforo.

#### Lab Test

You may develop an abnormal lab test (called Coombs test) that looks for certain antibodies which may act against your red blood cells. If the level of your red blood cells fall your doctor may check to see if these antibodies have caused this.

If any of the above apply to you (or you are not sure), talk to your doctor or nurse before using Zinforo.

#### Children

Zinforo should not be used in children below the age of 2 months as not enough data exist in these populations.

#### **Other medicines and Zinforo**

Tell your doctor or nurse if you are using, have recently used or might use any other medicines.

#### **Pregnancy and breast-feeding**

Tell your doctor before using Zinforo if you are pregnant. Do not use this medicine during pregnancy unless your doctor has told you to.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

#### **Driving and using machines**

Zinforo may cause side effects such as dizziness. This may impair your ability to drive or operate machinery.

### **3. How to use Zinforo**

Zinforo will be given to you by a doctor or nurse.

#### **How much to use**

The usual recommended dose for adults is 600 mg every 12 hours. Your doctor may increase your dose to 600 mg every 8 hours for some infections. The usual recommended dose for children depends on the age and weight of the child and is given every 8 or 12 hours. It is given as a drip into a vein lasting 5 to 60 minutes if you receive the usual dose or 120 minutes if you receive an increased dose.

A course of treatment usually lasts for 5 to 14 days for skin infections and 5 to 7 days for pneumonia.

#### **Patients with kidney problems**

If you have kidney problems your doctor may lower your dose because Zinforo is removed from your body by the kidneys.

#### **If you use more Zinforo than you should**

If you think you have been given too much Zinforo, tell your doctor or nurse straight away.

#### **If you miss a dose of Zinforo**

If you think you have missed a dose, tell your doctor or nurse straight away.

If you have any further questions on the use of this medicine, ask your doctor or nurse.

#### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. The following side effects may happen with this medicine:

##### **Tell your doctor straight away if you get these symptoms as you may need urgent medical treatment:**

- Sudden swelling of your lips, face, throat or tongue; a severe rash; and, swallowing or breathing problems. These may be signs of a severe allergic reaction (anaphylaxis) and may be life-threatening;
- Diarrhoea that becomes severe or does not go away or stool that contains blood or mucus during or after treatment with Zinfo. In this situation, you should not take medicines that stop or slow bowel movement.

##### **Very common (may affect more than 1 in 10 people)**

- Changes in a blood test called a 'Coombs test' commonly seen in patients receiving this type of antibiotic. This test looks for certain antibodies which may act against your red blood cells.

##### **Common (may affect up to 1 in 10 people)**

- Fever
- Headache
- Feeling dizzy
- Itching, skin rash
- Diarrhoea, stomach pain
- Feeling sick (nausea) or being sick (vomiting)
- More enzymes produced by your liver (as shown in blood tests)
- Pain and irritation of the veins
- Redness, pain or swelling where the injection was given.

##### **Uncommon (may affect up to 1 in 100 people)**

- Anaemia
- Raised itchy rash (hives)
- An increase in the level of creatinine in your blood. Creatinine shows how well your kidneys are working.
- Bleeding or bruising more than usual. This may be because the level of platelets in your blood has dropped.
- Changes in tests which measure how well your blood clots.
- A decrease in the total number of white blood cells, or a certain type of white blood cells in your blood (leucopenia and neutropenia).

##### **Rare (may affect up to 1 in 1,000 people)**

- A significant decrease in the number of certain white blood cells in your blood (agranulocytosis). You may experience fever, flu-like symptoms, sore throat, or any other infection which may be serious.
- An increase in the number of certain white blood cells in your blood (eosinophilia).

##### **Reporting of side effects**

If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## 5. How to store Zinforo

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the container. The expiry date refers to the last day of that month.

Store below 30°C.

Store in the original package in order to protect from light.

Medicines should not be disposed of via wastewater or household waste. The hospital will dispose of any waste materials safely. These measures will help to protect the environment.

## 6. Contents of the pack and other information

### What Zinforo contains

- Each vial contains 600 mg of ceftaroline fosamil.
- The other ingredient is arginine.

### What Zinforo looks like and contents of the pack

Zinforo is a pale yellowish-white to light yellow powder for concentrate for solution for infusion in a vial. It is available in packs containing 10 vials.

### Marketing Authorisation Holder

Pfizer Ireland Pharmaceuticals  
Operations Support Group  
Ringaskiddy, County Cork  
Ireland

### Manufacturer

ACS Dobfar S.p.A.  
Nucleo Industriale S. Atto  
64100 Teramo  
Italy

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

### België/Belgique/Belgien

Pfizer S.A./N.V.  
Tél/Tel: +32 (0)2 554 62 11

### Lietuva

Pfizer Luxembourg SARL filialas Lietuvoje  
Tel. +3705 2514000

### България

Пфайзер Люксембург САРЛ, Клон България  
Тел.: +359 2 970 4333

### Luxembourg/Luxemburg

Pfizer S.A.  
Tél/Tel: +32 (0)2 554 62 11

### Česká republika

Pfizer spol s r.o.  
Tel: +420-283-004-111

### Magyarország

PFIZER Kft.  
Tel. + 36 1 488 37 00

**Danmark**

Pfizer ApS  
Tlf: +45 44 20 11 00

**Deutschland**

Pfizer Pharma PFE GmbH  
Tel: +49 (0)800 8535555

**Eesti**

Pfizer Luxembourg SARL Eesti filiaal  
Tel: +372 666 7500

**Ελλάδα**

Pfizer ΕΛΛΑΣ Α.Ε.  
Τηλ.: +30 210 67 85 800

**España**

Pfizer, S.L.  
Tel: +34 91 490 99 00

**France**

Pfizer  
Tél: +33 (0)1 58 07 34 40

**Hrvatska**

Pfizer Croatia d.o.o.  
Tel: + 385 1 3908 777

**Ireland**

Pfizer Healthcare Ireland  
Tel: +1800 633 363 (toll free)  
+44 (0)1304 616161

**Ísland**

Icepharma hf.  
Sími: +354 540 8000

**Italia**

Pfizer Italia S.r.l.  
Tel: +39 06 33 18 21

**Κύπρος**

Pfizer ΕΛΛΑΣ Α.Ε. (CYPRUS BRANCH)  
Τηλ: +357 22 817690

**Latvija**

Pfizer Luxembourg SARL filiāle Latvijā  
Tel: +371 670 35 775

**Malta**

Vivian Corporation Ltd.  
Tel: +35621 344610

**Nederland**

Pfizer bv  
Tel: +31 (0)10 406 43 01

**Norge**

Pfizer AS  
Tlf: +47 67 52 61 00

**Österreich**

Pfizer Corporation Austria Ges.m.b.H.  
Tel: +43 (0)1 521 15-0

**Polska**

Pfizer Polska Sp. z o.o.  
Tel.: +48 22 335 61 00

**Portugal**

Laboratórios Pfizer, Lda.  
Tel: +351 21 423 5500

**România**

Pfizer România S.R.L.  
Tel: +40 (0)21 207 28 00

**Slovenija**

Pfizer Luxembourg SARL  
Pfizer, podružnica za svetovanje s področja  
farmacevtske dejavnosti, Ljubljana  
Tel: + 386 (0)1 52 11 400

**Slovenská republika**

Pfizer Luxembourg SARL, organizačná zložka  
Tel: +421-2-3355 5500

**Suomi/Finland**

Pfizer PFE Finland Oy  
Puh./Tel: +358 (0)9 43 00 40

**Sverige**

Pfizer AB  
Tel: +46 (0)8 550 520 00

**United Kingdom**

Pfizer Limited  
Tel: +44 (0) 1304 616161

**This leaflet was last revised in**

## Other sources of information

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>.

-----  
The following information is intended for medical or healthcare professionals only:

Important: Please refer to the Summary of Product Characteristics before prescribing.

Aseptic technique must be followed in preparing the infusion solution. The contents of Zinfo vial should be reconstituted with 20 mL of sterile water for injections. Instructions for the reconstitution of Zinfo vial are summarized below:

<b>Dosage strength (mg)</b>	<b>Volume of diluent to be added (mL)</b>	<b>Approximate ceftaroline concentration (mg/mL)</b>	<b>Amount to be withdrawn</b>
600	20	30	Total volume

The reconstituted solution must be further diluted to produce Zinfo solution for infusion. A 250 mL, 100 mL or 50 mL infusion bag can be used to prepare the infusion, based on the patient's volume requirements. Appropriate infusion diluents include: sodium chloride 9 mg/mL (0.9%) solution for injection, dextrose 50 mg/mL (5%) solution for injection, sodium chloride 4.5 mg/mL and dextrose 25 mg/mL solution for injection (0.45% sodium chloride and 2.5% dextrose) or Lactated Ringer's solution. The resulting solution should be administered according to the dose selected over 5 to 60 minutes for standard dose or 120 minutes for high dose in infusion volumes of 50 mL, 100 mL or 250 mL.

Infusion volumes for paediatric patients will vary according to the weight of the child. The infusion solution concentration during preparation and administration should not exceed 12 mg/mL ceftaroline fosamil.

Reconstitution time is less than 2 minutes. Mix gently to reconstitute and check to see that the contents have dissolved completely. Parenteral drug products should be inspected visually for particulate matter prior to administration.

The colour of Zinfo infusion solutions ranges from clear, light to dark yellow depending on the concentration and storage conditions. It is free of any particles. When stored as recommended, the product potency is not affected.

Studies have shown that Zinfo solutions for infusion are stable for up to 6 hours at room temperature. Alternatively they are stable for up to 24 hours under refrigerated storage. Once removed from refrigeration to room temperature, the diluted product must be used within 6 hours.

From a microbiological point of view, the medicinal product should be used immediately unless reconstitution and dilution has taken place in controlled and validated aseptic conditions. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user.

The compatibility of Zinfo with other medicines has not been established. Zinfo should not be mixed with or physically added to solutions containing other drugs.

Each vial is for single use only.

Any unused product or waste material should be disposed of in accordance with local requirements.