

**ANNEX I**

**LIST OF THE NAMES, PHARMACEUTICAL FORM, STRENGTHS OF THE MEDICINAL  
PRODUCTS, ROUTE OF ADMINISTRATION, MARKETING AUTHORISATION  
HOLDERS IN THE MEMBER STATES**

<b>Member State</b>	<b>Marketing Authorisation Holder</b>	<b>Product Name</b>	<b>Strength</b>	<b>Pharmaceutical Form</b>	<b>Route of administration</b>
Austria	Glaxo SmithKline Pharma GmbH, Albert-Schweitzer-Gasse 6 A-1140 Wien Austria	Seretide forte Dosieraerosol	25/250	Pressurised inhalation, suspension	Inhalation use
Austria	Glaxo SmithKline Pharma GmbH, Albert-Schweitzer-Gasse 6 A-1140 Wien Austria	Seretide standard Dosieraerosol	25/125	Pressurised inhalation, suspension	Inhalation use
Austria	Glaxo SmithKline Pharma GmbH Albert Schweitzer Gasse 6 1140 Wien Austria	Seretide junior Dosieraerosol	25/50	Pressurised inhalation, suspension	Inhalation use
Belgium	GlaxoSmithKline s.a./n.v. Rue du Tilleul 13 B-1332 Genval Belgium	Seretide 25/125	25/125	Pressurised inhalation, suspension	Inhalation use
Belgium	GlaxoSmithKline s.a./n.v., Rue du Tilleul 13 B-1332 Genval Belgium	Seretide 25/250	25/250	Pressurised inhalation, suspension	Inhalation use
Belgium	GlaxoSmithKline s.a./n.v. Rue du Tilleul 13 B-1332 Genval Belgium	Seretide 25/50	25/50	Pressurised inhalation, suspension	Inhalation use
Denmark	GlaxoSmithKline Pharma A/S Nykær 68 DK-2605 Brøndby Denmark	Seretide	25/50	Pressurised inhalation, suspension	Inhalation use
Denmark	GlaxoSmithKline Pharma A/S Nykær 68 DK-2605 Brøndby Denmark	Seretide	25/125	Pressurised inhalation, suspension	Inhalation use
Denmark	GlaxoSmithKline Pharma A/S, Nykær 68 DK-2605 Brøndby Denmark	Seretide	25/250	Pressurised inhalation, suspension	Inhalation use

Finland	GlaxoSmithKline Oy Piispansilta 9A, P.O. Box 24 FIN-02231, Espoo Finland	Seretide Evohaler	25/125	Pressurised inhalation, suspension	Inhalation use
Finland	GlaxoSmithKline Oy Piispansilta 9A, P.O. Box 24 FIN-02231, Espoo Finland	Seretide Evohaler	25/250	Pressurised inhalation, suspension	Inhalation use
Finland	GlaxoSmithKline Oy Piispansilta 9A, P.O. Box 24 FIN-02231, Espoo Finland	Seretide Evohaler	25/50	Pressurised inhalation, suspension	Inhalation use
France	Laboratoire GlaxoSmithKline 100 route de Versailles 78163 Marly-le-roi Cedex France	Seretide	25/50	Pressurised inhalation suspension	Inhalation use
France	Laboratoire GlaxoSmithKline 100 route de Versailles 78163 Marly-le-roi Cedex France	Seretide	25/125	Pressurised inhalation suspension	Inhalation use
France	Laboratoire GlaxoSmithKline 100 route de Versailles 78163 Marly-le-roi Cedex France	Seretide	25/250	Pressurised inhalation suspension	Inhalation use
Germany	Schwarz Pharma Deutschland GmbH Alfred-Nobel-Str. 10 D-40789 Monheim Germany	atmadisc mite Dossier- Aerosol FCKW-frei 25 µg/50 µg Druckgasinhalation, Suspension	25/50	Pressurised inhalation suspension	Inhalation use
Germany	Schwarz Pharma Deutschland GmbH Alfred-Nobel-Str. 10 D-40789 Monheim Germany	atmadisc Dossier-Aerosol FCKW-frei 25 µg/125 µg Druckgasinhalation, Suspension	25/125	Pressurised inhalation suspension	Inhalation use

Germany	Schwarz Pharma Deutschland GmbH Alfred-Nobel-Str. 10 D-40789 Monheim Germany	atmadisc forte Dosier-Aerosol FCKW-frei 25 µg/250 µg Druckgasinhalation, Suspension	25/250	Pressurised inhalation suspension	Inhalation use
Greece	GlaxoSmithKline A.E.B.E Leof. Kifisias 266 15232 Halandri, Athens Greece	Seretide Inhaler	25/250	Pressurised inhalation suspension	Inhalation use
Greece	GlaxoSmithKline A.E.B.E Leof. Kifisias 266 15232 Halandri, Athens Greece	Seretide Inhaler	25/50	Pressurised inhalation suspension	Inhalation use
Greece	GlaxoSmithKline A.E.B.E Leof. Kifisias 266 15232 Halandri, Athens Greece	Seretide Inhaler	25/125	Pressurised inhalation suspension	Inhalation use
Iceland	GlaxoSmithKline ehf Thverholt 14, 105 Reykjavik Iceland	Seretide	25/50	Pressurised inhalation suspension	Inhalation use
Iceland	GlaxoSmithKline ehf Thverholt 14, 105 Reykjavik Iceland	Seretide	25/125	Pressurised inhalation suspension	Inhalation use
Iceland	GlaxoSmithKline ehf Thverholt 14, 105 Reykjavik Iceland	Seretide	25/250	Pressurised inhalation suspension	Inhalation use
Ireland	GlaxoSmithKline (Ireland) Limited Stonemasons Way Rathfarnham, Dublin 16 Ireland	Seretide 50 Evohaler	25/50	Pressurised inhalation, suspension.	Inhalation use
Ireland	GlaxoSmithKline (Ireland) Limited Stonemasons Way Rathfarnham, Dublin 16 Ireland	Seretide 125 Evohaler	25/125	Pressurised inhalation, suspension.	Inhalation use

Ireland	GlaxoSmithKline (Ireland) Limited Stonemasons Way Rathfarnham, Dublin 16 Ireland	Seretide 250 Evohaler	25/250	Pressurised inhalation, suspension.	Inhalation use
Italy	GlaxoSmithKline S.p.A. Via A. Fleming, 2 37135 Verona Italy	Seretide	25/50	Pressurised inhalation, suspension	Inhalation use
Italy	GlaxoSmithKline S.p.A. Via A. Fleming, 2 37135 Verona Italy	Seretide	25/125	Pressurised inhalation, suspension	Inhalation use
Italy	GlaxoSmithKline S.p.A. Via A. Fleming, 2 37135 Verona Italy	Seretide	25/250	Pressurised inhalation, suspension	Inhalation use
Luxembourg	GlaxoSmithKline s.a./n.v. Rue du Tilleul 13 B-1332 Genval Belgium	Seretide 25/50	25/50	Pressurised inhalation, suspension	Inhalation use
Luxembourg	GlaxoSmithKline s.a./n.v. Rue du Tilleul 13 B-1332 Genval Belgium	Seretide 25/125	25/125	Pressurised inhalation, suspension	Inhalation use
Luxembourg	GlaxoSmithKline s.a./n.v. Rue du Tilleul 13 B-1332 Genval Belgium	Seretide 25/250	25/250	Pressurised inhalation, suspension	Inhalation use
Portugal	Glaxo Wellcome Farmacêutica, Lda. Rua Dr. António Loureiro Borges, 3, Arquiparque , Miraflores 1495- 131 Algés Portugal	Seretaide Inalador	25/50	Pressurised inhalation, suspension	Inhalation use

Portugal	Glaxo Wellcome Farmacêutica, Lda. Rua Dr. António Loureiro Borges, 3, Arquiparque , Miraflores 1495- 131 Algés Portugal	Seretaide Inalador	25/125	Pressurised inhalation, suspension	Inhalation use
Portugal	Glaxo Wellcome Farmacêutica, Lda. Rua Dr. António Loureiro Borges, 3, Arquiparque , Miraflores 1495- 131 Algés Portugal	Seretaide Inalador	25/250	Pressurised inhalation, suspension	Inhalation use
Spain	GlaxoSmithKline SA Parque Tecnológico de Madrid, Calle Severo Ochoa 2, 28760, Tres Cantos, Madrid Spain	Seretide 25/50	25/50	Pressurised inhalation suspension	Inhalation use
Spain	GlaxoSmithKline SA Parque Tecnológico de Madrid, Calle Severo Ochoa 2, 28760 Tres Cantos, Madrid Spain	Seretide 25/125	25/125	Pressurised inhalation suspension	Inhalation use
Spain	GlaxoSmithKline SA Parque Tecnológico de Madrid, Calle Severo Ochoa 2, 28760 Tres Cantos, Madrid Spain	Seretide 25/250	25/250	Pressurised inhalation suspension	Inhalation use
Sweden	GlaxoSmithKline AB Box 516, SE-169 29 Solna Sweden	Seretide Evohaler mite	25/50	Pressurised inhalation, suspension	Inhalation use
Sweden	GlaxoSmithKline AB Box 516, SE-169 29 Solna Sweden	Seretide Evohaler	25/125	Pressurised inhalation, suspension	Inhalation use
Sweden	GlaxoSmithKline AB Box 516, SE-169 29 Solna Sweden	Seretide Evohaler forte	25/250	Pressurised inhalation, suspension	Inhalation use

The Netherlands	GlaxoSmithKline B.V. Huis ter Heideweg 62 3705 LZ Zeist The Netherlands	Seretide 25/50	25/50	Pressurised inhalation, suspension	Inhalation use
The Netherlands	GlaxoSmithKline B.V. Huis ter Heideweg 62 3705 LZ Zeist The Netherlands	Seretide 25/125	25/125	Pressurised inhalation, suspension	Inhalation use
The Netherlands	GlaxoSmithKline B.V. Huis ter Heideweg 62 3705 LZ Zeist The Netherlands	Seretide 25/250	25/250	Pressurised inhalation, suspension	Inhalation use
United Kingdom	Glaxo Wellcome UK Limited, Stockley Park West, Uxbridge Middlesex UB11 1BT United Kingdom	Seretide 50 Evohaler	25/50	Pressurised inhalation, suspension	Inhalation use
United Kingdom	Glaxo Wellcome UK Limited, Stockley Park West, Uxbridge Middlesex UB11 1BT United Kingdom	Seretide 125 Evohaler	25/125	Pressurised inhalation, suspension	Inhalation use
United Kingdom	Glaxo Wellcome UK Limited, Stockley Park West, Uxbridge Middlesex UB11 1BT United Kingdom	Seretide 250 Evohaler	25/250	Pressurised inhalation, suspension	Inhalation use

**ANNEX II**

**SCIENTIFIC CONCLUSIONS AND GROUNDS FOR AMENDMENT OF THE SUMMARY  
OF PRODUCT CHARACTERISTICS, LABELLING AND PACKAGE LEAFLET  
PRESENTED BY THE EMEA**



## SCIENTIFIC CONCLUSIONS

### OVERALL SUMMARY OF THE SCIENTIFIC EVALUATION OF SERETIDE EVOHALER AND ASSOCIATED NAMES (SEE ANNEX I)

Seretide Evohaler and associated names (see Annex I of Opinion) contain salmeterol and fluticasone propionate, a fixed dose combination of the long acting  $\beta$ -agonist (LABA) salmeterol and the inhaled corticosteroid (ICS) fluticasone propionate (FP) indicated in the regular treatment of asthma where use of a combination product (long-acting beta-2-agonist and inhaled corticosteroid) is appropriate: - patients not adequately controlled with inhaled corticosteroids and 'as needed' inhaled short acting beta-2-agonist or - patients already adequately controlled on both inhaled corticosteroid and long-acting beta-2-agonist.

In several EU Member States Seretide Evohaler and associated names (see Annex I) are approved through the Mutual Recognition Procedure (MRP). In the MRP the United Kingdom is the Reference Member State and Austria, Belgium, Denmark, Finland, France, Greece, Germany, Italy, Ireland, Iceland, Luxembourg, The Netherlands, Portugal, Spain and Sweden are Concerned Member States.

The MAHs applied for a type II variation subject to MRP to extend the current approved indications to include initial maintenance therapy (IMT) with the fixed dose combination in patients with chronic persistent asthma. The Reference and Concerned Member States rejected the variation to extend the indication on 3 August 2005. The main objection relating to the identification of the patient population especially those patients who would respond to inhaled corticosteroids alone had not been adequately identified.

The MAHs were of the opinion that the issues identified during the MRP were issues of clinical management and as such cannot be addressed by conducting further clinical trials. On 12 August 2005, the MAHs presented to the EMEA a referral under Article 6(13) of Commission Regulation (EC) No 1084/2003.

The issue to be considered by CHMP was whether it is appropriate to introduce initial maintenance therapy with the fixed dose combination of salmeterol and fluticasone propionate in all patients with chronic persistent asthma.

The CHMP performed a re-evaluation of the type II variation application data. A discussion on the appropriateness of initial maintenance therapy with the fixed dose combination took place in CHMP based on the Rapporteur's and Co-Rapporteur's Assessment Reports and the data presented by the Marketing Authorisation Holders (MAHs).

## EFFICACY

The MAHs have demonstrated in a series of substantial and well conducted clinical studies that in patients with moderate chronic persistent asthma poorly controlled on a short acting  $\beta$ -agonist alone the introduction of combination therapy with salmeterol and fluticasone is more effective in improving disease control than the introduction of salmeterol or fluticasone alone.

A subpopulation from the GOAL study (Gaining Optimal Asthma Control study) i.e. stratum 1, addendum, step 1 (patients without ICS at entry who fulfilled the criteria for moderate persistent asthma and who are randomised to Seretide 50/100 or FP 100 mcg), can be considered as an adequate target population to assess the adequacy of Seretide as IMT. Patients with mild persistent asthma are not considered to be candidates for a combination therapy as IMT and patients with a more severe disease would deserve more intensive therapeutic approaches.

The primary endpoint in the GOAL study was the proportion of patients reaching asthma control (well-control of asthma). The CHMP considered the use of "Control of Asthma" as the primary efficacy variable a reliable parameter of the patient clinical status and, as such, clinically meaningful.

The CHMP had a concern that a general recommendation i.e. the therapeutic indication in the Summary of Product Characteristics (SPC), will result in over treatment particularly in patients who would have responded to inhaled corticosteroids alone. The MAH's proposed therapeutic indication to include "*patients with persistent asthma not adequately controlled on "as needed" inhaled short-acting beta-2-agonists alone who exhibit a combination of at least two of the following clinical features of asthma: airflow limitation, daily use of rescue medication, daily symptoms (day-time and/or night-time)*" was not considered acceptable as it does not effectively identify a population in need of combination therapy.

Nevertheless the CHMP acknowledged that current clinical guidelines (e.g. the Global Initiative for Asthma (GINA) guidelines) describe circumstances which could lead to the initiation of maintenance therapy with the combination. Therefore the CHMP considered it acceptable to give guidance to physicians as to the most appropriate dose strategy for initiating maintenance therapy with the fixed dose combination. A recommendation in section 4.2 "Posology and method of administration" was considered acceptable.

The recommendation included in section 4.2 allows a short term trial of Seretide Evohaler in adults and adolescents with moderate persistent asthma for whom rapid control of asthma is essential. The initial starting dose is 2 inhalations of 25 mcg salmeterol and 50 mcg FP twice daily. The wording also indicates the importance for stepping down to inhaled corticosteroids alone once asthma control is attained. It is also highlighted that Seretide is not appropriate as IMT in patients with mild and severe asthma and that in general inhaled corticosteroids remain the first line treatment for most patients.

In addition to the update in section 4.2 of the SPC, some factual data from the GOAL study in particular the timing of reaching more rapidly control with Seretide than with ICS alone have been included in section 5.1 "Pharmacodynamic properties" of the SPC. The information currently in section 5.1 of the SPC have also been amended in line with the recommendation in section 4.2.

## **SAFETY**

Data from six pivotal studies and two supporting studies have been presented to confirm the safety profile of salmeterol/FP when initiated as maintenance treatment. Although the safety profile of salmeterol/fluticasone propionate is well known, the CHMP considered it not justified to expose the whole intended target population to an increased incidence of beta-agonists related adverse events. In line with above arguments, the CHMP was not in agreement with a general recommendation for IMT of the fixed dose combination in section 4.1. but considered it acceptable to include guidance for a short term trial of initial maintenance therapy with Seretide in section 4.2.

## **GROUNDINGS FOR AMENDMENT OF THE SUMMARY OF PRODUCT CHARACTERISTICS, LABELLING AND PACKAGE LEAFLET**

Whereas

- the CHMP considered the referral made under Article 6(13) of Commission Regulation (EC) N 1084/2003 for Seretide Evohaler and associated names (see Annex I)
- the CHMP considered that an initial maintenance therapy indication with the fixed dose combination of salmeterol and fluticasone propionate in patients with moderate persistent asthma in section 4.1 of the SPC is not acceptable as it could result in overtreatment particularly in patients who would have responded to inhaled corticosteroids alone. CHMP considered it not acceptable to expose the whole target population to beta-agonists related adverse events.
- the CHMP agreed, however, that a recommendation for a short term trial with the fixed dose combination of salmeterol and fluticasone propionate as initial maintenance therapy in adults

and adolescents with moderate persistent asthma for whom rapid control of asthma is essential, in section 4.2 of the SPC is acceptable.

- the CHMP agreed that factual data from the GOAL study in particular the timing of reaching more rapidly control with Seretide than with ICS alone should be included in section 5.1 of the SPC and that some amendments in this section were necessary in line with the agreed recommendation in section 4.2.

The CHMP has recommended the granting of the variation of the Marketing Authorisations for which the Summary of Product Characteristics, labelling and package leaflet are set out in Annex III.

### **ANNEX III**

**Note: This Annex III (Summary of Product Characteristics, Labelling and Package Leaflet) is the one that was Annexed to the Commission Decision on this Article 6(13) referral for Seretide Evohaler and associated names. The text was valid at that time.**

**After the Commission Decision, the Member State competent authorities will update Annex III as required. Therefore, Annex III may not necessarily represent the current text.**

## **SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE MEDICINAL PRODUCT

Seretide Evohaler and associated names (see Annex I) 25/50 microgram/dose pressurised inhalation, suspension.

Seretide Evohaler and associated names (see Annex I) 25/125 microgram/dose pressurised inhalation, suspension.

Seretide Evohaler and associated names (see Annex I) 25/250 microgram/dose pressurised inhalation, suspension.

[See Annex I - To be completed nationally]

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each single actuation of Seretide provides:

25 micrograms of salmeterol (as salmeterol xinafoate) and 50, 125 or 250 micrograms of fluticasone propionate (delivered from the valve). This is equivalent to 21 micrograms of salmeterol and 44, 110 or 220 micrograms of fluticasone propionate delivered from the actuator (delivered dose).

For a full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Pressurised inhalation, suspension.

The canister contains a white to off white suspension.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Seretide is indicated in the regular treatment of asthma where use of a combination product (long-acting beta-2-agonist and inhaled corticosteroid) is appropriate:

- patients not adequately controlled with inhaled corticosteroids and ‘as needed’ inhaled short acting beta-2-agonist
- or
- patients already adequately controlled on both inhaled corticosteroid and long-acting beta-2-agonist.

### 4.2 Posology and method of administration

Seretide Evohaler is for inhalation use only.

Patients should be made aware that Seretide Evohaler must be used daily for optimum benefit, even when asymptomatic.

Patients should be regularly reassessed by a doctor, so that the strength of Seretide they are receiving remains optimal and is only changed on medical advice. **The dose should be titrated to the lowest dose at which effective control of symptoms is maintained. Where the control of symptoms is**

**maintained with the lowest strength of the combination given twice daily then the next step could include a test of inhaled corticosteroid alone.** As an alternative, patients requiring a long acting beta-2-agonist could be titrated to Seretide given once daily if, in the opinion of the prescriber, it would be adequate to maintain disease control. In the event of once daily dosing when the patient has a history of nocturnal symptoms the dose should be given at night and when the patient has a history of mainly day-time symptoms the dose should be given in the morning.

Patients should be given the strength of Seretide containing the appropriate fluticasone propionate dosage for the severity of their disease. Note: Seretide 25/50 microgram strength is not appropriate in adults and children with severe asthma. Prescribers should be aware that, in patients with asthma, fluticasone propionate is as effective as other inhaled steroids at approximately half the microgram daily dose. For example, 100mcg of fluticasone propionate is approximately equivalent to 200mcg of beclomethasone dipropionate (CFC containing) or budesonide. If an individual patient should require dosages outside the recommended regimen, appropriate doses of beta-agonist and/or corticosteroid should be prescribed.

#### Recommended Doses:

Adults and adolescents 12 years and older:

Two inhalations of 25 micrograms salmeterol and 50 micrograms fluticasone propionate twice daily.

or

Two inhalations of 25 micrograms salmeterol and 125 micrograms fluticasone propionate twice daily.

or

Two inhalations of 25 micrograms salmeterol and 250 micrograms fluticasone propionate twice daily.

A short term trial of Seretide may be considered as initial maintenance therapy in adults or adolescents with moderate persistent asthma (defined as patients with daily symptoms, daily rescue use and moderate to severe airflow limitation) for whom rapid control of asthma is essential. In these cases, the recommended initial dose is two inhalations of 25 micrograms salmeterol and 50 micrograms fluticasone propionate twice daily. Once control of asthma is attained treatment should be reviewed and consideration given as to whether patients should be stepped down to an inhaled corticosteroid alone.

A clear benefit has not been shown as compared to inhaled fluticasone propionate alone used as initial maintenance therapy when one or two of the criteria of severity are missing. In general inhaled corticosteroids remain the first line treatment for most patients. Seretide is not intended for the initial management of mild asthma. Seretide 25/50 micrograms strength is not appropriate in adults and children with severe asthma; it is recommended to establish the appropriate dosage of inhaled corticosteroid before any fixed combination can be used in patients with severe asthma.

Children 4 years and older:

Two inhalations of 25 micrograms salmeterol and 50 micrograms fluticasone propionate twice daily.

The maximum licensed dose of fluticasone propionate delivered by Seretide inhaler in children is 100mcg twice daily.

There are no data available for use of Seretide inhaler in children aged under 4 years.

The use of a spacer device with the inhaler in order to derive greater therapeutic benefit is recommended for patients (in particular young children) who have difficulty co-ordinating inhalation with actuation (see section 4.4 Special Warnings and Precautions for use)



Special patient groups:

There is no need to adjust the dose in elderly patients or in those with renal impairment. There are no data available for use of Seretide in patients with hepatic impairment.

#### Instructions for Use:

Patients should be instructed in the proper use of their inhaler (see patient information leaflet) During inhalation, the patient should preferably sit or stand. The inhaler has been designed for use in a vertical position.

Testing the inhaler:

Before using for the first time patients should remove the mouthpiece cover by gently squeezing the sides of the cover, shake the inhaler well, and release puffs into the air until the counter reads 120 to make sure that it works. If the inhaler has not been used for a week or more remove the mouthpiece cover, the patients should shake the inhaler well and release two puffs into the air. Each time the inhaler is activated the number on the counter will count down by one.

Use of the inhaler:

1. Patients should remove the mouthpiece cover by gently squeezing the sides of the cover and check the mouthpiece inside and outside to see that it is clean.
2. Patients should shake the inhaler well.
3. Patients should hold the inhaler upright between fingers and thumb with their thumb on the base, below the mouthpiece.
4. Patients should breathe out as far as is comfortable and then place the mouthpiece in your mouth between their teeth and close their lips around it, Patients should be instructed not to bite the mouth piece.
5. Just after starting to breathe in through their mouth, patients should press down on the top of the inhaler to release Seretide, while still breathing in steadily and deeply.
6. While holding their breath, patients should take the inhaler from their mouth and take their finger from the top of the inhaler. Patients should continue holding their breath for as long as is comfortable.
7. If patients are going to take a second puff they should keep the inhaler upright and wait about half a minute before repeating steps 2 to 6.
8. After use patients should always replace the mouthpiece cover by firmly pushing and snapping the cap into position.

#### IMPORTANT

Patients should not rush stages 4, 5 and 6. It is important that patients start to breathe in as slowly as possible just before operating their inhaler. Patients should practise in front of a mirror for the first few times. If they see "mist" coming from the top of their inhaler or the sides of their mouth they should start again from stage 2.

Patients should consider getting a replacement when the counter shows the number 020. Replace the inhaler when the counter reads 000.

Patients should never try to alter the numbers on the counter or detach the counter from the metal canister. The counter cannot be reset and is permanently attached to the canister.

Cleaning:

1. Remove the mouth piece cover.
2. Do not remove the canister from the plastic casing.
3. Wipe the inside and outside of the mouthpiece and the plastic casing with a dry cloth or tissue.
4. Replace the mouthpiece cover.

DO NOT PUT THE METAL CONTAINER IN WATER

### **4.3 Contraindications**

Seretide is contraindicated in patients with hypersensitivity to any of the active substances or to the excipient.

### **4.4 Special warnings and precautions for use**

The management of asthma should normally follow a stepwise programme and patient response should be monitored clinically and by lung function tests.

Seretide Evohaler should not be used to treat acute asthma symptoms for which a fast and short acting bronchodilator is required. Patients should be advised to have their medicinal product to be used for relief in an acute asthma attack available at all times.

Increasing use of short-acting bronchodilators to relieve asthma symptoms indicates deterioration of asthma control and patients should be reviewed by a physician.

Sudden and progressive deterioration in control of asthma is potentially life-threatening and the patient should undergo urgent medical assessment. Consideration should be given to increasing corticosteroid therapy. The patient should also be medically reviewed where the current dosage of Seretide has failed to give adequate control of asthma. Consideration should be given to additional corticosteroid therapies.

Treatment with Seretide should not be stopped abruptly.

As with all inhaled medication containing corticosteroids, Seretide should be administered with caution in patients with pulmonary tuberculosis.

Seretide should be administered with caution in patients with severe cardiovascular disorders, including heart rhythm abnormalities, diabetes mellitus, untreated hypokalaemia or thyrotoxicosis.

There have been very rare reports of increases in blood glucose levels (see 4.8 Undesirable Effects) and this should be considered when prescribing to patients with a history of diabetes mellitus.

Potentially serious hypokalaemia may result from systemic beta-2-agonist therapy, but following inhalation at therapeutic doses plasma levels of salmeterol are very low.

As with other inhalation therapy paradoxical bronchospasm may occur with an immediate increase in wheezing after dosing. Seretide Evohaler should be discontinued immediately, the patient assessed and alternative therapy instituted if necessary.

Care should be taken when transferring patients to Seretide therapy, particularly if there is any reason to suppose that adrenal function is impaired from previous systemic steroid therapy.

Systemic effects may occur with any inhaled corticosteroid, particularly at high doses prescribed for long periods. These effects are much less likely to occur than with oral corticosteroids. Possible

systemic effects include Cushing's syndrome, Cushingoid features, adrenal suppression, growth retardation in children and adolescents, decrease in bone mineral density, cataract and glaucoma.

**It is important, therefore, that the patient is reviewed regularly and the dose of inhaled corticosteroid is reduced to the lowest dose at which effective control of asthma is maintained.**

It is recommended that the height of children receiving prolonged treatment with inhaled corticosteroid is regularly monitored.

Prolonged treatment of patients with high doses of inhaled corticosteroids may result in adrenal suppression and acute adrenal crisis. Children and adolescents <16years taking high doses of fluticasone (typically  $\geq 1000\text{mcg/day}$ ) may be at particular risk. Very rare cases of adrenal suppression and acute adrenal crisis have also been described with doses of fluticasone propionate between 500 and less than 1000mcg. Situations, which could potentially trigger acute adrenal crisis, include trauma, surgery, infection or any rapid reduction in dosage. Presenting symptoms are typically vague and may include anorexia, abdominal pain, weight loss, tiredness, headache, nausea, vomiting, hypotension, decreased level of consciousness, hypoglycaemia, and seizures. Additional systemic corticosteroid cover should be considered during periods of stress or elective surgery.

As systemic absorption is largely through the lungs, the use of a spacer plus metered dose inhaler may increase drug delivery to the lungs. It should be noted that this could potentially lead to an increase in the risk of systemic adverse effects.

The benefits of inhaled fluticasone propionate therapy should minimise the need for oral steroids, but patients transferring from oral steroids may remain at risk of impaired adrenal reserve for a considerable time. Patients who have required high dose emergency corticosteroid therapy in the past may also be at risk. This possibility of residual impairment should always be borne in mind in emergency and elective situations likely to produce stress, and appropriate corticosteroid treatment must be considered. The extent of the adrenal impairment may require specialist advice before elective procedures.

Ritonavir can greatly increase the concentration of fluticasone propionate in plasma. Therefore, concomitant use should be avoided, unless the potential benefit to the patient outweighs the risk of systemic corticosteroid side-effects. There is also an increased risk of systemic side effects when combining fluticasone propionate with other potent CYP3A inhibitors (see 4.5 Interaction with Other Medicinal Products and Other Forms of Interaction).

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Both non-selective and selective beta-blockers should be avoided in patients with asthma, unless there are compelling reasons for their use.

Concomitant use of other beta-adrenergic containing drugs can have a potentially additive effect.

Under normal circumstances, low plasma concentrations of fluticasone propionate are achieved after inhaled dosing, due to extensive first pass metabolism and high systemic clearance mediated by cytochrome P450 3A4 in the gut and liver. Hence, clinically significant drug interactions mediated by fluticasone propionate are unlikely.

In an interaction study in healthy subjects with intranasal fluticasone propionate, ritonavir (a highly potent cytochrome P450 3A4 inhibitor) 100 mg b.i.d. increased the fluticasone propionate plasma concentrations several hundred fold, resulting in markedly reduced serum cortisol concentrations. Information about this interaction is lacking for inhaled fluticasone propionate, but a marked increase in fluticasone propionate plasma levels is expected. Cases of Cushing's syndrome and adrenal suppression have been reported. The combination should be avoided unless the benefit outweighs the increased risk of systemic glucocorticoid side-effects.

In a small study in healthy volunteers, the slightly less potent CYP3A inhibitor ketoconazole increased the exposure of fluticasone propionate after a single inhalation by 150%. This resulted in a greater reduction of plasma cortisol as compared with fluticasone propionate alone. Co-treatment with other potent CYP3A inhibitors, such as itraconazole, is also expected to increase the systemic fluticasone propionate exposure and the risk of systemic side-effects. Caution is recommended and long-term treatment with such drugs should if possible be avoided.

#### **4.6 Pregnancy and lactation**

There are insufficient data on the use of salmeterol and fluticasone propionate during pregnancy and lactation in man to assess the possible harmful effects. In animal studies foetal abnormalities occur after administration of beta-2-adrenoreceptor agonists and glucocorticosteroids (see 5.3 Preclinical Safety Data).

Administration of Seretide to pregnant women should only be considered if the expected benefit to the mother is greater than any possible risk to the foetus.

The lowest effective dose of fluticasone propionate needed to maintain adequate asthma control should be used in the treatment of pregnant women.

There are no data available for human breast milk. Both salmeterol and fluticasone propionate are excreted into breast milk in rats. Administration of Seretide to women who are breastfeeding should only be considered if the expected benefit to the mother is greater than any possible risk to the child.

#### **4.7 Effects on ability to drive and use machines**

No studies of the effect on the ability to drive and use machines have been performed.

#### **4.8 Undesirable effects**

As Seretide contains salmeterol and fluticasone propionate, the type and severity of adverse reactions associated with each of the compounds may be expected. There is no incidence of additional adverse events following concurrent administration of the two compounds.

Adverse events which have been associated with salmeterol/fluticasone propionate are given below, listed by system organ class and frequency. Frequencies are defined as: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  and  $< 1/10$ ), uncommon ( $\geq 1/1000$  and  $< 1/100$ ), and very rare ( $< 1/10,000$ ) including isolated reports. Very common, common and uncommon events were derived from clinical trial data. The incidence in placebo was not taken into account. Very rare events were derived from post-marketing spontaneous data.

System Organ Class	Adverse Event	Frequency
Infections & Infestations	Candidiasis of the mouth and throat	Common
Immune System Disorders	Hypersensitivity reactions with the following manifestations:  Cutaneous hypersensitivity reactions	Uncommon
	Angioedema (mainly facial and oropharyngeal oedema), Respiratory symptoms (dyspnoea and/or bronchospasm), Anaphylactic reactions	Very Rare
Endocrine Disorders	Cushing's syndrome, Cushingoid features, Adrenal suppression, Growth retardation in children and adolescents, Decreased bone mineral density, Cataract, Glaucoma	Very Rare
Metabolism & Nutrition Disorders	Hyperglycaemia	Very Rare
Psychiatric Disorders	Anxiety, sleep disorders and behavioural changes, including hyperactivity and irritability (predominantly in children)	Very Rare
Nervous System Disorders	Headache	*Very Common
	Tremor	Common
Cardiac Disorders	Palpitations	Common
	Tachycardia	Uncommon
	Cardiac arrhythmias (including atrial fibrillation, supraventricular tachycardia and extrasystoles).	Very Rare
Respiratory, Thoracic & Mediastinal Disorders	Throat irritation	Common
	Hoarseness/dysphonia	Common
	Paradoxical bronchospasm	Very Rare
Musculoskeletal & Connective Tissue Disorders	Muscle cramps	Common
	Arthralgia	Very Rare
	Myalgia	Very Rare

\*Reported commonly in placebo

The pharmacological side effects of beta-2-agonist treatment, such as tremor, palpitations and headache, have been reported, but tend to be transient and reduce with regular therapy.

Due to the fluticasone propionate component, hoarseness and candidiasis (thrush) of the mouth and throat can occur in some patients. Both hoarseness and incidence of candidiasis may be relieved by gargling with water after using the product. Symptomatic candidiasis can be treated with topical anti-fungal therapy whilst still continuing with the Seretide Evohaler.

Possible systemic effects include Cushing's syndrome, Cushingoid features, adrenal suppression, growth retardation in children and adolescents, decrease in bone mineral density, cataract and glaucoma (see 4.4 Special Warnings and Precautions For Use).

There have been very rare reports of hyperglycaemia (see 4.4 Special Warnings and Precautions for Use).

As with other inhalation therapy, paradoxical bronchospasm may occur (see 4.4 Special Warnings and Precautions for Use).

#### **4.9 Overdose**

There are no data available from clinical trials on overdose with Seretide, however data on overdose with both drugs are given below:

The signs and symptoms of salmeterol overdose are tremor, headache and tachycardia. The preferred antidotes are cardioselective beta-blocking agents, which should be used with caution in patients with a history of bronchospasm. If Seretide therapy has to be withdrawn due to overdose of the beta agonist component of the drug, provision of appropriate replacement steroid therapy should be considered. Additionally, hypokalaemia can occur and potassium replacement should be considered.

Acute: Acute inhalation of fluticasone propionate doses in excess of those recommended may lead to temporary suppression of adrenal function. This does not need emergency action as adrenal function is recovered in a few days, as verified by plasma cortisol measurements.

Chronic overdose of inhaled fluticasone propionate: Refer to section 4.4: risk of adrenal suppression. Monitoring of adrenal reserve may be necessary. In cases of fluticasone propionate overdose Seretide therapy may still be continued at a suitable dosage for symptom control.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic Group: Adrenergics and other anti-asthmatics.

ATC Code: R03AK06

#### **Seretide Asthma clinical trials**

A twelve month study (Gaining Optimal Asthma Control, GOAL), in 3416 adult and adolescent patients with persistent asthma, compared the safety and efficacy of Seretide versus inhaled corticosteroid (Fluticasone Propionate) alone to determine whether the goals of asthma management were achievable. Treatment was stepped up every 12 weeks until \*\*Total control was achieved or the highest dose of study drug was reached. GOAL showed more patients treated with Seretide achieved asthma control than patients treated with ICS alone and this control was attained at a lower corticosteroid dose.

Well-Controlled asthma was achieved more rapidly with Seretide than with ICS alone. The time on treatment for 50% of subjects to achieve a first individual Well-Controlled week was 16 days for Seretide compared to 37 days for the ICS group. In the subset of steroid naive asthmatics the time to an individual Well Controlled week was 16 days in the Seretide treatment compared to 23 days following treatment with ICS.

The overall study results showed:

<b>Percentage of Patients Attaining *Well Controlled (WC) and **Totally Controlled (TC) Asthma over 12 months</b>				
<b>Pre-Study Treatment</b>	<b>Salmeterol/FP</b>		<b>FP</b>	
	<b>WC</b>	<b>TC</b>	<b>WC</b>	<b>TC</b>
<b>No ICS (SABA alone)</b>	78%	50%	70%	40%
<b>Low dose ICS ( ≤500mcg BDP or equivalent/day)</b>	75%	44%	60%	28%
<b>Medium dose ICS (&gt;500-1000mcg BDP or equivalent/day)</b>	62%	29%	47%	16%
<b>Pooled results across the 3 treatment levels</b>	71%	41%	59%	28%

\*Well controlled asthma; occasional symptoms or SABA use or less than 80% predicted lung function plus no night-time awakenings, no exacerbations and no side effects enforcing a change in therapy

\*\*Total control of asthma; no symptoms, no SABA use, greater than or equal to 80% predicted lung function, no night-time awakenings, no exacerbations and no side effects enforcing a change in therapy.

The results of this study suggest that Seretide 50/100mcg bd may be considered as initial maintenance therapy in patients with moderate persistent asthma for whom rapid control of asthma is deemed essential (see section 4.2).

#### Mechanism of action:

Seretide contains salmeterol and fluticasone propionate which have differing modes of action.

The respective mechanisms of action of both drugs are discussed below.

#### Salmeterol:

Salmeterol is a selective long-acting (12 hour) beta-2-adrenoceptor agonist with a long side chain which binds to the exo-site of the receptor.

Salmeterol produces a longer duration of bronchodilation, lasting for at least 12 hours, than recommended doses of conventional short-acting beta-2-agonists.

#### Fluticasone propionate:

Fluticasone propionate given by inhalation at recommended doses has a glucocorticoid anti-inflammatory action within the lungs, resulting in reduced symptoms and exacerbations of asthma, with less adverse effects than when corticosteroids are administered systemically.

## **5.2 Pharmacokinetic properties**

When salmeterol and fluticasone propionate were administered in combination by the inhaled route, the pharmacokinetics of each component were similar to those observed when the drugs were administered separately. For pharmacokinetic purposes therefore each component can be considered separately.

#### Salmeterol:

Salmeterol acts locally in the lung therefore plasma levels are not an indication of therapeutic effects. In addition there are only limited data available on the pharmacokinetics of salmeterol because of the technical difficulty of assaying the drug in plasma due to the low plasma concentrations at therapeutic doses (approximately 200 picogram/ml or less) achieved after inhaled dosing.

### Fluticasone propionate:

The absolute bioavailability of inhaled fluticasone propionate in healthy subjects varies between approximately 10-30% of the nominal dose depending on the inhalation device used. In patients with asthma a lesser degree of systemic exposure to inhaled fluticasone propionate has been observed.

Systemic absorption occurs mainly through the lungs and is initially rapid then prolonged. The remainder of the inhaled dose may be swallowed but contributes minimally to systemic exposure due to the low aqueous solubility and pre-systemic metabolism, resulting in oral availability of less than 1%. There is a linear increase in systemic exposure with increasing inhaled dose.

The disposition of fluticasone propionate is characterised by high plasma clearance (1150ml/min), a large volume of distribution at steady-state (approximately 300l) and a terminal half-life of approximately 8 hours.

Plasma protein binding is 91%.

Fluticasone propionate is cleared very rapidly from the systemic circulation. The main pathway is metabolism to an inactive carboxylic acid metabolite, by the cytochrome P450 enzyme CYP3A4. Other unidentified metabolites are also found in the faeces.

The renal clearance of fluticasone propionate is negligible. Less than 5% of the dose is excreted in urine, mainly as metabolites. The main part of the dose is excreted as faeces as metabolites and unchanged drug.

### **5.3 Preclinical safety data**

The only safety concerns for human use derived from animal studies of salmeterol xinafoate and fluticasone propionate given separately were effects associated with exaggerated pharmacological actions.

In animal reproduction studies, glucocorticosteroids have been shown to induce malformations (cleft palate, skeletal malformations). However, these animal experimental results do not seem to be relevant for man given recommended doses. Animal studies with salmeterol xinafoate have shown embryofetal toxicity only at high exposure levels. Following co-administration, increased incidences of transposed umbilical artery and incomplete ossification of occipital bone were found in rats at doses associated with known glucocorticoid-induced abnormalities.

The non-CFC propellant, Norflurane, has been shown to have no toxic effect at very high vapour concentrations, far in excess of those likely to be experienced by patients, in a wide range of animal species exposed daily for periods of two years.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Norflurane (HFA 134a).

### **6.2 Incompatibilities**

Not applicable.



### **6.3 Shelf life**

1 year

### **6.4 Special precautions for storage**

The canister contains a pressurised liquid. Do not store above 25°C. The container should not be punctured, broken or burnt even when apparently empty.

As with most inhaled medicinal products in pressurised containers, the therapeutic effect of this medicinal product may decrease when the container is cold.

### **6.5 Nature and contents of container**

The suspension is contained in an internally lacquered, 8ml aluminium alloy pressurised container sealed with a metering valve. The containers are fitted into plastic actuators incorporating an atomising mouthpiece and fitted with dustcaps. The canister has a counter attached to it, which shows how many actuations of medicine are left. The number will show through a window in the back of the plastic actuator. One pressurised container delivers 120 actuations.

The devices are available in cardboard containers, which hold:

1 x 120 actuations Inhaler

or 3 x 120 actuations Inhaler

or 10 x 120 actuations Inhaler - hospital/pharmacy use only (for dispensing purposes)

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal**

No special requirements

## **7. MARKETING AUTHORISATION HOLDER**

[See Annex I - To be completed nationally]

## **8. MARKETING AUTHORISATION NUMBER(S)**

[To be completed nationally]

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

[To be completed nationally]

## **10. DATE OF REVISION OF THE TEXT**

[To be completed nationally]

## **LABELLING AND PACKAGE LEAFLET**

## **LABELLING**

## **PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

### **OUTER CARTON**

#### **1. NAME OF THE MEDICINAL PRODUCT**

Seretide Evohaler and associated names (see Annex I) 25/50 microgram/dose pressurised inhalation, suspension

[See Annex I – To be completed nationally]

Salmeterol xinafoate and fluticasone propionate

#### **2. STATEMENT OF ACTIVE SUBSTANCE(S)**

25 micrograms of salmeterol (as xinafoate) and 50 micrograms of fluticasone propionate per actuation

#### **3. LIST OF EXCIPIENTS**

Norflurane (HFA 134a)

#### **4. PHARMACEUTICAL FORM AND CONTENTS**

Pressurised inhalation, suspension

1 X 120 puffs

3 X 120 puffs

10 X 120 puffs (hospital or pharmacy use only)

#### **5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Shake well before use

Read the package leaflet before use

Inhalation use

#### **6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

Pressurised container. Do not puncture, break or burn even when apparently empty

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

[See Annex I - To be completed nationally]

**12. MARKETING AUTHORISATION NUMBER(S)**

[To be completed nationally]

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

[To be completed nationally]

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

[To be completed nationally]

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**  
**EVOHALER LABEL**

**1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

Seretide Evohaler and associated names (see Annex I) 25/50 microgram/dose pressurised inhalation, suspension

[See Annex I – To be completed nationally]

Salmeterol xinafoate and fluticasone propionate

Inhalation use

**2. METHOD OF ADMINISTRATION**

Shake well before use

Read the package leaflet before use

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

120 puffs

**6. OTHER**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Seretide Evohaler and associated names (see Annex I) 25/125 microgram/dose pressurised inhalation, suspension

[See Annex I – To be completed nationally]

Salmeterol xinafoate and fluticasone propionate

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

25 micrograms of salmeterol (as xinafoate) and 125 micrograms of fluticasone propionate per actuation

**3. LIST OF EXCIPIENTS**

Norflurane (HFA 134a)

**4. PHARMACEUTICAL FORM AND CONTENTS**

Pressurised inhalation, suspension

1 X 120 puffs

3 X 120 puffs

10 X 120 puffs (hospital or pharmacy use only)

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Shake well before use

Read the package leaflet before use

Inhalation use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

Pressurised container. Do not puncture, break or burn even when apparently empty

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

[See Annex I - To be completed nationally]

**12. MARKETING AUTHORISATION NUMBER(S)**

[To be completed nationally]

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

[To be completed nationally]

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

[To be completed nationally]



**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**  
**EVOHALER LABEL**

**1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

Seretide Evohaler and associated names (see Annex I) 25/125 microgram/dose pressurised inhalation, suspension

[See Annex I – To be completed nationally]

Salmeterol xinafoate and fluticasone propionate

Inhalation use

**2. METHOD OF ADMINISTRATION**

Shake well before use

Read the package leaflet before use

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

120 puffs

**6. OTHER**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Seretide Evohaler and associated names (see Annex I) 25/250 microgram/dose pressurised inhalation, suspension  
[See Annex I – To be completed nationally]

Salmeterol xinafoate and fluticasone propionate

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

25 micrograms of salmeterol (as xinafoate) and 250 micrograms of fluticasone propionate per actuation

**3. LIST OF EXCIPIENTS**

Norflurane (HFA 134a)

**4. PHARMACEUTICAL FORM AND CONTENTS**

Pressurised inhalation, suspension

1 X 120 puffs

3 X 120 puffs

10 X 120 puffs (hospital or pharmacy use only)

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Shake well before use

Read the package leaflet before use

Inhalation use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

Pressurised container. Do not puncture, break or burn even when apparently empty

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

[See Annex I - To be completed nationally]

**12. MARKETING AUTHORISATION NUMBER(S)**

[To be completed nationally]

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

[To be completed nationally]

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

[To be completed nationally]

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**  
**EVOHALER LABEL**

**1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

Seretide Evohaler and associated names (see Annex I) 25/250 microgram/dose pressurised inhalation, suspension

[See Annex I – To be completed nationally]

Salmeterol xinafoate and fluticasone propionate

Inhalation use

**2. METHOD OF ADMINISTRATION**

Shake well before use

Read the package leaflet before use

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

120 puffs

**6. OTHER**

**PACKAGE LEAFLET**

## PACKAGE LEAFLET: INFORMATION FOR THE USER

Seretide Evohaler and associated names (see Annex I) 25/50 microgram/dose pressurised inhalation, suspension

Seretide Evohaler and associated names (see Annex I) 25/125 microgram/dose pressurised inhalation, suspension

Seretide Evohaler and associated names (see Annex I) 25/250 microgram/dose pressurised inhalation, suspension

[See Annex I – To be completed nationally]

Salmeterol xinafoate and fluticasone propionate

### **Read all of this leaflet carefully before you start taking this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

### **In this leaflet:**

1. What Seretide is and what it is used for
2. Before you take Seretide
3. How to take Seretide
4. Possible side effects
5. How to store Seretide
6. Further information

## **1. WHAT SERETIDE IS AND WHAT IT IS USED FOR**

Seretide is supplied to you in a metered dose inhaler which delivers your medicine in a pressurised suspension for you to inhale. Each puff provides 25 micrograms of salmeterol (as xinafoate) together with either 50, 125 or 250 micrograms of fluticasone propionate.

Salmeterol is one of a group of medicines called long-acting bronchodilators (effect lasting for at least 12 hours). It helps the airways in the lung to stay open making it easier for air to get in and out. Fluticasone propionate is one of a group of medicines called corticosteroids which reduces swelling and irritation in the lungs. This combination of salmeterol and fluticasone propionate when used regularly prevents asthma attacks occurring. Therefore, it must be taken every day as directed by your doctor. Seretide does not work in controlling sudden attacks of breathlessness and wheezing where you would need to use your fast acting 'rescue' medication.

## **2. BEFORE YOU TAKE SERETIDE**

### **Do not take Seretide**

If you are allergic (hypersensitive) to salmeterol xinafoate, fluticasone propionate or to the other ingredient Norflurane (HFA 134a).

### **Take special care with Seretide**

Your doctor will supervise your treatment more closely if you have medical conditions such as heart disease including an irregular or fast heart beat, overactive thyroid gland, high blood pressure, diabetes mellitus (Seretide may increase your blood sugar), low potassium in your blood or if you are being, or have been treated for have tuberculosis (TB).

If you are using high doses of Seretide for a long time, one of the active ingredients, fluticasone propionate, may stop the natural production of steroid hormones by the adrenal gland. This may cause thinning of the bones, cataract, glaucoma, weight gain, rounded (moon shaped) face, high blood pressure and slowing of growth in children and adolescents. Your doctor will check you regularly for any of these side effects and to make sure you are taking the lowest dose of Seretide to control your asthma.

Very rarely side effects may occur if you have taken high doses of Seretide for a long time and you stop or reduce your dose suddenly. Side effects may also occur if you get an infection or at times of extreme stress (such as after a serious accident or if you have surgery). These effects include stomach pain, tiredness, loss of appetite, sickness, diarrhoea, weight loss, headache or drowsiness, low levels of potassium in your blood, low blood pressure and seizures. To prevent these symptoms occurring, your doctor may prescribe extra corticosteroids during this time.

### **Taking other medicines**

Please tell your doctor if you are taking or have recently taken any other medicines, including those for asthma or any medicines obtained without a prescription. In some cases, Seretide may not be suitable to be taken with other medicines.

Your doctor will need to know if you have recently been treated with corticosteroids (orally or by injection). This is to reduce the risk of affecting the function of your adrenal gland.

Seretide should not be taken with medicines called  $\beta$ -blockers (such as atenolol, propranolol, sotalol), unless your doctor tells you to do so. Some types of antiviral and antifungal medication (such as ritonavir, ketoconazole and itraconazole) may increase the amount of fluticasone propionate in your body and therefore increase your risk of having side effects. Only take these medicines whilst on Seretide, if told to do so by your doctor.

### **Pregnancy and breast-feeding**

You should ask your doctor for advice before taking Seretide whilst you are pregnant or breastfeeding. The doctor will assess whether you can take Seretide during this time.

### **Driving and using machines**

The possible side effects associated with Seretide are unlikely to affect your ability to drive or use machines.

## **3. HOW TO TAKE SERETIDE**

### **Dosage**

It is very important that you take your Seretide every day as directed, until you are advised to stop by your doctor.

Seretide Evohaler is for inhalation by mouth only.

#### Adults and adolescents 12 years and over

Seretide 25/50 Evohaler	Two puffs twice a day
Seretide 25/125 Evohaler	Two puffs twice a day
Seretide 25/250 Evohaler	Two puffs twice a day

#### Children 4 to 12 years of age

Seretide 25/50 Evohaler	Two puffs twice a day
-------------------------	-----------------------

This medicine is not recommended for use in children below 4 years of age.

The doctor will prescribe the lowest strength of Seretide Evohaler that will best control your symptoms. If your symptoms are well controlled using Seretide Evohaler twice daily, your doctor may decide to reduce your dose to once daily. This may be either once at night if you have night-time symptoms or once in the morning if you have daytime symptoms. It is very important to follow your doctor's instructions as to how many puffs to inhale and how often to take your medicine.

A spacer device may be used to improve treatment with Seretide particularly with young children, if you or your child has difficulty co-ordinating breathing in through your mouth and pressing the can to release a puff at the same time.

If you find it difficult to breathe or your wheezing gets worse after taking Seretide, stop using it immediately and tell your doctor straightaway so they can assess your treatment.

If your asthma gets worse or is not well controlled (you will feel wheezy and need more of your fast acting 'rescue' medicine), do not increase the number of puffs of Seretide you take. See your doctor immediately so they can review your condition and assess the medication you need to take.

Do not use Seretide to treat a sudden attack of breathlessness or wheezing as it will not help you immediately. You must use your fast acting 'rescue' medicine (such as salbutamol) for this purpose which you should carry with you at all times. Be careful not to confuse your Seretide inhaler with your 'rescue' inhaler.

### **Instructions for proper use**

Your doctor, nurse or pharmacist should instruct you in the proper use of your inhaler.


The medicine is contained in a pressurised can in a plastic casing with a mouthpiece. The can has a counter attached to show you how many puffs of medicine are left.



The number will show through a window at the back of the plastic casing.

### Testing the inhaler

If using your inhaler for the first time, remove the mouthpiece cover by holding the sides of the inhaler with your thumb and forefinger. Gently squeeze the sides of the cover and pull apart.



To make sure that it works, shake the inhaler well and press the can to release puffs into the air until the counter reads 120. Each time the can is pressed, a puff of medicine is released and the counter will count down by one.



Take care not to drop the inhaler as this may cause the counter to count down.

If your inhaler has not been used for a week or more remove the mouthpiece cover, shake the inhaler well and release two puffs into the air.

### Use of the inhaler

It is important to start to breathe in as slowly as possible just before using your inhaler.

1. You should be standing or sitting whilst using the inhaler.
2. To remove the mouthpiece cover, hold the sides of the inhaler with your thumb and forefinger, squeeze the cover gently and pull apart. Check inside and outside to make sure that the mouthpiece is clean and that there are no foreign objects.
3. Shake the inhaler before use.



Important, do not rush steps 4 to 7:

4. Hold the inhaler upright with your thumb on the base, below the mouthpiece. Breathe out as far as is comfortable.



5. Place the mouthpiece in your mouth between your teeth and close your lips around it. Do not



bite it.

6. Just after starting to breathe in through your mouth, press down on the top of the can to release a puff while still breathing in steadily and deeply.



7. Hold your breath, take the inhaler from your mouth and your finger from the top of the inhaler. Continue holding your breath for a few seconds, or as long as is comfortable.



8. Rinse your mouth with water and spit it out as this may help prevent the occurrence of thrush and hoarseness.
9. If your doctor has told you to take two puffs, wait about half a minute before you take another puff by repeating steps 3 to 8.
10. After use, always replace the mouthpiece cover immediately to keep out dust. Replace the cover by firmly pushing and clicking into position.

You should get a replacement when the counter shows the number 020. Stop using the inhaler when the counter reads 000. Never try to alter the numbers on the counter or detach the counter from the metal can.

#### Cleaning your inhaler

To prevent your inhaler blocking up, it is important to clean it at least once a week. To clean your inhaler:

- Remove the mouthpiece cover.
- Do not remove the metal can from the plastic casing during cleaning or at any other time.
- Wipe the inside and outside of the mouthpiece and the plastic casing with a dry cloth or tissue.
- Replace the mouthpiece cover.

Do not put the metal container in water.

#### **If you take more Seretide than you should**

It is important to take Seretide as your doctor recommended. If you accidentally take a larger dose, you may develop side effects such as your heart beating faster than usual, feeling shaky, headache, muscle weakness, aching joints and also a reduction of steroid hormones produced by the adrenal

gland. These effects usually wear off with continued treatment, however, if you have used larger doses for a long period of time, you should contact your doctor or pharmacist for advice.

#### **If you forget to take Seretide**

If you forget to take your medicine, take your next dose when it is due. Do not take a double dose to make up for a forgotten dose.

#### **If you stop taking Seretide**

It is very important that you take your Seretide every day as directed, until you are advised to stop by your doctor. Do not stop taking Seretide suddenly as this could make your asthma worse and could cause problems with the body's hormones.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

## **4. POSSIBLE SIDE EFFECTS**

Like all medicines, Seretide can cause side effects, although not everybody gets them. To help prevent side effects occurring, your doctor will make sure you are on the lowest dose of Seretide that will control your asthma. These are the side effects reported by people taking Seretide.

**Very Common effects:** this means that more than 1 in 10 people taking the medicine are affected.

- Headache, although this side effect usually lessens as treatment continues.

**Common effects:** means that between 1 in 10 and 1 in 100 people taking the medicine are affected.

- Thrush (sore, creamy-yellow, raised patches) in the mouth and throat. Soreness of the tongue, throat and hoarseness of the voice. Rinsing the mouth out with water and spitting it out immediately after taking each puff may help. Your doctor may prescribe additional anti fungal medication to treat the thrush.
- Feeling shaky and fast or irregular heartbeat (palpitations). These side effects are usually harmless and lessen as treatment continues.
- Muscle cramps.

**Uncommon effects:** means that between 1 in 100 and 1 in 1000 people taking the medicine are affected.

- Rash
- Very fast heart rate (tachycardia).

**Very rare:** means that fewer than 1 in 10,000 people taking the medicine are affected.

- Signs of allergy, such as itching, swelling (usually of the face, lips, tongue or throat), difficulty in breathing, wheezing or coughing (bronchospasm). If you notice these symptoms or they appear suddenly after taking Seretide, tell your doctor immediately as you could be allergic to Seretide.
- Seretide may affect the normal production of steroid hormones in the body, particularly if you have taken high doses for long periods of time. The effects are slowing of growth in children and adolescents, thinning of the bones, cataract, glaucoma, weight gain, high blood pressure and rounded (moon shaped) face (Cushing's Syndrome).
- Uneven heartbeat or heart gives an extra beat (arrhythmias). Tell your doctor, but do not stop taking Seretide unless they tell you to do so.

- Increases in the amount of sugar (glucose) in your blood (hyperglycaemia). If you have diabetes, more frequent blood sugar monitoring and possibly adjustment of your usual diabetic treatment may be required.
- Feeling anxious, disturbed sleep and behavioural changes, such as being unusually active and irritable (mainly in children).
- Breathing difficulties or wheezing (bronchospasm) worsens immediately after taking Seretide.
- Aching, swollen joints and muscle pain.

If any of the side effects become serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## **5. HOW TO STORE SERETIDE**

Keep out of the reach and sight of children.

Do not store above 25°C.

Do not store Seretide Evohaler in a cold place, as your medicine may not work as well.

The metal can contains a pressurised liquid. Do not puncture, break or burn it even if you think it is empty.

Do not use Seretide after the expiry date which is stated on the label and carton.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## **6. FURTHER INFORMATION**

### **What Seretide contains**

- The active substance(s) are 25 micrograms salmeterol (as xinafoate) and 50, 125 or 250 micrograms fluticasone propionate.
- The other ingredient is Norflurane (HFA 134a).

### **What Seretide looks like and contents of the pack**

Pressurised inhalation, suspension.

The pressurised can contains a white to off white suspension for inhalation.

The devices are packed in cartons of 1, 3 or 10 Evohalers.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

Marketing Authorisation Holder:  
[See Annex I - To be completed nationally]

Manufacturer:  
Glaxo Wellcome Production  
Zone Industrielle No.2, 23 Rue Lavoisier, la madeleine, 27000 Evreux, France.  
Tel: +33 2 3223 5500  
Fax: +33 2 3223 5558

**This medicinal product is authorised in the Member States of the EEA under the following names:**

Austria	Seretide Dosieraerosol
Belgium	Seretide
Denmark	Seretide
Finland	Seretide Evohaler
France	Seretide
Germany	atmadisc
Greece	Seretide Inhaler
Iceland	Seretide
Ireland	Seretide Evohaler
Italy	Seretide
Luxembourg	Seretide
The Netherlands	Seretide
Portugal	Seretide Inalador
Spain	Seretide
Sweden	Seretide Evohaler
United Kingdom	Seretide Evohaler

**This leaflet was last approved in {MM/YYYY}.**

[To be completed nationally]

<Detailed information on this medicine is available on the web site of {MA/Agency}>