

Summary of risk management plan for Wegovy (semaglutide s.c. 2.4 mg for weight management)

This is a summary of the risk management plan (RMP) for Wegovy. The RMP details important risks of Wegovy, how these risks can be minimised and how more information will be obtained about Wegovy's risks and uncertainties (missing information).

Wegovy's Summary of Product Characteristics (SmPC) and its package leaflet give essential information to healthcare professionals and patients on how Wegovy should be used.

This summary of the RMP for Wegovy should be read in the context of all this information including the assessment report of the evaluation and its plain-language summary, all of which is part of the EPAR.

Important new concerns or changes to the current ones will be included in updates of Wegovy's RMP.

I. The medicine and what it is used for

Wegovy is authorised for weight loss and weight maintenance. It contains the active substance semaglutide and it is given by subcutaneous route.

Wegovy is used together with diet and physical activity for weight loss and weight maintenance in adults who have:

- a BMI of 30 kg/m² or greater (with obesity) or
- a BMI of 27 kg/m² and less than 30 kg/m² (overweight) and weight-related health problems.

Wegovy is used together with diet and physical activity for weight management in adolescents ages 12 years and above, who have

- obesity and
- body weight >60 kg

Use of Wegovy should only be continued if patients lost at least 5% of their BMI after 12 weeks on the 2.4 mg dose or maximum tolerated dose.

BMI (body mass index) is a measure of your weight in relation to your height.

Further information about the evaluation of Wegovy's benefits can be found in Wegovy's EPAR, including its plain-language summary, available on the EMA website under the medicine's webpage: [EPAR link](#).

II. Risks associated with the medicine and activities to minimise or further characterise the risks

Important risks of Wegovy, together with measures to minimise such risks and the proposed studies for learning more about Wegovy's risks, are outlined below.

Measures to minimise the risks identified for medicinal products can be:

- Specific information, such as warnings, precautions and advice on correct use, in the package leaflet and SmPC addressed to patients and healthcare professionals
- Important advice on the medicine's packaging
- The authorised pack size – the amount of medicine in a pack is chosen so to ensure that the medicine is used correctly
- The medicine's legal status – the way a medicine is supplied to the public (e.g., with or without prescription) can help to minimise its risks.

Together, these measures constitute *routine risk minimisation* measures.

In addition to these measures, information about adverse events is collected continuously and analysed regularly, including the PSUR assessment, so that immediate action can be taken as necessary. These measures constitute *routine pharmacovigilance activities*.

II.A List of important risks and missing information

Important risks of Wegovy are risks that need special risk management activities to further investigate or minimise the risk, so that the medicinal product can be safely administered. Important risks can be regarded as identified or potential. Identified risks are concerns for which there is sufficient proof of a link with the use of Wegovy. Potential risks are concerns for which an association with the use of this medicine is possible based on available data, but this association has not been established yet and needs further evaluation. Missing information refers to information on the safety of the medicinal product that is currently missing and needs to be collected (e.g., on the long-term use of the medicine).

List of important risks and missing information	
Important identified risks	<ul style="list-style-type: none"> Diabetic retinopathy complications (only for patients with T2D)
Important potential risks	<ul style="list-style-type: none"> Pancreatic cancer Medullary thyroid cancer
Missing information	<ul style="list-style-type: none"> Pregnancy and lactation Patients with severe hepatic impairment

Abbreviations: T2D = type 2 diabetes mellitus.

II.B Summary of important risks

Diabetic retinopathy complications (only for patients with T2D)	
Evidence for linking the risk to the medicine	The risk is included for semaglutide s.c. 2.4 mg for weight management in patients with T2D based on the findings in the semaglutide s.c. for T2D (Ozempic) clinical development programme. In STEP 2 trial, few diabetic retinopathy events were reported, but with higher rates and proportions with semaglutide 2.4 mg than with placebo.
Risk factors and risk groups	Patient risk factors include increasing age, long duration of diabetes, poor glycaemic control, prior history of diabetic retinopathy and rapid decline in HbA _{1c} .
Risk minimisation measures	<p><i>Routine risk minimisation measures:</i> SmPC Sections 4.4 and 4.8, and PL Sections 2 and 4.</p> <p><i>Additional risk minimisation measures:</i> None</p>
Additional pharmacovigilance activities	<p><i>Additional pharmacovigilance activities:</i> Study NN9535-4352 (Long-term effects of semaglutide on diabetic retinopathy in participants with T2D [FOCUS]) will also be relevant for the ongoing evaluation of the risk for semaglutide s.c. 2.4 mg for weight management in patients with T2D.</p> <p>See section below of this summary for an overview of the post-authorisation development plan.</p>

Abbreviations: CVOT = cardiovascular outcomes trial; PL = package leaflet; s.c. = subcutaneous(-ly); SmPC = Summary of Product Characteristics; T2D = type 2 diabetes mellitus.

Pancreatic cancer	
Evidence for linking the risk to the medicine	Patients with T2D, as well as patients being overweight or with obesity, have an increased risk of certain types of cancer such as pancreatic cancer. There is no evidence from clinical trials that GLP-1-based therapies increase the risk of pancreatic cancer. However, pancreatic cancer is

	included as an important potential risk based on the EC regulation 726/2004 Article 5(3) referral procedure in 2013 (EMA/H/A-5(3)/1369).
Risk factors and risk groups	Patient risk factors for neoplasms include diabetes, chronic pancreatitis, obesity, physical inactivity, advanced age, smoking, alcohol abuse, environmental factors, history of neoplasms and family history of pancreatic cancer and other genetic predispositions.
Risk minimisation measures	No risk minimisation measures.
Additional pharmacovigilance activities	<i>Additional pharmacovigilance activities:</i> Study NN9535-4447 (Epidemiological assessment of the risk for pancreatic cancer associated with the use of semaglutide in patients with T2D) will also be relevant for the ongoing evaluation of the risk for semaglutide s.c. 2.4 mg for weight management. See section below of this summary for an overview of the post-authorisation development plan.

Abbreviations: GLP-1 = glucagon-like peptide-1; s.c. = subcutaneous(-ly); T2D = type 2 diabetes mellitus.

Medullary thyroid cancer	
Evidence for linking the risk to the medicine	This potential class risk is based on findings in mice and rats for all currently approved long-acting GLP-1 RAs. Data from the monitoring of calcitonin (a marker for MTC) in plasma in the semaglutide s.c. and oral semaglutide clinical development programmes did not support a semaglutide effect on calcitonin in humans.
Risk factors and risk groups	Patient risk factors for MTC include previous family history or personal medical history of multiple endocrine neoplasia 2 (MEN2), a group of medical disorders associated with tumours of the endocrine system.
Risk minimisation measures	<i>Routine risk minimisation measures:</i> SmPC Section 5.3. <i>Additional risk minimisation measures:</i> None
Additional pharmacovigilance activities	<i>Additional pharmacovigilance activities:</i> Study MTC-22341 (Medullary Thyroid Carcinoma Surveillance Study: a Case-Series Registry) See section below of this summary for an overview of the post-authorisation development plan.

Abbreviations: MEN2 = multiple endocrine neoplasia 2; MTC = medullary thyroid cancer; s.c. = subcutaneous(-ly); SmPC = Summary of Product Characteristics.

Pregnancy and lactation	
Risk minimisation measures	<i>Routine risk minimisation measures:</i> SmPC Section 4.6 and PL Section 2. <i>Additional risk minimisation measures:</i> None

Abbreviations: PL = package leaflet; SmPC = Summary of Product Characteristics.

Patients with severe hepatic impairment	
Risk minimisation measures	<i>Routine risk minimisation measures:</i> SmPC Sections 4.2 and 5.2. <i>Additional risk minimisation measures:</i> None

Abbreviations: SmPC = Summary of Product Characteristics.

II.C Post-authorisation development plan

II.C.1 Studies which are conditions of the marketing authorisation

There are no studies which are conditions of the marketing authorisation or specific obligation of Wegovy.

II.C.2 Other studies in post-authorisation development plan

NN9535-4352 (FOCUS)

Purpose of the study: The aim of this randomised clinical trial is to establish the long-term effects of semaglutide on diabetic retinopathy in participants with type 2 diabetes mellitus (T2D) using validated and standardised ophthalmic assessments. Results from the trial will be relevant also for the ongoing evaluation of the risk for semaglutide s.c. 2.4 mg for weight management in patients with T2D.

NN9535-4447

Purpose of the study: The aim of this study is to evaluate whether exposure to semaglutide increases the risk of pancreatic cancer in patients with T2D. Results from the study will also be relevant for the ongoing evaluation of the risk for semaglutide s.c. 2.4 mg for weight management

MTC-22341

Purpose of the study: This active surveillance programme for MTC has been established to evaluate further a potential association between treatment with long-acting GLP-1 RAs and the occurrence of MTC in humans. The MTC registry is an FDA post-marketing requirement for long-acting GLP-1 RA products.