

Summary of Risk Management Plan for Yentreve (duloxetine)

This is a summary of the RMP for Yentreve. The RMP details important risks of Yentreve, how these risks can be minimised, and how more information will be obtained about Yentreve's risks and uncertainties (missing information).

Yentreve's SmPC and its package leaflet give essential information to healthcare professionals and patients on how Yentreve should be used.

This summary of the RMP for Yentreve should be read in the context of all this information including the assessment report of the evaluation and its plain-language summary, all of which is part of the EPAR.

Important new concerns or changes to the current concerns will be included in updates of Yentreve's RMP.

I - The Medicine and What It Is Used for

Yentreve is authorised for SUI (see the SmPC for the full indication). Yentreve contains duloxetine as the active substance and is given by oral administration.

Further information about the evaluation of Yentreve's benefits can be found in Yentreve's EPAR, including in its plain-language summary, available on the EMA website, under the medicine's webpage:

http://www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/medicines/000545/human_med_001164.jsp&mid=WC0b01ac058001d124.

II - Risks Associated with the Medicine and Activities to Minimise or Further Characterise the Risks

Important risks of Yentreve, together with measures to minimise such risks and the proposed studies for learning more about Yentreve's risks, are outlined below.

Measures to minimise the risks identified for medicinal products can be:

- Specific information, such as warnings, precautions, and advice on correct use, in the package leaflet and SmPC addressed to patients and healthcare professionals.

Together, these measures constitute *routine risk minimisation* measures.

In addition to these measures, information about adverse reactions is collected continuously and regularly analysed including PSUR assessment so that immediate action can be taken as necessary. These measures constitute routine PV activities.

II.A Yentreve List of Important Risks and Missing Information

Important risks of Yentreve are risks that need special risk management activities to further investigate or minimise the risk, so that the medicinal product can be safely taken. Important risks can be regarded as identified or potential. Identified risks are concerns for which there is sufficient proof of a link with the use of Yentreve. Potential risks are concerns for which an association with the use of this medicine is possible based on available data, but this association

has not been established yet and needs further evaluation. Missing information refers to information on the safety of the medicinal product that is currently missing and needs to be collected (e.g., on the long-term use of the medicine).

List of important risks and missing information	
Important identified risks	Suicidality
Important potential risks	None
Missing information	None

II.B Yentreve Summary of Important Risks

Important identified risk: Suicidality	
Evidence for linking the risk to the medicine	<p>Suicidality has been considered a class effect across different antidepressant medicines for many years. With the exception of young adults and paediatric patients, the evidence linking duloxetine with suicidality is weak, with multiple evidence sources indicating that patients taking duloxetine are not at higher risk for suicidality when compared with placebo-treated patients. These data sources include meta-analyses conducted by Lilly and the US Food and Drug Administration (FDA) regulatory agency, as well as data from observational studies.</p> <p>The incidence estimates of suicidality in duloxetine-treated patients vary depending on data source and patient characteristics, but overall in postmarketing reporting (even taking into account recognised limitations of spontaneous data sources), the reporting rate of suicidality is very low. Although the results of the different meta-analyses conducted with duloxetine differ, with 1 analysis indicating a higher statistical risk than another, the overall trend across the class appears to be in children and in patients younger than 25 years old with psychiatric disorders.</p>
Risk factors and risk groups	Lilly's meta-analyses of placebo-controlled studies with duloxetine suggest that there is a potential but not statistically significantly increased risk of suicidal thinking and behaviour in young adults (aged 18 to 24 years) with duloxetine treatment.
Risk minimisation measures	<p>Routine risk minimisation measures: SmPC Sections 4.4 and 4.8 Section 4.4 advises to monitor and supervise patients for suicidality and not to use in treatment of children and adolescents under the age of 18 years</p> <p>Additional risk minimisation measures: No risk minimisation measures</p>

II.C Yentreve Postauthorisation Development Plan

II.C.1 Studies that are Conditions of the Marketing Authorisation

There are no studies that are conditions of the marketing authorisation or specific obligation of Yentreve.

II.C.2 Other Studies in Postauthorisation Development Plan

There are no studies required for Yentreve.