

19 December 2013
EMA/876333/2011 Rev 2* (superseded version)

Guideline on good pharmacovigilance practices (GVP)

Annex I - Definitions (Rev 2)

Draft of first version finalised by the Agency in collaboration with Member States	7 February 2012
Definitions agreed by ERMS FG	24 January 2012
Draft adopted by Executive Director	20 February 2012
Start of public consultation	21 February 2012
End of consultation (deadline for comments)	18 April 2012
Revised draft of first version finalised by the Agency in collaboration with Member States	20 June 2012
Revised draft agreed by ERMS FG	21 June 2012
Revised draft adopted by Executive Director as final	22 June 2012
Date for coming into effect	2 July 2012
Preparation of draft Revision 1	26 November 2012
Draft Revision 1 agreed by ERMS FG	6 December 2012
Draft Revision 1 adopted by Executive Director as final	12 December 2012
Date for coming into effect of Revision 1	13 December 2012
Draft of definitions for GVP P I finalised by the Agency in collaboration with Member States	21 February 2013
Draft agreed by ERMS FG	8 March 2013
Draft adopted by Executive Director	9 April 2013
Start of public consultation	12 April 2013
End of consultation (deadline for comments)	12 June 2013
Revised draft Revision 2* finalised by the Agency in collaboration with Member States	23 October 2013
Draft Revision 2 agreed by ERMS FG	11 November 2013
Draft Revision 2 adopted by Executive Director as final	19 December 2013

See websites for contact details



Date for coming into effect of Revision 2*	8 January 2014
--------------------------------------------	----------------

***Note:** Revision 2 includes the following:

- Addition of definitions consulted with the public and finalised for Module XV (Direct healthcare professional communication);
- Addition of definitions consulted with the public and finalised for Considerations P.I (Immunisation, Immunisation anxiety-related reaction, Immunisation error-related reaction, Target population (vaccine), Vaccination, Vaccination failure, Vaccine failure, Vaccine pharmacovigilance, Vaccine product-related reaction, Vaccine quality defect-related reaction);
- Addition of definitions from the European Union Regulatory Network Incident Management Plan for Medicines for Human Use (Crisis, Incident);
- Amendments to the definitions of Missing information and Safety concern in order to meet EU legal requirements;
- Move of definition of Important missing information to explanatory note to definition of Missing information;
- Addition of explanatory notes for the definition of Off-label use in accordance with Module V;
- Addition of an explanatory note on abbreviations used in the definition for Immunological medicinal product.

This version is **not valid anymore**, but kept on the Agency's website for the purpose of public access to historical documents. For the valid version, please refer to the Agency's GVP webpage for the latest revision of this GVP Annex.

Table of contents

Abuse of a medicinal product.....	6
Advanced therapy medicinal product (ATMP)	6
Adverse event (AE); synonym: Adverse experience.....	6
Adverse event following immunisation (AEFI)	6
Adverse reaction; synonyms: Adverse drug reaction (ADR), Suspected adverse (drug) reaction, Adverse effect, Undesirable effect.....	6
Audit	6
Audit finding(s).....	6
Audit plan	7
Audit programme.....	7
Audit recommendation	7
Clinical trial.....	7
Closed signal	7
Company core data sheet (CCDS)	8
Company core safety information (CCSI)	8
Compassionate use of a medicinal product.....	8
Completed clinical trial	8
Consumer	8
Crisis.....	8
Data lock point	9
Development international birth date (DIBD).....	9
Development safety update report (DSUR).....	9
Direct healthcare professional communication (DHPC)	9
EU reference date; synonym: Union reference date.....	9
Failure to vaccinate.....	9
Generic medicinal product.....	10
Good pharmacovigilance practices (GVP) for the European Union.....	10
Healthcare professional	10
Herbal medicinal product	10
Homeopathic medicinal product	10
Identified risk.....	10
Illegal purposes	11
Immunological medicinal product.....	11
Immunisation.....	11
Immunisation anxiety-related reaction	11
Immunisation error-related reaction.....	12
Important identified risk and Important potential risk.....	12
Important potential risk	12
Incident.....	12
Individual case safety report (ICSR); synonym: Adverse (drug) reaction report.....	13
International birth date (IBD)	13
Investigational drug	13
Investigational medicinal product.....	13
Labelling.....	13

Medicinal product.....	13
Medicinal product derived from human blood or human plasma	13
Minimum criteria for reporting	14
Missing information	14
Misuse of a medicinal product.....	14
Misuse of a medicinal product for illegal purposes	14
Name of the medicinal product	14
Newly identified signal.....	14
Non-interventional trial; synonym: Non-interventional study.....	15
Occupational exposure to a medicinal product	15
Off-label use	15
Ongoing clinical trial.....	15
Ongoing signal	16
Overdose	16
Package leaflet	16
Periodic safety update report (PSUR).....	16
Pharmacovigilance	16
Pharmacovigilance system	16
Pharmacovigilance system master file (PSMF)	17
Post-authorisation safety study (PASS).....	17
Potential risk	17
Quality adherence.....	17
Quality assurance	17
Quality control and assurance.....	17
Quality improvements	18
Quality of a pharmacovigilance system	18
Quality objectives	18
Quality planning	18
Quality requirements.....	18
Quality system of a pharmacovigilance system	18
Reference safety information.....	18
Registry.....	19
Risk-benefit balance	19
Risk management plan (RMP).....	19
Risk management system	19
Risk minimisation activity; synonym: Risk minimisation measure.....	19
Risks related to use of a medicinal product.....	19
Safety concern	19
Serious adverse reaction.....	20
Signal.....	20
Signal management process.....	20
Signal validation	20
Solicited sources of individual case safety reports	21
Spontaneous report, synonym: Spontaneous notification	21
Stimulated reporting	21
Substance.....	21

Summary of product characteristics (SmPC)	21
Target population (treatment); synonym: Treatment target population	22
Target population (vaccine); synonym: Vaccine target population	22
Traditional herbal medicinal product	22
Unexpected adverse reaction	22
Upper management	22
Vaccination	23
Vaccination failure	23
Vaccine	23
Vaccine failure	23
Vaccine pharmacovigilance	24
Vaccine product-related reaction	24
Vaccine quality defect-related reaction	24
Valid individual case safety report	25
Validated signal	25

Abuse of a medicinal product

Persistent or sporadic, intentional excessive use of medicinal products which is accompanied by harmful physical or psychological effects [DIR 2001/83/EC Art 1(16)].

Advanced therapy medicinal product (ATMP)

A medicinal product for human use that is either a gene therapy medicinal product, a somatic cell therapy product or a tissue engineered products as defined in Regulation (EC) No 1394/2007 [Reg (EC) No 1394/2007 Art 1(1)].

Adverse event (AE); synonym: Adverse experience

Any untoward medical occurrence in a patient or clinical trial subject administered a medicinal product and which does not necessarily have a causal relationship with this treatment [Dir 2001/20/EC Art 2(m)].

An adverse event can therefore be any unfavourable and unintended sign (e.g. an abnormal laboratory finding), symptom, or disease temporally associated with the use of a medicinal product, whether or not considered related to the medicinal product.

Adverse event following immunisation (AEFI)

See Vaccine pharmacovigilance, Vaccine product-related reaction, Vaccine quality defect-related reaction, Immunisation error-related reaction, Immunisation anxiety-related reaction

Adverse reaction; synonyms: Adverse drug reaction (ADR), Suspected adverse (drug) reaction, Adverse effect, Undesirable effect

A response to a medicinal product which is noxious and unintended [DIR 2001/83/EC Art 1(11)]¹.

Response in this context means that a causal relationship between a medicinal product and an adverse event is at least a reasonable possibility (see Annex IV, ICH-E2A Guideline).

Adverse reactions may arise from use of the product within or outside the terms of the marketing authorisation or from occupational exposure [DIR 2001/83/EC Art 101(1)]. Conditions of use outside the marketing authorisation include off-label use, overdose, misuse, abuse and medication errors.

See also Adverse event, Serious adverse reaction, Unexpected adverse reaction, Off-label use, Overdose, Misuse of a medicinal product, Abuse of a medicinal product, Occupational exposure to a medicinal product

Audit

A systematic, disciplined, independent and documented process for obtaining audit evidence and evaluating it objectively to determine the extent to which the audit criteria are fulfilled (see ISO 19011 (3.1)²).

Audit finding(s)

Results of the evaluation of the collected audit evidence against audit criteria (see ISO19011 (3.4)³).

¹ In the context of clinical trials, an adverse reaction is defined as all untoward and unintended responses to an investigational medicinal product related to any dose administered [Dir 2001/20/EC Art 2(n)].

² International Organization for Standardization (ISO); www.iso.org

Audit evidence is necessary to support the auditor's results of the evaluation, i.e. the auditor's opinion and report. It is cumulative in nature and is primarily obtained from audit procedures performed during the course of the audit.

See also Audit

Audit plan

Description of activities and arrangement for an individual audit (see ISO19011 (3.12)⁴).

See also Audit

Audit programme

Set of one or more audits planned for a specific timeframe and directed towards a specific purpose (see ISO 19011 (3.11)⁵).

See also Audit

Audit recommendation

Describes the course of action management might consider to rectify conditions that have gone awry, and to mitigate weaknesses in systems of management control (see Sawyer LB et al, 2003⁶).

Audit recommendations should be positive and as specific as possible. They should also identify who is to act on them (Sawyer LB et al, 2003⁶).

See also Audit

Clinical trial

Any investigation in human subjects intended to discover or verify the clinical, pharmacological and/or other pharmacodynamic effects of one or more investigational medicinal product(s), and/or to identify any adverse reactions to one or more investigational medicinal product(s) and/or to study absorption, distribution, metabolism and excretion of one or more investigational medicinal product(s) with the objective of ascertaining its (their) safety and/or efficacy. This includes clinical trials carried out in either one site or multiple sites, whether in one or more Member State [Dir 2001/20/EC Art 2(a)].

See also Ongoing clinical trial, Completed clinical trial, Investigational medicinal product

Closed signal

In periodic benefit-risk evaluation reports, a signal for which an evaluation was completed during the reporting interval (see Annex IV, ICH-E2C(R2) Guideline).

This definition is also applicable to periodic safety update reports.

See also Signal

³ International Organization for Standardization (ISO); www.iso.org

⁴ International Organization for Standardization (ISO); www.iso.org

⁵ International Organization for Standardization (ISO); www.iso.org

⁶ Sawyer LB, Dittenhofer MA. Sawyer's Internal Auditing. 5th ed. Altamonte Springs, FL: The IIA Research Foundation; 2003.

Company core data sheet (CCDS)

For medicinal products, a document prepared by the marketing authorisation holder containing, in addition to safety information, material related to indications, dosing, pharmacology and other information concerning the product (see Annex IV, ICH-E2C(R2) Guideline).

See also Company core safety information

Company core safety information (CCSI)

For medicinal products, all relevant safety information contained in the company core data sheet prepared by the marketing authorisation holder and which the marketing authorisation holder requires to be listed in all countries where the company markets the product, except when the local regulatory authority specifically requires a modification (see Annex IV, ICH-E2C(R2) Guideline).

It is the reference information by which listed and unlisted are determined for the purposes of periodic reporting for marketed products, but not by which expected and unexpected are determined for expedited reporting (see Annex IV, ICH-E2C(R2) Guideline).

See also Company core data sheet

Compassionate use of a medicinal product

Making a medicinal product available for compassionate reasons to a group of patients with a chronically or seriously debilitating disease or whose disease is considered to be life-threatening, and who cannot be treated satisfactorily by an authorised medicinal product (the medicinal product concerned must either be subject of an application for a central marketing authorisation or must be undergoing clinical trials) [REG (EC) No 726/2004 Art 83(2)].

Completed clinical trial

Study for which a final clinical study report is available (see ICH-E2F Guideline, Volume 10 of the Rules Governing Medicinal Products in the EU).

See also Clinical trial

Consumer

For the purpose of reporting cases of suspected adverse reactions, a person who is not a healthcare professional such as a patient, lawyer, friend or relative/parent/child of a patient (see Annex IV, ICH-E2D Guideline).

Crisis

In the context of the European Union Regulatory Network Incident Management Plan for Medicines for Human Use, a crisis is defined as a situation where, after assessment of the associated risks, urgent and coordinated action within the EU regulatory network is required to manage and control the situation (see European Union Regulatory Network Incident Management Plan for Medicines for Human Use⁷).

See also Incident

⁷ www.ema.europa.eu

Data lock point

For a periodic safety update report (PSUR), the date designated as the cut-off date for data to be included in a PSUR.

For a periodic benefit-risk evaluation report (PBRER), the date designated as the cut-off date for data to be included in a PBRER, based on the international birth date (see Annex IV, ICH-E2C(R2) Guideline).

For a development safety update report (DSUR), the date designated as the cut-off date for data to be included in a DSUR, based on the development international birth date (see ICH-E2F Guideline, Volume 10 of the Rules Governing Medicinal Products in the EU).

Date includes day and month (see ICH-E2F Guideline, Volume 10 of the Rules Governing Medicinal Products in the EU).

See also Periodic safety update report, Development safety update report, International birth date, Development international birth date

Development international birth date (DIBD)

Date of first approval (or authorisation) for conducting an interventional clinical trial in any country (see ICH-E2F Guideline, Volume 10 of the Rules Governing Medicinal Products in the EU).

Development safety update report (DSUR)

Format and content for periodic reporting on drugs under development (see ICH-E2F Guideline, Volume 10 of the Rules Governing Medicinal Products in the EU).

Direct healthcare professional communication (DHPC)

A communication intervention by which important information is delivered directly to individual healthcare professionals by a marketing authorisation holder or by a competent authority, to inform them of the need to take certain actions or adapt their practices in relation to a medicinal product.

DHPCs are not replies to enquiries from healthcare professionals.

EU reference date; synonym: Union reference date

For medicinal products containing the same active substance or the same combination of active substances, the date of the first marketing authorisation in the EU of a medicinal product containing that active substance or that combination of active substances; or if this date cannot be ascertained, the earliest of the known dates of the marketing authorisations for a medicinal product containing that active substance or that combination of active substances [DIR 2001/83/EC Art 107c(5)].

Failure to vaccinate

An indicated vaccine was not administered appropriately for any reason (see CIOMS-WHO⁸).

For interpreting what is appropriate, consider the explanatory note for Immunisation error-related reaction.

See also Vaccination failure

⁸ Council for International Organizations of Medical Sciences (CIOMS). Definition and application of terms of vaccine pharmacovigilance (report of CIOMS/WHO Working Group on Vaccine Pharmacovigilance). Genève: CIOMS; 2012.

Generic medicinal product

A medicinal product which has the same qualitative and quantitative composition in active substances and the same pharmaceutical form as the reference medicinal product, and whose bioequivalence with the reference medicinal product has been demonstrated by appropriate bioavailability studies [REG (EC) No 726/2004 Art 10(2)(b)].

Good pharmacovigilance practices (GVP) for the European Union

A set of guidelines for the conduct of pharmacovigilance in the EU, drawn up based on Article 108a of Directive 2001/83/EC, by the European Medicines Agency in cooperation with competent authorities in Member States and interested parties, and applying to marketing authorisation holders in the EU, the Agency and competent authorities in Member States.

Healthcare professional

For the purposes of reporting suspected adverse reactions, healthcare professionals are defined as medically qualified persons, such as physicians, dentists, pharmacists, nurses and coroners (see Annex IV, ICH-E2D Guideline).

Herbal medicinal product

Any medicinal product, exclusively containing as active ingredients one or more herbal substances or one or more herbal preparations, or one or more such herbal substances in combination with one or more such herbal preparations [DIR 2001/83/EC Art 1(30)].

Herbal substances are all mainly whole, fragmented or cut plants, plant parts, algae, fungi, lichen in an unprocessed, usually dried, form, but sometimes fresh. Certain exudates that have not been subjected to a specific treatment are also considered to be herbal substances. Herbal substances are precisely defined by the plant part used and the botanical name according to the binominal system [DIR 2001/83/EC Art 1(31)].

Herbal preparations are preparations obtained by subjecting herbal substances to treatments such as extraction, distillation, expression, fractionation, purification, concentration or fermentation. These include comminuted or powered herbal substances, tinctures, extracts, essential oils, expressed juices and processed exudates [DIR 2001/83/EC Art 1(32)].

Homeopathic medicinal product

Any medicinal product prepared from substances called homeopathic stocks in accordance with a homeopathic manufacturing procedure described by the European Pharmacopoeia or, in the absence thereof, by the pharmacopoeias currently used officially in the Member States. A homeopathic medicinal product may contain a number of principles [DIR 2001/83/EC Art 1(5)].

Identified risk

An untoward occurrence for which there is adequate evidence of an association with the medicinal product of interest (see ICH-E2F Guideline, Volume 10 of the Rules Governing Medicinal Products in the EU).

Examples include:

- an adverse reaction adequately demonstrated in non-clinical studies and confirmed by clinical data;

- an adverse reaction observed in well-designed clinical trials or epidemiological studies for which the magnitude of the difference, compared with the comparator group on a parameter of interest suggests a causal relationship;
- an adverse reaction suggested by a number of well-documented spontaneous reports where causality is strongly supported by temporal relationship and biological plausibility, such as anaphylactic reactions or application site reactions (see ICH-E2F Guideline, Volume 10 of the Rules Governing Medicinal Products in the EU).

In a clinical trial, the comparator may be placebo, an active substance or non-exposure.

Adverse reactions included in section 4.8 of the summary of product characteristics (SmPC) are also considered identified risks, unless they are class-related reactions which are mentioned in the SmPC but which are not specifically described as occurring with this product (these would normally be considered as a potential risk)).

See also Risks related to use of a medicinal product, Important identified risk and Important potential risk, Missing information, Unexpected adverse reaction

Illegal purposes

See Misuse for illegal purposes

Immunological medicinal product

Any medicinal product consisting of vaccines, toxins, serums or allergen products:

Vaccines, toxins and serums shall cover in particular agents used to produce active immunity (such as cholera vaccine, BCG, polio vaccine, smallpox vaccine), agents used to diagnose the state of immunity (including in particular tuberculin and tuberculin PPD, toxins for the Schick and Dick Tests, brucellin) and agents used to produce passive immunity (such as diphtheria antitoxin, anti-smallpox globulin, antilymphocytic globulin).

Allergen products shall mean any medicinal product which is intended to identify or induce a specific acquired alteration in the immunological response to an allergizing agent [DIR 2001/83/EC Art 1(4)].

BCG stands for Bacillus Calmette-Guérin vaccine and PPD for purified protein derivative.

Immunisation

The process of making a person immune.

For the context of Considerations P.I, immunisation refers to the process of making a person immune to an infection.

See also Vaccination

Immunisation anxiety-related reaction

An adverse event following immunisation arising from anxiety about the immunisation (see CIOMS-WHO⁹).

In this definition immunisation means the usage (handling, prescribing and administration) of a vaccine for the purpose of immunising individuals (see CIOMS-WHO⁹), which in the EU is preferably referred to as vaccination (in

⁹ Council for International Organizations of Medical Sciences (CIOMS). Definition and application of terms of vaccine pharmacovigilance (report of CIOMS/WHO Working Group on Vaccine Pharmacovigilance). Genève: CIOMS; 2012.

the report of CIOMS/WHO Working Group on Vaccine Pharmacovigilance the terms immunisation and vaccination are used interchangeably⁹).

See also Adverse reaction, Vaccine pharmacovigilance, Vaccination

Immunisation error-related reaction

An adverse event following immunisation that is caused by inappropriate vaccine handling, prescribing or administration and thus by its nature is preventable (see CIOMS-WHO¹⁰).

In this definition immunisation means the usage (handling, prescribing and administration) of a vaccine for the purpose of immunising individuals (see CIOMS-WHO¹⁰), which in the EU is preferably referred to as vaccination (in the report of CIOMS/WHO Working Group on Vaccine Pharmacovigilance the terms immunisation and vaccination are used interchangeably¹⁰).

Inappropriate refers to usage (handling, prescribing and administration) other than what is licensed and recommended in a given jurisdiction based on scientific evidence or expert recommendations (see CIOMS-WHO¹⁰).

See also Adverse reaction, Vaccine pharmacovigilance, Vaccination

Important identified risk and Important potential risk

An identified risk or potential risk that could have an impact on the risk-benefit balance of the product or have implications for public health (see ICH-E2F Guideline, Volume 10 of the Rules Governing Medicinal Products in the EU).

What constitutes an important risk will depend upon several factors, including the impact on the individual, the seriousness of the risk and the impact on public health. Normally, any risk that is likely to be included in the contraindications or warnings and precautions section of the product information should be considered important (see Annex IV, ICH-E2C(R2) Guideline).

See also Risk-benefit balance, Identified risk, Potential risk, Safety concern

Important potential risk

See Important identified risk and Important potential risk

Incident

A situation where an event occurs or new information arises, irrespective whether this is in the public domain or not, in relation to (an) authorised medicinal product(s) which could have a serious impact on public health.

The incident may be related to quality, efficacy or safety concerns, but most likely to safety and/or quality (and possibly subsequent supply shortages). In addition, situations that do not seem at a first glance to have a serious impact on public health, but are in the public domain - subject of media attention or not- and may lead to serious public concerns about the product, may also need to be considered as incidents. Likewise, other situations which might have a negative impact on the appropriate use of a medicinal products (e.g. resulting in patients stop taking their medicine) may fall within the definition of an incident.

In the context of this the European Union Regulatory Network Incident Management Plan for Medicines for Human Use Incident Management Plan, an incident relates to (a) medicinal product(s) authorised in the EU, irrespective of their route of authorisation.

¹⁰ Council for International Organizations of Medical Sciences (CIOMS). Definition and application of terms of vaccine pharmacovigilance (report of CIOMS/WHO Working Group on Vaccine Pharmacovigilance). Genève: CIOMS; 2012.

Individual case safety report (ICSR); synonym: Adverse (drug) reaction report

Format and content for the reporting of one or several suspected adverse reactions to a medicinal product that occur in a single patient at a specific point of time¹¹.

See also Minimum criteria for reporting

International birth date (IBD)

The date of the first marketing authorisation for any product containing the active substance granted to any company in any country in the world (see Annex IV, ICH-E2C(R2) Guideline).

Investigational drug

Experimental product under study or development. This term is more specific than investigational medicinal product, which includes comparators and placebos (see ICH-E2F Guideline, Volume 10 of the Rules Governing Medicinal Products in the EU).

See also Investigational medicinal product

Investigational medicinal product

An investigational medicinal product is a pharmaceutical form of an active substance or placebo being tested or used as a reference in a clinical trial, including products already with a marketing authorisation but used or assembled (formulated or packaged) in a way different from the authorised form, or when used for an unauthorised indication, or when used to gain further information about the authorised form [Dir 2001/20/EC Art 2(d)].

See also Clinical trial

Labelling

Information on the immediate or outer packaging [DIR 2001/83/EC Art 1(25)].

Medicinal product

Any substance or combination of substances

- presented as having properties for treating or preventing disease in human beings; or
- which may be used in or administered to human beings either with a view to restoring, correcting or modifying physiological functions by exerting a pharmacological, immunological or metabolic action, or to making a medical diagnosis [DIR 2001/83/EC Art 1(2)].

Medicinal product derived from human blood or human plasma

Any medicinal product based on blood constituents which is prepared industrially by a public or private establishment, such as a medicinal product including, in particular, albumin, coagulating factor(s) and immunoglobulin(s) of human origin [DIR 2001/83/EC Art 1(10)].

¹¹ In the context of a clinical trial, an individual case is the information provided by a primary source to describe suspected unexpected serious adverse reactions related to the administration of one or more investigational medicinal products to an individual patient at a particular point of time.

Minimum criteria for reporting

For the purpose of reporting cases of suspected adverse reactions, the minimum data elements for a case are: an identifiable reporter, an identifiable patient, an adverse reaction and a suspect medicinal product (see Annex IV, ICH-E2D Guideline).

For the purpose of validation of individual case safety reports as qualifying for reporting in the EU, see Module VI.

See also Individual case safety report

Missing information

Gaps in knowledge, related to safety or particular patient populations, which could be clinically significant.

It is noted that there is an ICH definition for important missing information, which is: critical gaps in knowledge for specific safety issues or populations that use the marketed product (see Annex IV, ICH-E2C(R2) Guideline).

Misuse of a medicinal product

Situations where the medicinal product is intentionally and inappropriately used not in accordance with the authorised product information.

See also Misuse of a medicinal product for illegal purposes

Misuse of a medicinal product for illegal purposes

Misuse for illegal purposes is misuse with the additional connotation of an intention of misusing the medicinal product to cause an effect in another person. This includes, amongst others: the sale, to other people, of medicines for recreational purposes and use of a medicinal product to facilitate assault.

See also Misuse of a medicinal product

Name of the medicinal product

The name which may be either an invented name not liable to confusion with the common name, or a common or scientific name accompanied by a trade mark or the name of the marketing authorisation holder [DIR 2001/83/EC Art 1(20)].

The common name is the international non-proprietary name (INN) recommended by the World Health Organization, or, if one does not exist, the usual common name [DIR 2001/83/EC Art 1(21)].

The complete name of the medicinal product is the name of the medicinal product followed by the strength and pharmaceutical form.

Newly identified signal

In periodic benefit-risk evaluation reports, a signal first identified during the reporting interval, prompting further actions or evaluation (see Annex IV, ICH-E2C(R2) Guideline).

This definition could also apply to a previously closed signal for which new information becomes available in the reporting interval prompting further action or evaluation (see Annex IV, ICH-E2C(R2) Guideline).

This definition is also applicable to periodic safety update reports.

See also Signal, Closed signal

Non-interventional trial; synonym: Non-interventional study

A study where the medicinal product(s) is (are) prescribed in the usual manner in accordance with the terms of the marketing authorisation. The assignment of the patient to a particular therapeutic strategy is not decided in advance by a trial protocol but falls within current practice and the prescription of the medicine is clearly separated from the decision to include the patient in the study. No additional diagnostic or monitoring procedures shall be applied to the patients and epidemiological methods shall be used for the analysis of collected data [Dir 2001/20/EC Art 2(c)].

Thus, a trial is non-interventional if the following requirements are cumulatively fulfilled:

- the medicinal product is prescribed in the usual manner in accordance with the terms of the marketing authorisation;
- the assignment of the patient to a particular therapeutic strategy is not decided in advance by a trial protocol but falls within current practice and the prescription of the medicine is clearly separated from the decision to include the patient in the study; and
- no additional diagnostic or monitoring procedures are applied to the patients and epidemiological methods are used for the analysis of collected data (see Volume 10 of the Rules Governing Medicinal Products in the EU, Questions & Answers Version 10.0).

Non-interventional studies are defined by the methodological approach used and not by the scientific objectives. Non-interventional studies include database research or review of records where all the events of interest have already happened (this may include case-control, cross-sectional, cohort and other study designs making secondary use of data). Non-interventional studies also include those involving primary data collection (e.g. prospective observational studies and registries in which the data collected derive from routine clinical care), provided that the conditions set out above are met. In these studies, interviews, questionnaires and blood samples may be performed as normal clinical practice.

Non-interventional trials do not fall in the scope of Directive 2001/20/EC.

Occupational exposure to a medicinal product

For the purpose of reporting cases of suspected adverse reactions, an exposure to a medicinal product as a result of one's professional or non-professional occupation.

Off-label use

Situations where a medicinal product is intentionally used for a medical purpose not in accordance with the authorised product information.

Off-label use includes use in non-authorised paediatric age categories. Unless specifically requested, it does not include use outside the EU in an indication authorised in that territory which is not authorised in the EU.

Ongoing clinical trial

Trial where enrolment has begun, whether a hold is in place or analysis is complete, but for which a final clinical study report is not available (see ICH-E2F Guideline, Volume 10 of the Rules Governing Medicinal Products in the EU).

See also Clinical trial, Completed clinical trial

Ongoing signal

In periodic benefit-risk evaluation reports, a signal that remains under evaluation at the data lock point (see Annex IV, ICH-E2C(R2) Guideline).

This definition is also applicable to periodic safety update reports.

See also Signal, Data lock point

Overdose

Administration of a quantity of a medicinal product given per administration or cumulatively which is above the maximum recommended dose according to the authorised product information. Clinical judgement should always be applied.

Package leaflet

A leaflet containing information for the user which accompanies the medicinal product [Dir 2011/83/EC Art 1(26)].

Periodic safety update report (PSUR)

Format and content for providing an evaluation of the risk-benefit balance of a medicinal product for submission by the marketing authorisation holder at defined time points during the post-authorisation phase.

In the EU, periodic safety update reports should follow the format described in Module VII.

Pharmacovigilance

Science and activities relating to the detection, assessment, understanding and prevention of adverse effects or any other medicine-related problem (see WHO¹²).

In line with this general definition, underlying objectives of pharmacovigilance in accordance with the applicable EU legislation for are:

- preventing harm from adverse reactions in humans arising from the use of authorised medicinal products within or outside the terms of marketing authorisation or from occupational exposure; and
- promoting the safe and effective use of medicinal products, in particular through providing timely information about the safety of medicinal products to patients, healthcare professionals and the public.

Pharmacovigilance is therefore an activity contributing to the protection of patients' and public health.

Pharmacovigilance system

A system used by the marketing authorisation holder and by Member States to fulfil the tasks and responsibilities listed in Title IX of Directive 2001/83/EC and designed to monitor the safety of authorised medicinal products and detect any change to their risk-benefit balance [DIR 2001/83/EC Art 1(28d)].

In general, a pharmacovigilance system is a system used by an organisation to fulfil its legal tasks and responsibilities in relation to pharmacovigilance and designed to monitor the safety of authorised medicinal products and detect any change to their risk-benefit balance.

¹² World Health Organization (WHO). The importance of pharmacovigilance: safety monitoring of medicinal products. Genève: WHO; 2002.

Pharmacovigilance system master file (PSMF)

A detailed description of the pharmacovigilance system used by the marketing authorisation holder with respect to one or more authorised medicinal products [DIR 2001/83/EC Art 1(28e)].

See also Pharmacovigilance system

Post-authorisation safety study (PASS)

Any study relating to an authorised medicinal product conducted with the aim of identifying, characterising or quantifying a safety hazard, confirming the safety profile of the medicinal product, or of measuring the effectiveness of risk management measures [DIR 2001/83/EC Art 1(15)].

A post-authorisation safety study may be an interventional clinical trial or may follow an observational, non-interventional study design.

See also Clinical trial, Non-interventional trial

Potential risk

An untoward occurrence for which there is some basis for suspicion of an association with the medicinal product of interest but where this association has not been confirmed (see ICH-E2F Guideline, Volume 10 of the Rules Governing Medicinal Products in the EU).

Examples include:

- non-clinical toxicological findings that have not been observed or resolved in clinical studies;
- adverse events observed in clinical trials or epidemiological studies for which the magnitude of the difference, compared with the comparator group (placebo or active substance, or unexposed group), on the parameter of interest raises a suspicion of, but is not large enough to suggest, a causal relationship;
- a signal arising from a spontaneous adverse reaction reporting system;
- an event known to be associated with other active substances within the same class or which could be expected to occur based on the properties of the medicinal product (see ICH-E2F Guideline, Volume 10 of the Rules Governing Medicinal Products in the EU).

See also Adverse event, Signal

Quality adherence

Carrying out tasks and responsibilities in accordance with quality requirements [IR 520/2012 Art 8(3)].

See also Quality requirements

Quality assurance

See Quality control and assurance

Quality control and assurance

Monitoring and evaluating how effectively the structures and processes have been established and how effectively the processes are being carried out [IR 520/2012 Art 8(3)].

This applies for the purpose of fulfilling quality requirements.

See also Quality requirements

Quality improvements

Correcting and improving the structures and processes where necessary [IR 520/2012 Art 8(3)].

This applies for the purpose of fulfilling quality requirements.

See also Quality requirements

Quality of a pharmacovigilance system

All characteristics of the pharmacovigilance system which are considered to produce, according to estimated likelihoods, outcomes relevant to the objectives of pharmacovigilance.

See also Pharmacovigilance system, Quality system of a pharmacovigilance system

Quality objectives

See Quality requirements

Quality planning

Establishing structures and planning integrated and consistent processes [IR 520/2012 Art 8(3)].

This applies for the purpose of fulfilling quality requirements.

See also Quality requirements

Quality requirements

Those characteristics of a system that are likely to produce the desired outcome, or quality objectives.

See also Pharmacovigilance system, Quality system of a pharmacovigilance system

Quality system of a pharmacovigilance system

The organisational structure, responsibilities, procedures, processes and resources of the pharmacovigilance system as well as appropriate resource management, compliance management and record management [IR 520/2012 Art 8(2)].

The quality system is part of the pharmacovigilance system.

See also Pharmacovigilance system, Quality of a pharmacovigilance system

Reference safety information

In periodic benefit-risk evaluation reports for medicinal products, all relevant safety information contained in the reference product information (e.g. the company core data sheet) prepared by the marketing authorisation holder and which the marketing authorisation holder requires to be listed in all countries where it markets the product, except when the local regulatory authority specifically requires a modification (see Annex IV, ICH-E2C(R2) Guideline).

It is a subset of information contained within the marketing authorisation holder's reference product information for the periodic benefit-risk evaluation report. Where the reference product information is the company core data sheet, the reference safety information is the company core safety information (see Annex IV, ICH-E2C(R2) Guideline).

See also Company core data sheet, Company core safety information

Registry

An organised system that uses observational methods to collect uniform data on specified outcomes in a population defined by a particular disease, condition or exposure.

Risk-benefit balance

An evaluation of the positive therapeutic effects of the medicinal product in relation to the risks [DIR 2001/83/EC Art 1(28a)], i.e. any risk relating to the quality, safety or efficacy of the medicinal product as regards patients' health or public health [DIR 2001/83/EC Art 1(28)].

See also Risks related to use of a medicinal product

Risk management plan (RMP)

A detailed description of the risk management system [DIR 2001/83/EC Art 1(28c)].

To this end, it must identify or characterise the safety profile of the medicinal product(s) concerned, indicate how to characterise further the safety profile of the medicinal product(s) concerned, document measures to prevent or minimise the risks associated with the medicinal product, including an assessment of the effectiveness of those interventions and document post-authorisation obligations that have been imposed as a condition of the marketing authorisation [IR 520/2012 Art 30].

See also Risk management system, Risk minimisation activity

Risk management system

A set of pharmacovigilance activities and interventions designed to identify, characterise, prevent or minimise risks relating to a medicinal product, including the assessment of the effectiveness of those interventions [DIR 2001/83/EC Art 1(28b)].

Risk minimisation activity; synonym: Risk minimisation measure

A public health intervention intended to prevent or reduce the probability of the occurrence of an adverse reaction associated with the exposure to a medicine, or to reduce its severity should it occur.

These activities may consist of routine risk minimisation (e.g. product information) or additional risk minimisation activities (e.g. healthcare professional or patient communications/educational materials).

Risks related to use of a medicinal product

Any risk relating to the quality, safety or efficacy of the medicinal product as regards patients' health or public health and any risk of undesirable effects on the environment [DIR 2001/83/EC Art 1(28)].

Safety concern

An important identified risk, important potential risk or missing information.

It is noted that the ICH definition of safety concern is: an important identified risk, important potential risk or important missing information, i.e. includes the qualifier "important" in relation to missing information (see Annex IV, ICH-E2C(R2) Guideline). The ICH-E2E Guideline (see Annex IV) uses the terms safety issue and safety concern interchangeably with the same definition for safety concern as defined in the ICH-E2C(R2) Guideline.

See also Important identified risk and Important potential risk, Missing information

Serious adverse reaction

An adverse reaction which results in death, is life-threatening, requires in-patient hospitalisation or prolongation of existing hospitalisation, results in persistent or significant disability or incapacity, or is a congenital anomaly/birth defect [DIR 2001/83/EC Art 1(12)].

Life-threatening in this context refers to a reaction in which the patient was at risk of death at the time of the reaction; it does not refer to a reaction that hypothetically might have caused death if more severe (see Annex IV, ICH-E2D Guideline).

Medical and scientific judgement should be exercised in deciding whether other situations should be considered serious reactions, such as important medical events that might not be immediately life threatening or result in death or hospitalisation but might jeopardise the patient or might require intervention to prevent one of the other outcomes listed above. Examples of such events are intensive treatment in an emergency room or at home for allergic bronchospasm, blood dyscrasias or convulsions that do not result in hospitalisation or development of dependency or abuse (see Annex IV, ICH-E2D Guideline).

Any suspected transmission via a medicinal product of an infectious agent is also considered a serious adverse reaction.

See also Adverse reaction

Signal

Information arising from one or multiple sources, including observations and experiments, which suggests a new potentially causal association, or a new aspect of a known association between an intervention and an event or set of related events, either adverse or beneficial, that is judged to be of sufficient likelihood to justify verificatory action [IR 520/2012 Art 19(1)].

For the purpose of monitoring data in the EudraVigilance database, only signals related to an adverse reaction shall be considered [IR 520/2012, Art 19(1)].

For the purpose of Section 16.2 of the periodic benefit-risk evaluation report, signals relate to adverse effects (see Annex IV, ICH-E2C(R2) Guideline).

See also Validated signal, Newly identified signal, Closed signal, Ongoing signal, Signal management process, Adverse reaction

Signal management process

Includes the following activities: signal detection, signal validation, signal confirmation, signal analysis and prioritisation, signal assessment and recommendation for action [IR 520/2012 Art 21(1)].

It therefore is a set of activities performed to determine whether, based on an examination of individual case safety reports (ICSRs), aggregated data from active surveillance systems or studies, literature information or other data sources, there are new risks causally associated with an active substance or a medicinal product or whether known risks have changed.

See also Signal validation

Signal validation

Process of evaluating the data supporting a detected signal in order to verify that the available documentation contains sufficient evidence demonstrating the existence of a new potentially causal association, or a new aspect of a known association, and therefore justifies further analysis of the signal [IR 520/2012 Art 21(1)].

See also Validated signal

Solicited sources of individual case safety reports

Organised data collection systems, which include clinical trials, registries, post-authorisation named-patients use programmes, other patient support and disease management programmes, surveys of patients or healthcare providers or information gathering on efficacy or patient compliance. For the purpose of safety reporting, solicited reports should not be considered spontaneous but classified as individual case safety reports from studies and therefore should have an appropriate causality assessment by a healthcare professional or the marketing authorisation holder (see Annex IV, ICH-E2D).

See also Clinical trial, Post-authorisation safety study, Non-interventional trial

Spontaneous report, synonym: Spontaneous notification

An unsolicited communication by a healthcare professional or consumer to a company, regulatory authority or other organisation (e.g. the World Health Organization, a regional centre, a poison control centre) that describes one or more adverse reactions in a patient who was given one or more medicinal products and that does not derive from a study or any organised data collection scheme (see Annex IV, ICH-E2D).

In this context, an adverse reaction refers to a suspected adverse reaction.

Stimulated reporting can occur in certain situations, such as after a direct healthcare professional communication (DHPC), a publication in the press or questioning of healthcare professionals by company representatives, and adverse reaction reports arising from these situations are considered spontaneous reports (see Annex IV, ICH-E2D), provided the report meets the definition above. Reporting can also be stimulated by invitation from patients' or consumers' organisations to their members. Reporting made in the context of early post-marketing phase vigilance (EPPV), e.g. in Japan, is also considered stimulated reporting.

See also Adverse reaction

Stimulated reporting

See Spontaneous report

Substance

Any matter irrespective of origin which may be human (e.g. human blood and human blood products), animal (e.g. micro-organisms, whole animals, parts of organs, animal secretions, toxins, extracts, blood products), vegetable (e.g. micro-organisms, plants, part of plants, vegetable secretions, extracts), chemical (e.g. elements, naturally occurring chemical materials and chemical products obtained by chemical change or synthesis) [DIR 2001/83/EC Art 1(3)].

Summary of product characteristics (SmPC)

Part of the marketing authorisation of a medicinal product setting out the agreed position of the product as distilled during the course of the assessment process which includes the information described in Article 11 of Directive 2001/83/EC. It is the basis of information for healthcare professionals on how to use the product safely and effectively. The package leaflet is drawn in accordance with the summary of product characteristics (based on A Guideline on Summary of Product Characteristics, Volume 2C of the Rules Governing Medicinal Products in the EU).

Target population (treatment); synonym: Treatment target population

The patients who might be treated with the medicinal product in accordance with the indication(s) and contraindications in the authorised product information.

Target population (vaccine); synonym: Vaccine target population

Persons who might be vaccinated in accordance with the indication(s) and contraindications in the authorised product information and official recommendations for vaccinations.

Traditional herbal medicinal product

A herbal medicinal product that fulfils the conditions laid down in Article 16a(1) of Directive 2001/83/EC [DIR 2001/83/EC Art 1(29)], i.e.

(a) it has (an)indication(s) exclusively appropriate to traditional herbal medicinal products which, by virtue of their composition and purpose, are intended and designed for use without the supervision of a medical practitioner for diagnostic purposes or for prescription or monitoring of treatment;

(b) it is exclusively for administration in accordance with a specified strength and posology;

(c) it is an oral, external and/or inhalation preparation;

(d) the period of traditional use as laid down in Article 16c(1)(c) has elapsed;

(e) the data on the traditional use of the medicinal product are sufficient; in particular the product proves not to be harmful in the specified conditions of use and the pharmacological effects or efficacy of the medicinal product are plausible on the basis of long-standing use and experience [DIR 2001/83/EC Art 16a].

Regarding (d), the product must have been in medicinal use throughout a period of at least 30 years, including at least 15 years within the EU (see DIR 2001/83/EC Art 16c(c) and European Commission Questions & Answers Document on Registration of Traditional Herbal Medicinal Products, 2011).

See also Herbal medicinal product

Unexpected adverse reaction

An adverse reaction, the nature, severity or outcome of which is not consistent with the summary of product characteristics [DIR 2001/83/EC Art 1(13)]¹³.

This includes class-related reactions which are mentioned in the summary of product characteristics (SmPC) but which are not specifically described as occurring with this product. For products authorised nationally, the relevant SmPC is that authorised by the competent authority in the Member State to whom the reaction is being reported. For centrally authorised products, the relevant SmPC is the SmPC authorised by the European Commission. During the time period between a CHMP opinion in favour of granting a marketing authorisation and the Commission decision granting the marketing authorisation, the relevant SmPC is the SmPC annexed to the CHMP opinion.

See also Summary of product characteristics

Upper management

Group of persons in charge of the highest executive management of an organisation.

¹³ For investigational medicinal products, an unexpected adverse reaction is an adverse reaction, the nature or severity of which is not consistent with the applicable product information (e.g. the investigator's brochure for an unauthorised investigational product or the summary of product characteristics for an authorised product) [Dir 2001/20/EC Art 2(p)].

Membership of this group is determined by the governance structure of the organisation. While it is envisaged that the upper management usually is a group, the head of the organisation is the one person at the top of the organisation with ultimate responsibility for ensuring that the organisation complies with relevant legislation.

Vaccination

The administration of a vaccine with the aim to produce immune response.

See also Immunisation

Vaccination failure

Vaccination failure due to actual vaccine failure or failure to vaccinate (see CIOMS-WHO¹⁴).

Vaccination failure may be defined based on clinical endpoints or immunological criteria, where correlates or surrogate markers for disease protection exist. Primary failure (e.g. lack of seroconversion or seroprotection) needs to be distinguished from secondary failure (waning immunity) (see CIOMS-WHO¹⁴).

See also Vaccine failure, Failure to vaccinate

Vaccine

See Immunological medicinal product

Vaccine failure

Confirmed or suspected vaccine failure.

Confirmed clinical vaccine failure

Occurrence of the specific vaccine-preventable disease in a person who is appropriately and fully vaccinated taking into account the incubation period and the normal delay for the protection to be acquired as a result of immunisation (see CIOMS-WHO¹⁵).

Suspected clinical vaccine failure

Occurrence of disease in an appropriately and fully vaccinated person, but the disease is not confirmed to be the specific vaccine-preventable disease, e.g. disease of unknown serotype in a fully vaccinated person (based on CIOMS-WHO¹⁵).

Confirmed immunological vaccine failure

Failure of the vaccinated person to develop the accepted marker of protective immune response after being fully and appropriately vaccinated, as demonstrated by having tested or examined the vaccinated person at an appropriate time interval after completion of immunisation (based on CIOMS-WHO¹⁵).

Suspected immunological vaccine failure

Failure of the vaccinated person to develop the accepted marker of protective immune response after being fully and appropriately vaccinated, but with the testing or examination of the vaccinated person done at an inappropriate time interval after completion of immunisation (based on CIOMS-WHO¹⁵).

¹⁴ Council for International Organizations of Medical Sciences (CIOMS). Definition and application of terms of vaccine pharmacovigilance (report of CIOMS/WHO Working Group on Vaccine Pharmacovigilance). Genève: CIOMS; 2012.

¹⁵ Council for International Organizations of Medical Sciences (CIOMS). Definition and application of terms of vaccine pharmacovigilance (report of CIOMS/WHO Working Group on Vaccine Pharmacovigilance). Genève: CIOMS; 2012.

For interpreting what means appropriately vaccinated, consider the explanatory note for Immunisation error-related reaction.

See also Vaccination failure

Vaccine pharmacovigilance

The science and activities relating to the detection, assessment, understanding and communication of adverse events following immunisation and other vaccine- or immunisation-related issues, and to the prevention of untoward effects of the vaccine or immunisation (see CIOMS-WHO¹⁶).

In this definition, immunisation means the usage of a vaccine for the purpose of immunising individuals (see CIOMS-WHO¹⁶), which in the EU is preferably referred to as vaccination (in the report of CIOMS/WHO Working Group on Vaccine Pharmacovigilance the terms immunisation and vaccination are used interchangeably¹⁶). Usage includes all processes that occur after a vaccine product has left the manufacturing/packaging site, i.e. handling, prescribing and administration of the vaccine (see CIOMS-WHO¹⁶).

An adverse event following immunisation (AEFI) is any untoward medical occurrence which follows immunisation and which does not necessarily have a causal relationship with the usage of the vaccine. The adverse event may be any unfavourable or unintended sign, abnormal laboratory finding, symptom or disease. While this AEFI definition is compatible with the definition of adverse event applied in the EU, the AEFI definition is not needed to describe pharmacovigilance for vaccines in the EU. However, EU guidance on pharmacovigilance for vaccines makes use of the terminology suggested by CIOMS-WHO¹⁶ regarding possible causes of adverse events, turning them into suspected adverse reactions. A coincidental event is an AEFI that is caused by something other than the vaccine product, immunisation error or immunisation anxiety (see CIOMS-WHO¹⁶).

See also Adverse event, Immunisation anxiety-related reaction, Immunisation error-related reaction, Vaccine product-related reaction, Vaccine quality defect-related reaction, Vaccination

Vaccine product-related reaction

An adverse event following immunisation that is caused or precipitated by a vaccine due to one or more of the inherent properties of the vaccine product (see CIOMS-WHO¹⁷).

In this definition immunisation means the usage (handling, prescribing and administration) of a vaccine for the purpose of immunising individuals (see CIOMS-WHO¹⁷), which in the EU is preferably referred to as vaccination (in the report of CIOMS/WHO Working Group on Vaccine Pharmacovigilance the terms immunisation and vaccination are used interchangeably¹⁷).

See also Adverse reaction, Vaccine pharmacovigilance

Vaccine quality defect-related reaction

An adverse event following immunisation that is caused or precipitated by a vaccine that is due to one or more quality defects of the vaccine product including its administration device as provided by the manufacturer (see CIOMS-WHO¹⁸).

In this definition immunisation means the usage (handling, prescribing and administration) of a vaccine for the purpose of immunising individuals (see CIOMS-WHO¹⁸), which in the EU is preferably referred to as vaccination (in

¹⁶ Council for International Organizations of Medical Sciences (CIOMS). Definition and application of terms of vaccine pharmacovigilance (report of CIOMS/WHO Working Group on Vaccine Pharmacovigilance). Genève: CIOMS; 2012.

¹⁷ Council for International Organizations of Medical Sciences (CIOMS). Definition and application of terms of vaccine pharmacovigilance (report of CIOMS/WHO Working Group on Vaccine Pharmacovigilance). Genève: CIOMS; 2012.

¹⁸ Council for International Organizations of Medical Sciences (CIOMS). Definition and application of terms of vaccine pharmacovigilance (report of CIOMS/WHO Working Group on Vaccine Pharmacovigilance). Genève: CIOMS; 2012.

the report of CIOMS/WHO Working Group on Vaccine Pharmacovigilance the terms immunisation and vaccination are used interchangeably¹⁸).

For the purpose of this definition, a vaccine quality defect is defined as any deviation of the vaccine product as manufactured from its set quality specifications (see CIOMS-WHO¹⁸).

See also Adverse reaction, Vaccine pharmacovigilance

Valid individual case safety report

See Individual case safety report

Validated signal

A signal where the signal validation process of evaluating the data supporting the detected signal has verified that the available documentation contains sufficient evidence demonstrating the existence of a new potentially causal association, or a new aspect of a known association, and therefore justifies further analysis of the signal [based on IR 520/2012 Art 21(1)].

See also Signal