



EUROPEAN MEDICINES AGENCY
SCIENCE MEDICINES HEALTH

10 December 2020
EMA/CHMP/660100/2020

Summary of opinion¹ (post authorisation)

Iscover clopidogrel

On 10 December 2020, the Committee for Medicinal Products for Human Use (CHMP) adopted a positive opinion recommending a change to the terms of the marketing authorisation for the medicinal product Iscover. The marketing authorisation holder for this medicinal product is sanofi-aventis groupe.

The CHMP adopted a new indication as follows:

In patients with moderate to high-risk Transient Ischemic Attack (TIA) or minor Ischemic Stroke (IS)

Clopidogrel in combination with ASA is indicated in:

- Adult patients with moderate to high-risk TIA (ABCD2 score ≥ 4) or minor IS (NIHSS ≤ 3) within 24 hours of either the TIA or IS event.

For information, the full indications for Iscover will now be:

Secondary prevention of atherothrombotic events

Clopidogrel is indicated in:

- Adult patients suffering from myocardial infarction (from a few days until less than 35 days), ischemic stroke (from 7 days until less than 6 months) or established peripheral arterial disease.
- Adult patients suffering from acute coronary syndrome:
 - Non-ST segment elevation acute coronary syndrome (unstable angina or non-Q-wave myocardial infarction), including patients undergoing a stent placement following percutaneous coronary intervention, in combination with acetylsalicylic acid (ASA).
 - ST segment elevation acute myocardial infarction, in combination with ASA in medically treated patients eligible for thrombolytic therapy.

¹ Summaries of positive opinion are published without prejudice to the Commission decision, which will normally be issued 67 days from adoption of the opinion



In patients with moderate to high-risk Transient Ischemic Attack (TIA) or minor Ischemic Stroke (IS)

Clopidogrel in combination with ASA is indicated in:

- Adult patients with moderate to high-risk TIA (ABCD2 score ≥ 4) or minor IS (NIHSS ≤ 3) within 24 hours of either the TIA or IS event.

Prevention of atherothrombotic and thromboembolic events in atrial fibrillation

In adult patients with atrial fibrillation who have at least one risk factor for vascular events, are not suitable for treatment with Vitamin K antagonists (VKA) and who have a low bleeding risk, clopidogrel is indicated in combination with ASA for the prevention of atherothrombotic and thromboembolic events, including stroke.

Detailed recommendations for the use of this product will be described in the updated summary of product characteristics (SmPC), which will be published in the revised European public assessment report (EPAR), and will be available in all official European Union languages after a decision on this change to the marketing authorisation has been granted by the European Commission.