



EUROPEAN MEDICINES AGENCY
SCIENCE MEDICINES HEALTH

25 April 2013
EMA/CHMP/236919/2013
Committee for Medicinal Products for Human Use (CHMP)

Summary of opinion¹ (post authorisation)

RoActemra tocilizumab

On 25 April 2013, the Committee for Medicinal Products for Human Use (CHMP) adopted a positive opinion recommending a variation to the terms of the marketing authorisation for the medicinal product RoActemra. The marketing authorisation holder for this medicinal product is Roche Registration Limited. They may request a re-examination of the CHMP opinion, provided that they notify the European Medicines Agency in writing of their intention within 15 days of receipt of the opinion.

The CHMP adopted a new indication as follows:

RoActemra in combination with methotrexate (MTX) is indicated for the treatment of juvenile idiopathic polyarthritis (rheumatoid factor positive or negative and extended oligoarthritis) in patients 2 years of age and older, who have responded inadequately to previous therapy with MTX. RoActemra can be given as monotherapy in case of intolerance to MTX or where continued treatment with MTX is inappropriate.

Detailed conditions for the use of this product will be described in the updated summary of product characteristics (SmPC), which will be published in the revised European public assessment report (EPAR), and will be available in all official European Union languages after the variation to the marketing authorisation has been granted by the European Commission.

For information, the full indications for RoActemra will be as follows²:

RoActemra, in combination with methotrexate (MTX), is indicated for the treatment of moderate to severe active rheumatoid arthritis (RA) in adult patients who have either responded inadequately to, or who were intolerant to, previous therapy with one or more disease-modifying anti-rheumatic drugs (DMARDs) or tumour necrosis factor (TNF) antagonists. In these patients, RoActemra can be given as monotherapy in case of intolerance to MTX or where continued treatment with MTX is inappropriate.

¹ Summaries of positive opinion are published without prejudice to the Commission decision, which will normally be issued within 44 days (Type II variations) and 67 days (Annex II applications) from adoption of the opinion.

² The text in bold represents the new or the amended indication.



RoActemra has been shown to reduce the rate of progression of joint damage as measured by X-ray and to improve physical function when given in combination with methotrexate.

RoActemra is indicated for the treatment of active systemic juvenile idiopathic arthritis (sJIA) in patients 2 years of age and older, who have responded inadequately to previous therapy with NSAIDs and systemic corticosteroids. RoActemra can be given as monotherapy (in case of intolerance to MTX or where treatment with MTX is inappropriate) or in combination with MTX.

RoActemra in combination with methotrexate (MTX) is indicated for the treatment of juvenile idiopathic polyarthritis (rheumatoid factor positive or negative and extended oligoarthritis) in patients 2 years of age and older, who have responded inadequately to previous therapy with MTX. RoActemra can be given as monotherapy in case of intolerance to MTX or where continued treatment with MTX is inappropriate.