

EU RISK MANAGEMENT PLAN (RMP) FOR
pembrolizumab

RMP version to be assessed as part of this application:

RMP Version number: 46.0

Data lock point for this RMP: 03-SEP-2024

Date of final sign off: 17-SEP-2025

Rationale for submitting an updated RMP:

Addition of a new indication for pembrolizumab as monotherapy for the treatment of resectable locally advanced head and neck squamous cell carcinoma as neoadjuvant treatment, continued as adjuvant treatment in combination with radiation therapy with or without concomitant cisplatin and then as monotherapy in adults whose tumours express PD-L1 with a CPS \geq 1.

Addition of pembrolizumab subcutaneous route of administration with a new pharmaceutical form (solution for injection) and two new strengths (395 mg and 790 mg).

Summary of significant changes in this RMP:

Addition of study KEYNOTE-689 and MK-3475A-D77 in Modules SIII, SVII, and SVIII; no changes to the risk profile in Modules SIII, SVII, and SVIII.

Removal of the fulfilled commitments for KEYNOTE-716 and KEYNOTE-158 from Part IV and Part VI.

Change of the due date for the KEYNOTE-204 final CSR from 4Q2025 to 4Q2026 in Part IV.

Other RMP versions under evaluation:

- **Version number: 50.1**

Submitted on: 27-AUG-2025

Procedure number: EMA/VR/0000293815

Details of the currently approved RMP:

- **Version number: 45.0**

Approved with procedure: EMEA/H/C/003820/II/0154

Date of approval (opinion date): 14-NOV-2024

Qualified Person for Pharmacovigilance (QPPV) name: Peter De Veene, MD

QPPV oversight declaration: The content of this RMP has been reviewed and approved by the marketing authorisation holder's QPPV. The electronic signature is available on file.

TABLE OF CONTENTS

TABLE OF CONTENTS.....	3
LIST OF TABLES.....	5
LIST OF ABBREVIATIONS.....	15
PART I: PRODUCT(S) OVERVIEW.....	19
PART II: SAFETY SPECIFICATION.....	28
PART II: MODULE SI - EPIDEMIOLOGY OF THE INDICATION(S) AND TARGET POPULATION(S).....	28
PART II: MODULE SII - NON-CLINICAL PART OF THE SAFETY SPECIFICATION.....	48
PART II: MODULE SIII - CLINICAL TRIAL EXPOSURE.....	50
PART II: MODULE SIV - POPULATIONS NOT STUDIED IN CLINICAL TRIALS.....	120
SIV.1 Exclusion Criteria in Pivotal Clinical Studies Within the Development Program.....	120
SIV.2 Limitations to Detect Adverse Reactions in Clinical Trial Development Program.....	123
SIV.3 Limitations in Respect to Populations Typically Under-represented in Clinical Trial Development Program.....	123
PART II: MODULE SV - POST-AUTHORISATION EXPERIENCE.....	125
SV.1 Post-Authorisation Exposure.....	125
SV.1.1 Method Used to Calculate Exposure.....	125
SV.1.2 Exposure.....	125
PART II: MODULE SVI - ADDITIONAL EU REQUIREMENTS FOR THE SAFETY SPECIFICATION.....	126
PART II: MODULE SVII - IDENTIFIED AND POTENTIAL RISKS.....	127
SVII.1 Identification of Safety Concerns in the Initial RMP Submission.....	127
SVII.1.1 Risks Not Considered Important for Inclusion in the List of Safety Concerns in the RMP.....	127
SVII.1.2 Risks Considered Important for Inclusion in the List of Safety Concerns in the RMP.....	127
SVII.2 New Safety Concerns and Reclassification With a Submission of an Updated RMP.....	127
SVII.3 Details of Important Identified Risks, Important Potential Risks, and Missing Information.....	127

SVII.3.1	Presentation of Important Identified Risks and Important Potential Risks	127
SVII.3.2	Presentation of the Missing Information.....	230
PART II:	MODULE SVIII - SUMMARY OF THE SAFETY CONCERNS	231
PART III:	PHARMACOVIGILANCE PLAN (INCLUDING POST-AUTHORISATION SAFETY STUDIES).....	232
III.1	Routine Pharmacovigilance Activities	232
III.2	Additional Pharmacovigilance Activities.....	233
III.3	Summary Table of Additional Pharmacovigilance Activities.....	234
PART IV:	PLANS FOR POST-AUTHORISATION EFFICACY STUDIES	235
PART V:	RISK MINIMISATION MEASURES (INCLUDING EVALUATION OF THE EFFECTIVENESS OF RISK MINIMISATION ACTIVITIES).....	237
V.1	Routine Risk Minimisation Measures	237
V.2	Additional Risk Minimisation Measures.....	238
V.3	Summary of Risk Minimisation Measures	239
PART VI:	SUMMARY OF THE RISK MANAGEMENT PLAN BY PRODUCT	241
I.	The Medicine and What it is Used For.....	241
II.	Risks Associated With the Medicine and Activities to Minimise or Further Characterise the Risks.....	249
II.A	List of Important Risks and Missing Information.....	249
II.B	Summary of Important Risks	250
II.C	Post-Authorisation Development Plan.....	254
II.C.1	Studies Which are Conditions of the Marketing Authorisation.....	254
II.C.2	Other Studies in Post-Authorisation Development Plan.....	254
REFERENCES	255
ANNEXES	283

LIST OF TABLES

Table I.1:	Product Overview	19
Table SI.1:	Background Frequencies of Relevant Adverse Events in Target Populations.....	45
Table SII.1:	Summary of Important Safety Findings from Non-clinical Studies.....	48
Table SIII.1:	Summary of Drug Exposure Melanoma Subjects KN001, KN002 and KN006 Monotherapy (APaT Population)	65
Table SIII.2:	Summary of Drug Exposure Melanoma Participants KN054 and KN716 Monotherapy (APaT Population).....	66
Table SIII.3:	Summary of Drug Exposure NSCLC Subjects KN001, KN010, KN024, and KN042 Monotherapy (APaT Population)	66
Table SIII.4:	Summary of Drug Exposure to Pembrolizumab NSCLC Subjects KN021 (Cohorts A, C and G), KN189 and KN407 Combination Therapy (APaT Population).....	67
Table SIII.5:	Summary of Drug Exposure NSCLC Participants KN091 Monotherapy (APaT Population).....	67
Table SIII.6:	Summary of Drug Exposure NSCLC Participants in KN671 Combination Therapy (APaT Population).....	68
Table SIII.7:	Summary of Drug Exposure NSCLC Participants in MK-3475A-D77 Combination Therapy (APaT Population).....	68
Table SIII.8:	Summary of Drug Exposure HL Subjects KN013, KN087, and KN204 Monotherapy (APaT Population).....	69
Table SIII.9:	Summary of Drug Exposure UC Subjects KN045 and KN052 Monotherapy (APaT Population).....	69
Table SIII.10:	Summary of Drug Exposure to Pembrolizumab Urothelial Cancer Participants in KNA39 and KN869 Pembrolizumab + EV Combination Therapy (APaT Population).....	70
Table SIII.11:	Summary of Drug Exposure HNSCC Subjects KN012 B+B2 Cohorts, KN040, KN055 and KN048 Monotherapy (ASaT Population).....	70
Table SIII.12:	Summary of Drug Exposure to Pembrolizumab HNSCC Subjects KN048 Combination Therapy (ASaT Population).....	71
Table SIII.13:	Summary of Drug Exposure to Pembrolizumab HNSCC Participants in KN689 Monotherapy or Combination Therapy (APaT Population)	71
Table SIII.14:	Summary of Drug Exposure to Pembrolizumab RCC Subjects KN426 Combination Therapy (ASaT Population)	72

Table SIII.15:	Summary of Drug Exposure to Pembrolizumab RCC Participants KN581 Combination Therapy (APaT Population)	72
Table SIII.16:	Summary of Drug Exposure RCC Participants KN564 Monotherapy (APaT Population).....	73
Table SIII.17:	Summary of Drug Exposure CRC Subjects KN177 Monotherapy (ASaT Population).....	73
Table SIII.18:	Summary of Drug Exposure to Pembrolizumab Esophageal Cancer Subjects KN590 Combination Therapy (ASaT Population).....	74
Table SIII.19:	Summary of Drug Exposure to Pembrolizumab TNBC Subjects KN355 Combination Therapy (ASaT Population)	74
Table SIII.20:	Summary of Drug Exposure to Pembrolizumab TNBC Subjects KN522 Combination Therapy (APaT Population)	75
Table SIII.21:	Summary of Drug Exposure to Pembrolizumab Endometrial Carcinoma Participants KN146 and KN775 Combination Therapy (APaT Population).....	75
Table SIII.22:	Summary of Drug Exposure to Pembrolizumab Endometrial Carcinoma Participants Subjects KN868 Combination Therapy (APaT Population)	76
Table SIII.23:	Summary of Drug Exposure MSI-H Subjects KN158 Cohort K and KN164 Cohorts A and B Monotherapy (ASaT Population)	76
Table SIII.24:	Summary of Drug Exposure to Pembrolizumab Cervical Cancer Participants in KNA18 + KN826 Combination Therapy (APaT Population)	77
Table SIII.25:	Summary of Drug Exposure Gastric Cancer Participants in KN811 Combination Therapy (APaT Population)	77
Table SIII.26:	Summary of Drug Exposure to Pembrolizumab Gastric Cancer Participants in KN859 Combination Therapy (APaT Population).....	78
Table SIII.27:	Summary of Drug Exposure to Pembrolizumab BTC Participants in KN966 Combination Therapy (APaT Population).....	78
Table SIII.28:	Summary of Drug Exposure to Pembrolizumab Mesothelioma Participants in KN483 and KNA17 Combination Therapy (APaT Population)	79
Table SIII. 29:	Clinical Trial Exposure by Dose and Duration Melanoma Subjects KN001, KN002 and KN006 Monotherapy (APaT Population).....	81

Table SIII.30:	Clinical Trial Exposure by Dose and Duration Melanoma Participants in KN054 and KN716 Monotherapy (APaT Population).....	82
Table SIII.31:	Clinical Trial Exposure by Dose and Duration NSCLC Subjects KN001, KN010, KN024 and KN042 Monotherapy (APaT Population).....	82
Table SIII.32:	Clinical Trial Exposure to Pembrolizumab by Dose and Duration NSCLC Subjects KN021 (Cohorts A, C and G), KN189 and KN407 Combination Therapy (APaT Population).....	83
Table SIII.33:	Clinical Trial Exposure by Dose and Duration NSCLC Participants KN091 Monotherapy (APaT Population).....	83
Table SIII. 34:	Clinical Trial Exposure by Dose and Duration NSCLC Participants in KN671 Combination Therapy (APaT Population).....	84
Table SIII.35:	Clinical Trial Exposure by Dose and Duration NSCLC Participants in MK-3475A-D77 Combination Therapy (APaT Population).....	84
Table SIII.36:	Clinical Trial Exposure by Dose and Duration HL Subjects in KN013, KN087, and KN204 Monotherapy (APaT Population).....	85
Table SIII.37:	Clinical Trial Exposure by Dose and Duration UC Subjects KN045 and KN052 Monotherapy (APaT Population).....	85
Table SIII.38:	Clinical Trial Exposure to Pembrolizumab by Dose and Duration Urothelial Cancer Participants in KNA39 and KN869 Pembrolizumab + EV Combination Therapy (APaT Population).....	85
Table SIII.39:	Clinical Trial Exposure by Dose and Duration HNSCC Subjects KN012 B+B2 Cohorts, KN040, KN055 and KN048 Monotherapy (ASaT Population).....	86
Table SIII.40:	Clinical Trial Exposure to Pembrolizumab by Dose and Duration HNSCC Subjects in KN048 Combination Therapy (ASaT Population).....	86
Table SIII.41:	Clinical Trial Exposure to Pembrolizumab by Dose and Duration HNSCC Participants in KN689 Monotherapy or Combination Therapy (APaT Population).....	86
Table SIII.42:	Clinical Trial Exposure to Pembrolizumab by Dose and Duration RCC Subjects KN426 Combination Therapy (ASaT Population).....	87
Table SIII.43:	Clinical Trial Exposure to Pembrolizumab by Dose and Duration RCC Participants KN581 Combination Therapy (APaT Population).....	87

Table SIII.44:	Clinical Trial Exposure by Dose and Duration RCC Participants KN564 Monotherapy (APaT Population).....	88
Table SIII.45:	Clinical Trial Exposure by Dose and Duration CRC Subjects KN177 Monotherapy (ASaT Population).....	88
Table SIII.46:	Clinical Trial Exposure to Pembrolizumab by Dose and Duration Esophageal Cancer Subjects KN590 Combination Therapy (ASaT Population).....	89
Table SIII.47:	Clinical Trial Exposure to Pembrolizumab by Dose and Duration TNBC Subjects KN355 Combination Therapy (ASaT Population).....	89
Table SIII.48:	Clinical Trial Exposure to Pembrolizumab by Dose and Duration TNBC Subjects KN522 Combination Therapy (APaT Population).....	90
Table SIII.49:	Clinical Trial Exposure to Pembrolizumab by Dose and Duration Endometrial Carcinoma Participants KN146 and KN775 Combination Therapy (APaT Population)	90
Table SIII.50:	Clinical Trial Exposure to Pembrolizumab by Dose and Duration Endometrial Carcinoma Participants in KN868 Combination Therapy (APaT Population).....	91
Table SIII.51:	Clinical Trial Exposure by Dose and Duration MSI-H Subjects KN158 Cohort K and KN164 Cohorts A and B Monotherapy (ASaT Population)	91
Table SIII.52:	Clinical Trial Exposure to Pembrolizumab by Dose and Duration Cervical Cancer Participants in KNA18 + KN826 Combination Therapy (APaT Population).....	92
Table SIII.53:	Clinical Trial Exposure to Pembrolizumab by Dose and Duration Gastric Cancer Participants KN811 Combination Therapy (APaT Population).....	92
Table SIII.54:	Clinical Trial Exposure to Pembrolizumab by Dose and Duration Gastric Cancer Participants KN859 Combination Therapy (APaT Population).....	93
Table SIII.55:	Clinical Trial Exposure to Pembrolizumab by Dose and Duration BTC Participants in KN966 Combination Therapy (APaT Population)	93
Table SIII.56:	Clinical Trial Exposure to Pembrolizumab by Dose and Duration Mesothelioma Participants in KN483 and KNA17 Combination Therapy (APaT Population).....	94
Table SIII.57:	Clinical Trial Exposure by Age and Gender Melanoma Subjects KN001, KN002 and KN006 Monotherapy (APaT Population).....	98

Table SIII.58:	Clinical Trial Exposure by Age and Gender Melanoma Participants KN054 and KN716 Monotherapy (APaT Population).....	98
Table SIII.59:	Clinical Trial Exposure by Age and Gender NSCLC Subjects KN001, KN010, KN024 and KN042 Monotherapy (APaT Population).....	98
Table SIII.60:	Clinical Trial Exposure to Pembrolizumab by Age and Gender NSCLC Subjects KN021 (Cohorts A, C and G), KN189 and KN407 Combination Therapy (APaT Population)	99
Table SIII.61:	Clinical Trial Exposure by Age and Gender NSCLC Participants KN091 Monotherapy (APaT Population).....	99
Table SIII.62:	Clinical Trial Exposure to Pembrolizumab by Age and Gender NSCLC Participants in KN671 Combination Therapy (APaT Population).....	99
Table SIII.63:	Clinical Trial Exposure to Pembrolizumab SC by Age and Gender NSCLC Participants in MK-3475A-D77 Combination Therapy (APaT Population).....	100
Table SIII.64:	Clinical Trial Exposure by Age and Gender HL Subjects KN013, KN087, and KN204 Monotherapy (APaT Population).....	100
Table SIII.65:	Clinical Trial Exposure by Age and Gender KN051 Parts I and II Monotherapy (APaT Population).....	100
Table SIII.66:	Clinical Trial Exposure by Age and Gender UC Subjects KN045 and KN052 Monotherapy (APaT Population)	101
Table SIII.67:	Clinical Trial Exposure to Pembrolizumab by Age and Gender UC Participants KNA39 and KN869 Pembrolizumab + EV Combination Therapy (APaT Population).....	101
Table SIII.68:	Clinical Trial Exposure by Age and Gender HNSCC Subjects KN012 B+B2 Cohorts, KN040, KN055 and KN048 Monotherapy (ASaT Population).....	101
Table SIII.69:	Clinical Trial Exposure to Pembrolizumab by Age and Gender HNSCC Subjects in KN048 Combination Therapy (ASaT Population).....	102
Table SIII.70:	Clinical Trial Exposure to Pembrolizumab by Age and Gender HNSCC Subjects in KN689 Monotherapy or Combination Therapy (APaT Population).....	102
Table SIII.71:	Clinical Trial Exposure to Pembrolizumab by Age and Gender RCC Subjects KN426 Combination Therapy (ASaT Population).....	102

Table SIII.72:	Clinical Trial Exposure to Pembrolizumab by Age and Gender RCC Participants KN581 Combination Therapy (APaT Population).....	102
Table SIII.73:	Clinical Trial Exposure by Age and Gender RCC Participants KN564 Monotherapy (APaT Population)	103
Table SIII.74:	Clinical Trial Exposure by Age and Gender CRC Subjects in KN177 Monotherapy (ASaT Population).....	103
Table SIII.75:	Clinical Trial Exposure to Pembrolizumab by Age and Gender Esophageal Cancer Subjects KN590 Combination Therapy (ASaT Population)	103
Table SIII.76:	Clinical Trial Exposure to Pembrolizumab by Age and Gender TNBC Subjects KN355 Combination Therapy (ASaT Population).....	103
Table SIII.77:	Clinical Trial Exposure to Pembrolizumab by Age and Gender TNBC Subjects KN522 Combination Therapy (APaT Population).....	104
Table SIII.78:	Clinical Trial Exposure to Pembrolizumab by Age Endometrial Carcinoma Participants KN146 and KN775 Combination Therapy (APaT Population).....	104
Table SIII.79:	Clinical Trial Exposure to Pembrolizumab by Age Endometrial Carcinoma Participants KN868 Combination Therapy (APaT Population).....	104
Table SIII.80:	Clinical Trial Exposure by Age and Gender MSI-H Subjects KN158 Cohort K and KN164 Cohorts A and B Monotherapy (ASaT Population)	105
Table SIII.81:	Clinical Trial Exposure to Pembrolizumab by Age Cervical Cancer Participants in KNA18 and KN826 Combination Therapy (APaT Population).....	105
Table SIII.82:	Clinical Trial Exposure to Pembrolizumab by Age and Gender Gastric Cancer Participants KN811 Combination Therapy (APaT Population)	105
Table SIII.83:	Clinical Trial Exposure to Pembrolizumab by Age and Gender Gastric Cancer Participants KN859 Combination Therapy (APaT Population)	106
Table SIII.84:	Clinical Trial Exposure to Pembrolizumab by Age and Gender BTC Participants in KN966 Combination Therapy (APaT Population).....	106
Table SIII.85:	Clinical Trial Exposure to Pembrolizumab by Age and Gender Mesothelioma Participants in KN483 and KNA17 Combination Therapy (APaT Population).....	106

Table SIII.86:	Clinical Trial Exposure by Race Melanoma Subjects KN001, KN002 and KN006 Monotherapy (APaT Population)	110
Table SIII.87:	Clinical Trial Exposure by Race Melanoma Participants KN054 and KN716 Monotherapy (APaT Population)	110
Table SIII.88:	Clinical Trial Exposure by Race NSCLC Subjects KN001, KN010, KN024 and KN042 Monotherapy (APaT Population).....	111
Table SIII.89:	Clinical Trial Exposure to Pembrolizumab by Race NSCLC Subjects KN021 (Cohorts A, C and G), KN189 and KN407 Combination Therapy (APaT Population).....	111
Table SIII.90:	Clinical Trial Exposure to Pembrolizumab by Race NSCLC Participants KN091 Monotherapy Therapy (APaT Population).....	111
Table SIII.91:	Clinical Trial Exposure to Pembrolizumab by Race NSCLC Participants in KN671 Combination Therapy (APaT Population).....	112
Table SIII.92:	Clinical Trial Exposure to Pembrolizumab SC by Race NSCLC Participants in MK-3475A-D77 Combination Therapy (APaT Population).....	112
Table SIII.93:	Clinical Trial Exposure by Race HL Subjects KN013, KN087, and KN204 Monotherapy (APaT Population)	112
Table SIII.94:	Clinical Trial Exposure by Race UC Subjects KN045 and KN052 Monotherapy (APaT Population).....	113
Table SIII.95:	Clinical Trial Exposure by Race UC Participants KNA39 and KN869 Pembrolizumab + EV Combination Therapy (APaT Population).....	113
Table SIII.96:	Clinical Trial Exposure by Race HNSCC Subjects KN012 B+B2 Cohorts, KN040, KN055 and KN048 Monotherapy (ASaT Population)	113
Table SIII.97:	Clinical Trial Exposure to Pembrolizumab by Race HNSCC Subjects in KN048 Combination Therapy (ASaT Population)	114
Table SIII.98:	Clinical Trial Exposure to Pembrolizumab by Race HNSCC Participants in KN689 Monotherapy or Combination Therapy (APaT Population)	114
Table SIII.99:	Clinical Trial Exposure to Pembrolizumab by Race RCC Subjects KN426 Combination Therapy (ASaT Population).....	114
Table SIII.100:	Clinical Trial Exposure to Pembrolizumab by Race RCC Participants KN581 Combination Therapy (APaT).....	115
Table SIII.101:	Clinical Trial Exposure by Race RCC Participants KN564 Monotherapy (APaT Population).....	115
Table SIII.102:	Clinical Trial Exposure by Race CRC Subjects KN177 Monotherapy (ASaT Population).....	115

Table SIII.103:	Clinical Trial Exposure to Pembrolizumab by Race Esophageal Cancer Subjects KN590 Combination Therapy (ASaT Population).....	116
Table SIII.104:	Clinical Trial Exposure to Pembrolizumab by Race TNBC Subjects in KN355 Combination Therapy (ASaT Population)	116
Table SIII.105:	Clinical Trial Exposure to Pembrolizumab by Race TNBC Subjects in KN522 Combination Therapy (APaT Population)	116
Table SIII.106:	Clinical Trial Exposure to Pembrolizumab by Race Endometrial Carcinoma Participants KN146 and KN775 Combination Therapy (APaT Population).....	117
Table SIII.107:	Clinical Trial Exposure to Pembrolizumab by Race Endometrial Carcinoma Participants in KN868 Combination Therapy (APaT)	117
Table SIII.108:	Clinical Trial Exposure by Race MSI-H Subjects in KN158 Cohort K and KN164 Cohorts A and B Monotherapy (ASaT Population).....	118
Table SIII.109:	Clinical Trial Exposure to Pembrolizumab by Race Cervical Cancer Participants in KNA18 + KN826 Combination Therapy (APaT Population)	118
Table SIII.110:	Clinical Trial Exposure to Pembrolizumab by Race in Gastric Cancer Participants KN811 Combination Therapy (APaT Population).....	118
Table SIII.111:	Clinical Trial Exposure to Pembrolizumab by Race in Gastric Cancer Participants KN859 Combination Therapy (APaT Population).....	119
Table SIII.112:	Clinical Trial Exposure to Pembrolizumab by Race BTC Participants in KN966 Combination Therapy (APaT Population).....	119
Table SIII.113:	Clinical Trial Exposure to Pembrolizumab by Race Mesothelioma Participants in KN483 and KNA17 Combination Therapy (APaT Population).....	119
Table SIV.1.1:	Exclusion Criteria in Pivotal Clinical Studies Within the Development Program	120
Table SIV.3.1:	Exposure of Special Populations Included or not in Clinical Trial Development Programs.....	124
Table SVII.3.1.1:	Details of Important Identified Risk: Immune-Mediated Adverse Reactions-Pneumonitis	135
Table SVII.3.1.2:	Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Colitis	146

Table SVII.3.1.3:	Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Hepatitis.....	156
Table SVII.3.1.4:	Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Nephritis	166
Table SVII.3.1.5:	Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies – Hypophysitis (Including Hypopituitarism)	175
Table SVII.3.1.6:	Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies –Adrenal Insufficiency (Primary and Secondary)	184
Table SVII.3.1.7:	Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis).....	195
Table SVII.3.1.8:	Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies -Type 1 Diabetes Mellitus	220
Table SVII.3.1.9:	Details of Important Potential Risk: For Hematologic Malignancies: Increased Risk of Severe Complications of Allogeneic Stem Cell Transplantation (SCT) in Patients Who Have Previously Received Pembrolizumab.....	229
Table SVII.3.1.10:	Details of Important Potential Risk: Graft Versus Host Disease (GVHD) After Pembrolizumab Administration in Patients With a History of Allogeneic Stem Cell Transplant (SCT).....	230
Table SVIII.1:	Summary of Safety Concerns	231
Table III.1.1:	Overview of Pharmacovigilance Actions	233
Table IV.1:	Planned and On-Going Post-Authorisation Efficacy Studies that are Conditions of the Marketing Authorisation or that are Specific Obligations.....	235
Table V.1.1:	Description of Routine Risk Minimisation Measures by Safety Concern	237
Table V.2.1:	Description of Additional Risk Minimisation Measures by Safety Concern.....	238
Table V.3.1:	Summary Table of Pharmacovigilance Activities and Risk Minimisation Activities by Safety Concern.....	239
Table II.A.1:	List of Important Risks and Missing Information.....	250
Table II.B.1:	Important Identified Risk: Immune-Mediated Adverse Reactions.....	250

Table II.B.2:	Important Potential Risk: For Hematologic Malignancies: Increased Risk of Severe Complications of Allogeneic Stem Cell Transplantation (SCT) in Patients Who Have Previously Received Pembrolizumab	253
Table II.B.3:	Important Potential Risk: Graft Versus Host Disease (GVHD) After Pembrolizumab Administration in Patients With a History of Allogeneic Stem Cell Transplant (SCT).....	253
Table II.C.1.1:	Studies Which are Conditions of the Marketing Authorisation.....	254

LIST OF ABBREVIATIONS

5-FU	5-Fluorouracil
ADR	Adverse Drug Reaction
AE	Adverse Event
AEOSI	Adverse Events of Special Interest
AJCC	American Joint Committee on Cancer
ALK	Anaplastic Lymphoma Kinase
ALT	Alanine Aminotransferase
APaT	All Patients as Treated
ASaT	All Subjects as Treated
ASCT	Autologous Stem Cell Transplant
ASE	Aggregate Safety Evaluation
AST	Aspartate Aminotransferase
ATC	Anatomical Therapeutic Chemical classification system
AUC	Area Under the Curve
BC	Breast Cancer
BICR	Blinded Independent Central Review
BRAF	B-Raf Proto-Oncogene
BRCA	Breast Cancer Gene
BTC	Biliary Tract Carcinoma
BV	Brentuximab vedotin
BW	Body weight
CCA	Cholangiocarcinoma
cHL	classical Hodgkin Lymphoma
CNS	Central Nervous System
COPD	Chronic Obstructive Pulmonary Disease
CPS	Combined Positive Score
CR	Complete response
CRC	Colorectal Cancer
CT	Clinical Trial
CTCAE	Common Terminology Criteria for Adverse Events
CTLA-4	Cytotoxic T-Lymphocyte-Associated Antigen 4
CVD	Cardiovascular Disease
DBL	Database Lock
DLT	Dose Limiting Toxicity
dMMR	Mismatch Repair Deficient
EBRT	External Beam Radiotherapy

EBV	Epstein-Barr Virus
EC	Endometrial carcinoma
EEA	European Economic Area
EGFR	Epidermal Growth Factor Receptor
EMA	European Medicines Agency
EORTC	European Organisation for Research and Treatment of Cancer
EPAR	European Public Assessment Report
EU	European Union
EV	Enfortumab vedotin
FIGO	International Federation of Gynecology and Obstetrics
FDA	Food and Drug Administration
GC	Gastric Cancer
GEJ	Gastro-esophageal Junction
GVHD	Graft Versus Host Disease
HBV	Hepatitis B Virus
HCV	Hepatitis C Virus
HCP	Health Care Provider
HER2	Human Epithelial Growth Factor Receptor 2
HGB	Hemoglobin
HIV	Human Immunodeficiency Virus
HL	Hodgkin Lymphoma
HNC	Head and Neck Cancer
HNSCC	Head and Neck Squamous Cell Carcinoma
HPV	Human Papillomavirus
HSCT	Hematopoietic Stem Cell Transplant
IgG4	Immunoglobulin G4
IM	Intramuscular(ly)
INN	International Nonproprietary Name
IPI	Ipilimumab
IV	Intravenous(ly)
KG	Kilogram
KN	KEYNOTE
LACC	Locally advanced cervical cancer
la/mUC	Locally advanced or metastatic urothelial cancer
mAb	Humanized monoclonal antibody
MAH	Marketing Authorisation Holder
MedDRA	Medical Dictionary for Regulatory Activities
MEK	Mitogen-activated protein kinase kinase

MG	Milligram
MIBC	Muscle-invasive bladder cancer
ML	Milliliter
MPM	Malignant pleural mesothelioma
MNP	Malignant neoplasm progression
MSI-H	Microsatellite Instability-High
MSS	Microsatellite Stable
mTOR	Mammalian Target of Rapamycin
N/A	Not Applicable
NAC	Neoadjuvant Chemotherapy
NAFLD	Nonalcoholic fatty liver disease
NASH	Nonalcoholic steatohepatitis
NCCN	National Comprehensive Cancer Network
NCI	National Cancer Institute
NED	No Evidence of Disease
NHL	Non-Hodgkin Lymphoma
NLP	Nodular lymphocyte predominant
NLPHL	Nodular lymphocyte predominant Hodgkin's lymphoma
NOAEL	No observed adverse effect level
NSCLC	Non-Small Cell Lung Cancer
ORR	Objective Response Rate
OS	Overall Survival
PAES	Post-Authorisation Efficacy Studies
pCR	Pathological Complete Response
PD	Pharmacodynamics
PD-1	Programmed Cell Death-1
PD-L1	Programmed Cell Death 1 Ligand
PD-L2	Programmed Cell Death 2 Ligand
PFS	Progression Free Survival
PK	Pharmacokinetics
PIP	Pediatric Investigation Plans
PM	Postmarketing
PR	Partial Response
PRAC	Pharmacovigilance Risk Assessment Committee
PSUR	Periodic Safety Update Report
QW	Every week
Q2W	Every 2 weeks
Q3W	Every 3 weeks

Q6W	Every 6 weeks
QPPV	Qualified Person for Pharmacovigilance
RC+PLND	Radical cystectomy and Pelvic lymph node dissection
RCC	Renal Cell Carcinoma
RECIST	Response Evaluation Criteria in Solid Tumors
R/M	Recurrent/Metastatic
RMP	Risk Management Plan
SAE	Serious Adverse Event
SC	Subcutaneous
SCT	Stem Cell Transplantation
SOC	Standard of Care
SmPC	Summary of Product Characteristics
SMQ	Standardized MedDRA Query
TDAR	T-Dependent Antibody Response
TKI	Tyrosine Kinase Inhibitor
TMB-H	Tumor Mutational Burden-High
TNBC	Triple-Negative Breast Cancer
TPC	Treatment of Physician's Choice
TPS	Tumor Proportion Score
U	Unit
UC	Urothelial Carcinoma
ULN	Upper Limit of Normal
US	United States
VEGF	Vascular Endothelial Growth Factor
VEGFR2	Vascular Endothelial Growth Factor Receptor 2
VOD	Veno-occlusive Disease
WFRS	Worldwide Financial Reporting System

PART I: PRODUCT(S) OVERVIEW

Table I.1: Product Overview

Active substance(s) (International Nonproprietary Name (INN) or common name)	Pembrolizumab
Pharmacotherapeutic group(s) (Anatomical Therapeutic Chemical classification system (ATC) Code)	L01FF02
Marketing Authorisation	Merck Sharp & Dohme B.V. Waarderweg 39 2031 BN Haarlem The Netherlands
Number of medicinal products to which this RMP refers	1
Invented name(s) in the European Economic Area (EEA)	KEYTRUDA®
Marketing authorisation procedure	Centralised
Brief description of the product	<p>Chemical class: Humanized monoclonal anti PD-1 antibody</p> <p>Summary of mode of action: Pembrolizumab is a potent and highly-selective humanized monoclonal antibody (mAb) of the Immunoglobulin G4 (IgG4)/kappa isotype directed to the programmed cell death-1 (PD-1) receptor and designed to directly block the interaction between the receptor and its ligands, PD-L1 and PD-L2. The PD-1 pathway represents a major immune control switch which may be engaged by tumor cells to overcome active T-cell immune surveillance.</p> <p>Important information about its composition: Pembrolizumab for IV administration is presented as: - a sterile, preservative-free, clear to slightly opalescent, colourless to slightly yellow solution. Each vial of 4 mL contains 100 mg of pembrolizumab.</p> <p>Pembrolizumab for SC administration is presented as: - a sterile, preservative-free, clear to slightly opalescent, colourless to slightly yellow solution. Each vial of 2.4 mL contains 395 mg of pembrolizumab and each vial of 4.8 mL contains 790 mg of pembrolizumab.</p>
Hyperlink to the Prescribing Information	Refer to the proposed Product Information in module 1.3.1.
Indication(s) in the EEA	<p>Current:</p> <p>KEYTRUDA Administered IV:</p> <p><u>Melanoma</u> KEYTRUDA as monotherapy is indicated for the treatment of adults and adolescents aged 12 years and older with advanced (unresectable or metastatic) melanoma.</p> <p>KEYTRUDA as monotherapy is indicated for the adjuvant treatment of adults and adolescents aged 12 years and older with Stage IIB, IIC or III melanoma and who have undergone complete resection.</p>

Table I.1: Product Overview

	<p><u>Non-small cell lung carcinoma (NSCLC)</u> KEYTRUDA in combination with platinum-containing chemotherapy as neoadjuvant treatment, and then continued as monotherapy as adjuvant treatment, is indicated for the treatment of resectable non-small cell lung carcinoma at high risk of recurrence in adults.</p> <p>KEYTRUDA as monotherapy is indicated for the adjuvant treatment of adults with non-small cell lung carcinoma who are at high risk of recurrence following complete resection and platinum based chemotherapy.</p> <p>KEYTRUDA as monotherapy is indicated for the first-line treatment of metastatic non-small cell lung carcinoma (NSCLC) in adults whose tumours express PD-L1 with a $\geq 50\%$ tumour proportion score (TPS) with no EGFR or ALK positive tumour mutations.</p> <p>KEYTRUDA, in combination with pemetrexed and platinum chemotherapy, is indicated for the first-line treatment of metastatic non-squamous NSCLC in adults whose tumours have no EGFR or ALK positive mutations.</p> <p>KEYTRUDA in combination with carboplatin and either paclitaxel or nab-paclitaxel is indicated for the first-line treatment of metastatic squamous NSCLC in adults.</p> <p>KEYTRUDA as monotherapy is indicated for the treatment of locally advanced or metastatic NSCLC in adults whose tumours express PD-L1 with a $\geq 1\%$ TPS and who have received at least one prior chemotherapy regimen. Patients with EGFR or ALK positive tumour mutations should also have received targeted therapy before receiving KEYTRUDA.</p> <p><u>Malignant pleural mesothelioma (MPM)</u> KEYTRUDA, in combination with pemetrexed and platinum chemotherapy, is indicated for the first-line treatment of adults with unresectable non-epithelioid malignant pleural mesothelioma.</p> <p><u>Classical Hodgkin lymphoma (cHL)</u> KEYTRUDA as monotherapy is indicated for the treatment of adult and paediatric patients aged 3 years or older with relapsed or refractory classical Hodgkin Lymphoma (cHL) who have failed autologous stem cell transplant (ASCT) or following at least two prior therapies when ASCT is not a treatment option.</p> <p><u>Urothelial carcinoma</u> KEYTRUDA, in combination with enfortumab vedotin is indicated for the first-line treatment of unresectable or metastatic urothelial carcinoma in adults.</p> <p>KEYTRUDA as monotherapy is indicated for the treatment of locally advanced or metastatic urothelial carcinoma in adults who have received prior platinum-containing chemotherapy.</p> <p>KEYTRUDA as monotherapy is indicated for the treatment of locally advanced or metastatic urothelial carcinoma in adults who are not eligible for cisplatin-containing chemotherapy, and whose tumours express PD L1 with a combined positive score (CPS) ≥ 10.</p> <p><u>Head and neck squamous cell carcinoma (HNSCC)</u> KEYTRUDA as monotherapy is indicated for the treatment of resectable locally advanced head and neck squamous cell carcinoma (HNSCC) as neoadjuvant treatment, continued as adjuvant treatment in combination with radiation therapy with or without concomitant cisplatin and then as monotherapy in adults whose tumours express PD-L1 with a CPS ≥ 1.</p> <p>KEYTRUDA, as monotherapy or in combination with platinum and 5-fluorouracil (5-FU) chemotherapy, is indicated for the first-line treatment of metastatic or unresectable recurrent HNSCC in adults whose tumours express PD-L1 with a CPS ≥ 1.</p>
--	---

Table I.1: Product Overview

	<p>KEYTRUDA as monotherapy is indicated for the treatment of recurrent or metastatic HNSCC in adults whose tumours express PD-L1 with a $\geq 50\%$ TPS and progressing on or after platinum-containing chemotherapy.</p> <p>KEYTRUDA as monotherapy is indicated for the treatment of resectable locally advanced HNSCC as neoadjuvant treatment, continued as adjuvant treatment in combination with radiation therapy with or without platinum-containing chemotherapy and then as monotherapy in adults.</p> <p><u>Renal cell carcinoma (RCC)</u> KEYTRUDA, in combination with axitinib, is indicated for the first-line treatment of advanced renal cell carcinoma (RCC) in adults.</p> <p>KEYTRUDA, in combination with lenvatinib, is indicated for the first-line treatment of advanced renal cell carcinoma (RCC) in adults.</p> <p>KEYTRUDA as monotherapy is indicated for the adjuvant treatment of adults with renal cell carcinoma at increased risk of recurrence following nephrectomy, or following nephrectomy and resection of metastatic lesions.</p> <p><u>Microsatellite instability high (MSI-H) or mismatch repair deficient (dMMR) cancers</u></p> <p><u>Colorectal cancer (CRC)</u> KEYTRUDA as monotherapy is indicated for adults with microsatellite instability high (MSI-H) or mismatch repair deficient (dMMR) colorectal cancer in the following settings:</p> <ul style="list-style-type: none"> - first-line treatment of metastatic colorectal cancer; - treatment of unresectable or metastatic colorectal cancer after previous fluoropyrimidine-based combination therapy. <p><u>Non-colorectal cancers</u> KEYTRUDA as monotherapy is indicated for the treatment of the following MSI H or dMMR tumours in adults with:</p> <ul style="list-style-type: none"> - advanced or recurrent endometrial carcinoma, who have disease progression on or following prior treatment with a platinum containing therapy in any setting and who are not candidates for curative surgery or radiation; - unresectable or metastatic gastric, small intestine, or biliary cancer, who have disease progression on or following at least one prior therapy. <p><u>Oesophageal carcinoma</u> KEYTRUDA, in combination with platinum and fluoropyrimidine based chemotherapy, is indicated for the first-line treatment of patients with locally advanced unresectable or metastatic carcinoma of the oesophagus in adults whose tumours express PD-L1 with a CPS ≥ 10.</p> <p><u>Triple negative breast cancer (TNBC)</u> KEYTRUDA, in combination with chemotherapy as neoadjuvant treatment, and then continued as monotherapy as adjuvant treatment after surgery, is indicated for the treatment of adults with locally advanced, or early-stage triple-negative breast cancer (TNBC) at high risk of recurrence.</p> <p>KEYTRUDA, in combination with chemotherapy, is indicated for the treatment of locally recurrent unresectable or metastatic triple-negative breast cancer (TNBC) in adults whose tumors express PD-L1 with a CPS ≥ 10 and who have not received prior chemotherapy for metastatic disease.</p>
--	--

Table I.1: Product Overview

	<p><u>Endometrial carcinoma (EC)</u> KEYTRUDA, in combination with carboplatin and paclitaxel, is indicated for the first-line treatment of primary advanced or recurrent endometrial carcinoma in adults who are candidates for systemic therapy.</p> <p>KEYTRUDA, in combination with lenvatinib, is indicated for the treatment of advanced or recurrent endometrial carcinoma in adults who have disease progression on or following prior treatment with a platinum-containing therapy in any setting and who are not candidates for curative surgery or radiation.</p> <p><u>Cervical cancer</u> KEYTRUDA, in combination with chemoradiotherapy (external beam radiation therapy followed by brachytherapy), is indicated for the treatment of FIGO 2014 Stage III - IVA locally advanced cervical cancer in adults who have not received prior definitive therapy.</p> <p>KEYTRUDA, in combination with chemotherapy with or without bevacizumab, is indicated for the treatment of persistent, recurrent, or metastatic cervical cancer in adults whose tumours express PD-L1 with a CPS \geq 1.</p> <p><u>Gastric or gastro-oesophageal junction adenocarcinoma</u> KEYTRUDA, in combination with trastuzumab, fluoropyrimidine- and platinum-containing chemotherapy, is indicated for the first-line treatment of locally advanced unresectable or metastatic HER2-positive gastric or gastro-oesophageal junction adenocarcinoma in adults whose tumours express PD-L1 with a CPS \geq 1.</p> <p>KEYTRUDA, in combination with fluoropyrimidine and platinum -containing chemotherapy, is indicated for the first-line treatment of locally advanced unresectable or metastatic HER2-negative gastric or gastro-oesophageal junction adenocarcinoma in adults whose tumours express PD-L1 with a CPS \geq 1.</p> <p><u>Biliary tract carcinoma (BTC)</u> KEYTRUDA, in combination with gemcitabine and cisplatin, is indicated for the first-line treatment of locally advanced unresectable or metastatic biliary tract carcinoma in adults. KEYTRUDA administered SC:</p> <p><u>Melanoma</u> KEYTRUDA as monotherapy is indicated for the treatment of adults with advanced (unresectable or metastatic) melanoma.</p> <p>KEYTRUDA as monotherapy is indicated for the adjuvant treatment of adults with Stage IIB, IIC or III melanoma and who have undergone complete resection.</p> <p><u>Non-small cell lung carcinoma (NSCLC)</u> KEYTRUDA, in combination with platinum-containing chemotherapy as neoadjuvant treatment, and then continued as monotherapy as adjuvant treatment, is indicated for the treatment of resectable non-small cell lung carcinoma at high risk of recurrence in adults.</p> <p>KEYTRUDA as monotherapy is indicated for the adjuvant treatment of adults with non-small cell lung carcinoma who are at high risk of recurrence following complete resection and platinum-based chemotherapy.</p> <p>KEYTRUDA as monotherapy is indicated for the first line treatment of metastatic non-small cell lung carcinoma in adults whose tumours express PD L1 with a \geq 50% tumour proportion score (TPS) with no EGFR or ALK positive tumour mutations.</p>
--	--

Table I.1:

Product Overview

	<p>KEYTRUDA, in combination with pemetrexed and platinum chemotherapy, is indicated for the first-line treatment of metastatic non squamous non-small cell lung carcinoma in adults whose tumours have no EGFR or ALK positive mutations.</p> <p>KEYTRUDA, in combination with carboplatin and either paclitaxel or nab paclitaxel, is indicated for the first line treatment of metastatic squamous non-small cell lung carcinoma in adults.</p> <p>KEYTRUDA as monotherapy is indicated for the treatment of locally advanced or metastatic non-small cell lung carcinoma in adults whose tumours express PD L1 with a $\geq 1\%$ TPS and who have received at least one prior chemotherapy regimen. Patients with EGFR or ALK positive tumour mutations should also have received targeted therapy before receiving KEYTRUDA.</p> <p><u>Classical Hodgkin lymphoma (cHL)</u> KEYTRUDA as monotherapy is indicated for the treatment of adults with relapsed or refractory classical Hodgkin lymphoma who have failed autologous stem cell transplant (ASCT) or following at least two prior therapies when ASCT is not a treatment option.</p> <p><u>Urothelial carcinoma</u> KEYTRUDA, in combination with enfortumab vedotin, is indicated for the first-line treatment of unresectable or metastatic urothelial carcinoma in adults. KEYTRUDA as monotherapy is indicated for the treatment of locally advanced or metastatic urothelial carcinoma in adults who have received prior platinum containing chemotherapy.</p> <p>KEYTRUDA as monotherapy is indicated for the treatment of locally advanced or metastatic urothelial carcinoma in adults who are not eligible for cisplatin containing chemotherapy and whose tumours express PD L1 with a combined positive score (CPS) ≥ 10.</p> <p><u>Head and neck squamous cell carcinoma (HNSCC)</u> KEYTRUDA as monotherapy is indicated for the treatment of resectable locally advanced head and neck squamous cell carcinoma (HNSCC) as neoadjuvant treatment, continued as adjuvant treatment in combination with radiation therapy with or without concomitant cisplatin and then as monotherapy in adults whose tumours express PD-L1 with a CPS ≥ 1.</p> <p>KEYTRUDA, as monotherapy or in combination with platinum and 5 fluorouracil (5 FU) chemotherapy, is indicated for the first line treatment of metastatic or unresectable recurrent head and neck squamous cell carcinoma in adults whose tumours express PD L1 with a CPS ≥ 1. KEYTRUDA as monotherapy is indicated for the treatment of recurrent or metastatic head and neck squamous cell carcinoma in adults whose tumours express PD L1 with a $\geq 50\%$ TPS and progressing on or after platinum containing chemotherapy.</p> <p><u>Renal cell carcinoma (RCC)</u> KEYTRUDA, in combination with axitinib, is indicated for the first line treatment of advanced renal cell carcinoma in adults.</p> <p>KEYTRUDA, in combination with lenvatinib, is indicated for the first line treatment of advanced renal cell carcinoma in adults.</p> <p>KEYTRUDA as monotherapy is indicated for the adjuvant treatment of adults with renal cell carcinoma at increased risk of recurrence following nephrectomy, or following nephrectomy and resection of metastatic lesions.</p>
--	--

Table I.1: Product Overview

	<p><u>Microsatellite instability high (MSI-H) or mismatch repair deficient (dMMR) cancers</u></p> <p><u>Colorectal cancer (CRC)</u></p> <p>KEYTRUDA as monotherapy is indicated for adults with MSI-H or dMMR colorectal cancer in the following settings:</p> <ul style="list-style-type: none"> - first line treatment of metastatic colorectal cancer; - treatment of unresectable or metastatic colorectal cancer after previous fluoropyrimidine based combination therapy. <p><u>Non-colorectal cancers</u></p> <p>KEYTRUDA as monotherapy is indicated for the treatment of the following MSI H or dMMR tumours in adults with:</p> <ul style="list-style-type: none"> - advanced or recurrent endometrial carcinoma, who have disease progression on or following prior treatment with a platinum containing therapy in any setting and who are not candidates for curative surgery or radiation. - unresectable or metastatic gastric, small intestine, or biliary cancer, who have disease progression on or following at least one prior therapy. <p><u>Oesophageal carcinoma</u></p> <p>KEYTRUDA, in combination with platinum and fluoropyrimidine based chemotherapy, is indicated for the first-line treatment of locally advanced unresectable or metastatic carcinoma of the oesophagus in adults whose tumours express PD L1 with a CPS \geq 10.</p> <p><u>Triple negative breast cancer (TNBC)</u></p> <p>KEYTRUDA, in combination with chemotherapy as neoadjuvant treatment, and then continued as monotherapy as adjuvant treatment after surgery, is indicated for the treatment of adults with locally advanced, or early stage triple negative breast cancer at high risk of recurrence.</p> <p>KEYTRUDA, in combination with chemotherapy, is indicated for the treatment of locally recurrent unresectable or metastatic triple negative breast cancer in adults whose tumours express PD L1 with a CPS \geq 10 and who have not received prior chemotherapy for metastatic disease.</p> <p><u>Endometrial carcinoma (EC)</u></p> <p>KEYTRUDA, in combination with carboplatin and paclitaxel, is indicated for the first-line treatment of primary advanced or recurrent endometrial carcinoma in adults who are candidates for systemic therapy.</p> <p>KEYTRUDA, in combination with lenvatinib, is indicated for the treatment of advanced or recurrent endometrial carcinoma in adults who have disease progression on or following prior treatment with a platinum containing therapy in any setting and who are not candidates for curative surgery or radiation.</p> <p><u>Cervical cancer</u></p> <p>KEYTRUDA, in combination with chemoradiotherapy (external beam radiation therapy followed by brachytherapy), is indicated for the treatment of FIGO 2014 Stage III - IVA locally advanced cervical cancer in adults who have not received prior definitive therapy.</p> <p>KEYTRUDA, in combination with chemotherapy with or without bevacizumab, is indicated for the treatment of persistent, recurrent, or metastatic cervical cancer in adults whose tumours express PD L1 with a CPS \geq 1.</p> <p><u>Gastric or gastro-oesophageal junction (GEJ) adenocarcinoma</u></p> <p>KEYTRUDA, in combination with trastuzumab, fluoropyrimidine and platinum-containing chemotherapy, is indicated for the first-line treatment of locally advanced unresectable or metastatic HER2-positive gastric or gastro-oesophageal junction adenocarcinoma in adults whose tumours express PD-L1 with a CPS \geq 1.</p>
--	--

Table I.1: Product Overview

	<p>KEYTRUDA, in combination with fluoropyrimidine and platinum-containing chemotherapy, is indicated for the first-line treatment of locally advanced unresectable or metastatic HER2-negative gastric or gastro-oesophageal junction adenocarcinoma in adults whose tumours express PD L1 with a CPS \geq 1.</p> <p><u>Biliary tract carcinoma (BTC)</u></p> <p>KEYTRUDA, in combination with gemcitabine and cisplatin, is indicated for the first-line treatment of locally advanced unresectable or metastatic biliary tract carcinoma in adults.</p>
<p>Pharmaceutical form(s) and strengths</p>	<p>Pembrolizumab solution for infusion for intravenous (IV) administrations: 100 mg/4 mL (25 mg/mL) solution in a single-use vial</p> <p>Pembrolizumab solution for injection for subcutaneous (SC) administration: 395 mg/2.4mL (165 mg/mL) solution in a single-use vial 790 mg/4.8mL (165 mg/mL) solution in a single-use vial</p>
<p>Posology and route of administration in the EEA</p>	<p><u>Pembrolizumab IV</u></p> <p>The recommended dose of KEYTRUDA solution for infusion in adults is either 200 mg every 3 weeks or 400 mg every 6 weeks administered as an intravenous infusion over 30 minutes.</p> <p>The recommended dose of KEYTRUDA solution for infusion as monotherapy in paediatric patients aged 3 years and older with cHL or patients aged 12 years and older with melanoma is 2 mg/kg bodyweight (bw) (up to a maximum of 200 mg), every 3 weeks administered as an intravenous infusion over 30 minutes.</p> <p>For use in combination, see the Summary of Product Characteristics (SmPC) for the concomitant therapies.</p> <p>Patients should be treated with KEYTRUDA until disease progression or unacceptable toxicity (and up to maximum duration of therapy if specified for an indication). Atypical responses (i.e. an initial transient increase in tumour size or small new lesions within the first few months followed by tumour shrinkage) have been observed. It is recommended to continue treatment for clinically stable patients with initial evidence of disease progression until disease progression is confirmed.</p> <p>For the adjuvant treatment of melanoma, NSCLC, or RCC, KEYTRUDA should be administered until disease recurrence, unacceptable toxicity, or for a duration of up to one year.</p> <p>For the neoadjuvant and adjuvant treatment of resectable NSCLC, patients should be treated with neoadjuvant KEYTRUDA in combination with chemotherapy for 4 doses of 200 mg every 3 weeks or 2 doses of 400 mg every 6 weeks or until disease progression that precludes definitive surgery or unacceptable toxicity, followed by adjuvant treatment with KEYTRUDA as monotherapy for 13 doses of 200 mg every 3 weeks or 7 doses of 400 mg every 6 weeks or until disease recurrence or unacceptable toxicity. Patients who experience disease progression that precludes definitive surgery or unacceptable toxicity related to KEYTRUDA as neoadjuvant treatment in combination with chemotherapy should not receive KEYTRUDA monotherapy as adjuvant treatment. For the neoadjuvant and adjuvant treatment of resectable locally advanced HNSCC, patients should be treated with neoadjuvant KEYTRUDA as monotherapy for 2 doses of 200 mg every 3 weeks or 1 dose of 400 mg every 6 weeks or until disease progression that precludes definitive surgery or unacceptable toxicity, followed by adjuvant treatment with KEYTRUDA in combination with radiation with or without concomitant cisplatin for 3 doses of 200 mg every 3 weeks or 2 doses of 400 mg every 6 weeks followed by KEYTRUDA as monotherapy for 12 doses of 200 mg every 3 weeks or 6 doses of 400 mg every 6 weeks or until disease recurrence or unacceptable toxicity.</p>

Table I.1:**Product Overview**

	<p>For the neoadjuvant and adjuvant treatment of TNBC, patients should be treated with neoadjuvant KEYTRUDA in combination with chemotherapy for 8 doses of 200 mg every 3 weeks or 4 doses of 400 mg every 6 weeks or until disease progression that precludes definitive surgery or unacceptable toxicity, followed by adjuvant treatment with KEYTRUDA as monotherapy for 9 doses of 200 mg every 3 weeks or 5 doses of 400 mg every 6 weeks or until disease recurrence or unacceptable toxicity. Patients who experience disease progression that precludes definitive surgery or unacceptable toxicity related to KEYTRUDA as neoadjuvant treatment in combination with chemotherapy should not receive KEYTRUDA monotherapy as adjuvant treatment.</p> <p>For locally advanced cervical cancer, patients should be treated with KEYTRUDA concurrent with chemoradiotherapy, followed by KEYTRUDA as monotherapy. KEYTRUDA can be administered as either 200 mg every 3 weeks or 400 mg every 6 weeks until disease progression, unacceptable toxicity or up to 24 months.</p> <p><u>Pembrolizumab SC</u></p> <p>The recommended dose of KEYTRUDA solution for injection in adults is either:</p> <ul style="list-style-type: none"> - 395 mg every 3 weeks administered as a subcutaneous injection over 1 minute or - 790 mg every 6 weeks administered as a subcutaneous injection over 2 minutes. <p>For use in combination, see the Summary of Product Characteristics (SmPC) for the concomitant therapies.</p> <p>Patients should be treated with KEYTRUDA until disease progression or unacceptable toxicity (and up to maximum duration of therapy if specified for an indication). Atypical responses (i.e. an initial transient increase in tumour size or small new lesions within the first few months followed by tumour shrinkage) have been observed. It is recommended to continue treatment for clinically stable patients with initial evidence of disease progression until disease progression is confirmed.</p> <p>For the adjuvant treatment of melanoma, NSCLC, or RCC, KEYTRUDA should be administered until disease recurrence, unacceptable toxicity, or for a duration of up to one year.</p> <p>For the neoadjuvant and adjuvant treatment of resectable NSCLC, patients should be treated with neoadjuvant KEYTRUDA in combination with chemotherapy for 4 doses of 395 mg every 3 weeks or 2 doses of 790 mg every 6 weeks or until disease progression that precludes definitive surgery or unacceptable toxicity, followed by adjuvant treatment with KEYTRUDA as monotherapy for 13 doses of 395 mg every 3 weeks or 7 doses of 790 mg every 6 weeks or until disease recurrence or unacceptable toxicity. Patients who experience disease progression that precludes definitive surgery or unacceptable toxicity related to KEYTRUDA as neoadjuvant treatment in combination with chemotherapy should not receive KEYTRUDA monotherapy as adjuvant treatment.</p> <p>For the neoadjuvant and adjuvant treatment of resectable locally advanced HNSCC, patients should be treated with neoadjuvant KEYTRUDA as monotherapy for 2 doses of 395 mg every 3 weeks or 1 dose of 790 mg or until disease progression that precludes definitive surgery or unacceptable toxicity, followed by adjuvant treatment with KEYTRUDA in combination with radiation with or without concomitant cisplatin for 3 doses of 395 mg every 3 weeks or 2 doses of 790 mg every 6 weeks followed by KEYTRUDA as monotherapy for 12 doses of 395 mg every 3 weeks or 6 doses of 790 mg every 6 weeks or until disease recurrence or unacceptable toxicity. Patients who experience disease progression that precludes definitive surgery or unacceptable toxicity related to KEYTRUDA monotherapy as neoadjuvant treatment should not receive</p>
--	--

Table I.1: Product Overview

	<p>KEYTRUDA in combination with radiation with or without concomitant cisplatin as adjuvant treatment.</p> <p>For the neoadjuvant and adjuvant treatment of TNBC, patients should be treated with neoadjuvant KEYTRUDA in combination with chemotherapy for 8 doses of 395 mg every 3 weeks or 4 doses of 790 mg every 6 weeks or until disease progression that precludes definitive surgery or unacceptable toxicity, followed by adjuvant treatment with KEYTRUDA as monotherapy for 9 doses of 395 mg every 3 weeks or 5 doses of 790 mg every 6 weeks or until disease recurrence or unacceptable toxicity. Patients who experience disease progression that precludes definitive surgery or unacceptable toxicity related to KEYTRUDA as neoadjuvant treatment in combination with chemotherapy should not receive KEYTRUDA monotherapy as adjuvant treatment.</p> <p>For locally advanced cervical cancer, patients should be treated with KEYTRUDA concurrent with chemoradiotherapy, followed by KEYTRUDA as monotherapy. KEYTRUDA can be administered as either 395 mg every 3 weeks or 790 mg every 6 weeks until disease progression, unacceptable toxicity or up to 24 months.</p>
<p>Is/will the product be subject to additional monitoring in the EU?</p>	<p>No</p>

PART II: SAFETY SPECIFICATION

PART II: MODULE SI - EPIDEMIOLOGY OF THE INDICATION(S) AND TARGET POPULATION(S)

Melanoma

Incidence and prevalence:

Over 80% of the world's 287,700 annual new cases of melanoma occur in Australia, Europe, or North America [Ref. 5.4: 05DXHB]. Incidence varies according to geographic latitude, with the highest rates (per 100,000) found in Australia and New Zealand (33.6) and lowest in South-Central Asia (0.3) [Ref. 5.4: 05DXHB]. In Europe, incidence (per 100,000) ranges from 5.3 in the Central/Eastern region to 18.8 in Western countries, with 144,209 incident cases expected in 2018 [Ref. 5.4: 05DXHB]. In the United States (US), incidence is 22.2 per 100,000, corresponding to 96,480 new cases in 2019 [Ref. 5.4: 05DN03]. The 5-year prevalence in Europe is 494,111 cases [Ref. 5.4: 05DXHB].

Demographics of the indication population and risk factors for the disease:

Although melanoma incidence increases by age worldwide with a median age at diagnosis of 65 years [Ref. 5.4: 05DN03], it is one of the most common cancers among adolescents and young adults [Ref. 5.4: 05DR9D]. Incidence is higher for females until age 45-55, after which rates are higher among males [Ref. 5.4: 04KB3D, 03MS03]. Risk factors include fair skin, multiple dysplastic nevi, and family history of melanoma; ultraviolet light also plays a role [Ref. 5.4: 03MS0N].

The main existing treatment options

Most patients with early-stage disease will be cured by excision [Ref. 5.4: 05DLL8]. Adjuvant radiation therapy may be used for select patients with positive nodes or other features predicting high risk of relapse. For metastatic or unresectable melanoma, multiple targeted therapies have been FDA-approved, including immune checkpoint inhibitors, BRAF V600 targeted treatment, and BRAF/MEK combinations. Several drugs within these classes have been approved in the adjuvant setting.

Natural history of the indicated condition, including mortality and morbidity:

Mortality rates (per 100,000) are highest in Australia and New Zealand (3.4) and lowest in South-Central Asia (0.19) [Ref. 5.4: 05DXHB]. In Europe, mortality (per 100,000) ranges from 1.5 in the Southern region to 2.0 in Northern countries, with 27,147 deaths expected in 2018 [Ref. 5.4: 05DXHB]. In the US and Europe, the 5-year relative survival is 92% and 83%, respectively [Ref. 5.4: 04KJL, 05DN03]. Stage-specific overall survival varies across Europe, as follows: 95%–100% (stage I), 65%–92.8% (stage II), 41%–71% (stage III), and 9%–28% (stage IV) [Ref. 5.4: 05DRW3].

Important co-morbidities

Most melanomas occur among adults over age 50 [Ref. 5.4: 05DN03] for whom there is an increased risk of cardiovascular or cerebrovascular diseases, diabetes, and secondary cancers compared to younger adults [Ref. 5.4: 057XQG]. A higher prevalence of comorbidity may be associated with higher melanoma stage [Ref. 5.4: 05DQKR, 03N0KS].

Non-Small Cell Lung Cancer (NSCLC)

Incidence and prevalence:

Approximately 2,480,301 new lung cancer cases were diagnosed worldwide in 2022 [Ref. 5.4: 08LCQN], with 89% of all lung cancers being classified as NSCLC [Ref. 5.4: 08JD46]. Globally, lung cancer is the most commonly diagnosed tumor type accounting for 12.4% of new cancer cases [Ref. 5.4: 08LCQN], although the geographical distribution of lung cancer incidence varies widely with smoking prevalence [Ref. 5.4: 08LCQN]. The age-standardized incidence rate of lung cancer is highest in Eastern Asia (39.4 per 100,000), Polynesia (37.5), and North America (31.9); rates are lowest in Western, Middle, and Eastern Africa (2.1, 2.3, and 3.2 respectively) [Ref. 5.4: 08QGQ2]. In Europe, lung cancer incidence rates range from 27.6 per 100,000 in Eastern Europe to 31.2 in Western Europe, with 484,554 new cases diagnosed in 2022 [Ref. 5.4: 08QGQ2]. The 5-year prevalence of lung cancer in Europe is 610,169 cases [Ref. 5.4: 08QGQ2]. In the US, lung cancer incidence is 31.9 per 100,000, corresponding to 226,033 new cases in 2022, with a 5-year prevalence of 288,303 cases [Ref. 5.4: 08QGQ2].

Demographics of the indication population and risk factors for the disease:

Lung cancer is most frequently diagnosed between the ages of 65 and 74 years [Ref. 5.4: 058TT3]. The incidence and mortality rates of lung cancer are consistently higher in men than women regardless of geographical region [Ref. 5.4: 08LCQN], with incidence rates rising among women [Ref. 5.4: 05DQP9]. Tobacco exposure is the primary risk factor for developing lung cancer [Ref. 5.4: 05DQP9] and two-thirds of lung cancer deaths are attributable to smoking [Ref. 5.4: 05QDW7].

The main existing treatment options

Surgery provides the best chance for a cure among patients with stage I or II disease [Ref. 5.4: 05DMDQ]. Chemoradiation may improve survival for locally advanced unresectable disease. Patients with metastatic disease may achieve improved survival and palliation of symptoms with chemotherapy (typically a platinum doublet), targeted agents, checkpoint inhibitors, and other supportive measures combined or alone.

Natural history of the indicated condition, including mortality and morbidity:

Worldwide, lung cancer was the leading cause (18.7%) of all cancer deaths (1.8 million) in 2022 [Ref. 5.4: 08LCQN]. Age-standardized lung cancer mortality rates range from 31.7 per 100,000 in Polynesia, 30.5 in Micronesia, and 25.1 in Eastern Asia to 2.0 in Western Africa, 2.2 in Middle Africa, and 3.0 in Eastern Africa [Ref. 5.4: 08QGQ2]. Overall, lung cancer

survival remains poor with 5-year survival rates below 20% in most countries [Ref. 5.4: 08LCQN].

Important co-morbidities:

As lung cancer is primarily a disease of the aged and most common among smokers, cardiovascular disease (CVD), chronic obstructive pulmonary (COPD), diabetes, and other malignancies are expected to be the most prevalent comorbidities in NSCLC [Ref. 5.4: 0439J6].

Hodgkin Lymphoma (HL)

Incidence and prevalence:

Hodgkin lymphoma (HL) is a rare lymphoid malignancy of B-cell origin involving the lymph nodes and lymphatic system [Ref. 5.4: 052ZM2]. There are two major variants of HL, the uncommon nodular lymphocyte predominant (NLP) subtype (<10%) and the classical variety divided into four subtypes (lymphocyte-rich, 5%; nodular sclerosis, 70%; mixed cellularity, 20-25%; or lymphocyte-depleted, <1%) [Ref. 5.4: 052ZM2]. The estimated new cases globally 2018 was estimated to be 79,990 (1.0 per 100,000 world populations) [Ref. 5.4: 05DXHB]. Incidence (per 100,000) is highest in Southern (2.8) and Northern (2.6) Europe; rates are lowest in Eastern Asia (0.35) [Ref. 5.4: 05DXHB]. Nearly one quarter of the world's HL cases are diagnosed in Europe, where an estimated 19,193 new cases (2.4 per 100,000) of HL were reported in 2018 [Ref. 5.4: 05DXHB]. The incidence of HL in European children 0-19 years of age is 1.3 per 100,000, yielding 2193 new cases in 2018 [Ref. 5.4: 05DXHB]. In the US, incidence is 2.7 per 100,000, corresponding to 8100 new cases in 2019; 12% of cases occur in persons under age 20 [Ref. 5.4: 05DMHL]. The 5-year overall prevalence in Europe is 77,315 cases, with 6289 children ages 0-19 years [Ref. 5.4: 05DXHB].

Demographics of the indication population and risk factors for the disease:

HL is more common in males and is most frequently diagnosed in young adults ages 20-34 years [Ref. 5.4: 052ZM2]. HL is one of the few malignancies with shared features of biology and natural history for children and adults [Ref. 5.4: 05G5VD]. Risk factors are not well described, but a viral etiology has long been hypothesized [Ref. 5.4: 052ZM2]. Epstein-Barr virus (EBV) is highly prevalent among HL patients and accounts for up to 40-50% of all global cases [Ref. 5.4: 04HTK5]. However, the rate of EBV-positive HL varies considerably by geography with lower a prevalence observed in North America (32%) and Europe (36%) compared to Africa (74%), Central/South America (61%), and Asia (56%) [Ref. 5.4: 04HTK5]. HL in young children is more likely to be EBV-associated compared to adolescents and young adults [Ref. 5.4: 05G5VD].

The main existing treatment options:

Historically, treatment of classical HL in childhood and adults generally utilized the same strategy and agents, with high cure rates of at least 80% of patients overall; 90-95% of children can be cured [Ref. 5.4: 05DMDL, 05G5VD]. Current management of classical HL involves initial treatment with chemotherapy alone or combined with radiation therapy for early stage

disease. Several chemotherapy options and brentuximab vedotin are available for stage III-IV disease. High-dose salvage chemotherapy followed by autologous stem cell transplantation and checkpoint inhibitors are options for refractory or relapsed disease. The NLP subtype is characterized by an indolent course and occasional late relapse; most patients present with early-stage disease. Radiation therapy is the preferred treatment for early stage NLPHL; chemotherapy with or without rituximab/radiation is recommended for stage III-IV NLPHL.

Natural history of the indicated condition, including mortality and morbidity:

In 2018, 26,167 HL deaths (0.3 per 100,000) occurred in the world. Mortality (per 100,000) is highest in Northern Africa (0.63) and lowest in East Asia (0.13); in Europe, mortality ranges from 0.19 in Western countries to 0.43 in Central and Eastern region, resulting in 4307 deaths in 2018 [Ref. 5.4: 05DXHB]. The overall 5-year survival in Europe is 80.3% [Ref. 5.4: 05DPN4]. The overall 5-year survival in the US is 86.6%, with stage-specific survival as follows: 91.6% (stage I), 93.5% (stage II), 82.9% (stage III), and 73.5% (stage IV) [Ref. 5.4: 05DMHL]. Among children and adolescents, 5-year survival is more than 95% [Ref. 5.4: 05G5VD]. Of the 1000 HL deaths expected in 2019 in the US, 0.9% occurred among persons under age 20 [Ref. 5.4: 05DMHL].

Important co-morbidities:

Given the relatively young average age of HL patients, fewer common age-related comorbid conditions would be expected compared to patients with other cancers.

Urothelial Carcinoma

Incidence and prevalence:

Bladder cancer comprises 90-95% of all urothelial carcinomas (UCs); the remaining are upper tract urothelial cancers [Ref. 5.4: 05DMLD]. Overall, the age standardized incidence (per 100,000) is 5.7 globally [Ref. 5.4: 05DXHB]. The age standardized incidence per 100,000 persons is highest in Southern Europe (15.2) and lowest in Middle Africa (1.2) [Ref. 5.4: 05DXHB]. Rates are generally declining in developed countries and increasing in less developed regions [Ref. 5.4: 04GXQB]. In the US, there are 82,290 new cases of bladder cancer expected in 2023 [Ref. 5.4: 08CRCF]. The 5-year bladder cancer prevalence is 308,011 in North America and 631,337 in Europe [Ref. 5.4: 05DXHB].

Demographics of the indication population and risk factors for the disease:

Bladder cancer incidence is 3-5 times higher in males than females [Ref. 5.4: 05DXHB]. Bladder cancer usually occurs among older adults; the median age at diagnosis is 73 years in the US [Ref. 5.4: 057K9G]. Tobacco consumption is a significant risk factor for urothelial cancer [Ref. 5.4: 05DMLD].

The main existing treatment options:

Muscle-invasive bladder cancer (MIBC) represents approximately 20-25% of primary diagnoses with a higher risk of mortality (compared with non-muscle -invasive bladder cancer

(NMIBC)). Standard treatment options for MIBC include radical cystectomy and pelvic lymph node dissection (RC + PLND) with or without neoadjuvant (cisplatin-based) or adjuvant therapies and bladder-preserving approaches for MIBC, and systemic therapy for locally advanced and metastatic disease.

Natural history of the indicated condition, including mortality and morbidity:

Cancer-related mortality most commonly occurs in advanced/metastatic disease. Approximately 5-8% of bladder cancer patients have locally advanced/metastatic disease at the time of initial diagnosis; additionally, roughly half of all patients experience relapse after cystectomy, with distant metastases most common [Ref. 5.4: 059FM9]. Mortality rates (per 100,000) are highest in Northern Africa (4.4) and lowest in Central America (0.75); European rates range from 2.6 (Northern) to 3.2 (Southern) [Ref. 5.4: 05DXHB]. The 5-year survival by stage is as follows: in situ: 95.8%, local: 69.5%, regional: 36.3%, distant: 4.6% [Ref. 5.4: 057K9G]. The overall 5-year relative survival in Europe is 68% [Ref. 5.4: 04D76G].

Important co-morbidities:

As bladder cancer is primarily a disease of the elderly, CVD, COPD, diabetes and other malignancies are expected to be the most prevalent comorbidities in patients with urothelial cancers [Ref. 5.4: 04H4GM, 04GXN0].

Head and Neck Squamous Cell Carcinoma (HNSCC)

Incidence and prevalence:

Head and neck cancer (HNC) encompasses a heterogeneous group of cancers of the upper aero-digestive tract, including the oral cavity, salivary glands, pharynx, and larynx [Ref. 5.4: 04DP5M]. Cancers of the lip and oral cavity, oropharynx, hypopharynx, and larynx, which are primarily (>90%) comprised of squamous cell carcinoma histology, are collectively termed head and neck squamous cell carcinoma (HNSCC) [Ref. 5.4: 04DGDP, 04DP5M]. Worldwide, HNSCC incidence varies widely by country and tumor site; the most common sites in Europe are lip and oral cavity (42%) and larynx (27%). Across Europe, combined HNSCC age-standardized incidence (per 100,000) ranges by region from 8.6 (Southern Europe) to 11.9 (Eastern Europe), corresponding to approximately 148,346 new cases and 475,745 5-year prevalent cases in 2022 [Ref. 5.4: 08PQSD]. In the US, it is projected that in 2024 there will be 71,100 new cases and 16,110 deaths of HNC located in the oral cavity, pharynx, and larynx [Ref. 5.4: 08HZXM].

Demographics of the population in the indication population and risk factors for the disease:

HNC occurs most often among adults over age 60 years [Ref. 5.4: 057SQL, 05DMHM]. However, some oropharyngeal cancer cases are related to human papillomavirus (HPV) and tend to occur in younger adults [Ref. 5.4: 04DGDY]. Males are 2-3 times more likely to have HNSCC compared to females [Ref. 5.4: 05DXHB]. Tobacco and alcohol consumption are the dominant risk factors, particularly for cancers of the oral cavity and larynx,

[Ref. 5.4: 04DP5M]. Increased HPV prevalence is largely responsible for a dramatic rise in oropharyngeal cancer in North America and Western Europe [Ref. 5.4: 04DGDY].

The main existing treatment options:

HNSCC treatment is complex with specific site, stage and pathology directing management. Single-modality surgery or radiation therapy is generally recommended for early-stage disease [Ref. 5.4: 05DLLB] [Ref. 5.4: 08PQSB]. Standard options for locally advanced stage III or IV tumors include surgery plus postoperative radiotherapy. Combined concomitant chemoradiation may be recommended post-operatively for high-risk features or unresectable tumors. Commonly used agents include cisplatin, carboplatin, paclitaxel, docetaxel, 5-FU, methotrexate, capecitabine, and cetuximab. In addition to cytotoxic chemotherapy, several targeted therapies and checkpoint inhibitors are available as treatment options. In 2019, pembrolizumab was approved as a first-line treatment for patients with metastatic or unresectable recurrent HNSCC with PD-L1 CPS>1 and in combination with platinum and 5-fluorouracil for any HNSCC cancer regardless of PD-L1 status [Ref. 5.4: 08PQSB].

Natural history of the indicated condition, including mortality and morbidity:

Across European regions, combined HNSCC mortality rates range from 2.8 to 6.6 per 100,000 [Ref. 5.4: 08PKM9]. In Europe, five-year survival ranges from 33-44%, with the majority of cases diagnosed at stage III or IV [Ref. 5.4: 04SRJJ]. Patients with HPV-associated HNC generally have an improved prognosis and respond better to chemotherapy [Ref. 5.4: 05DLLB].

Important co-morbidities:

Studies of HNSCC patients in Europe found the most common co-morbidities to be cardiovascular disease, asthma/COPD, diabetes, liver disease, and depression [Ref. 5.4: 04D0C7, 04DDMM].

Renal Cell Carcinoma (RCC)

Incidence and prevalence:

The age-standardized incidence of kidney cancer (per 100,000) is highest in North America (12.2) and Northern Europe/Australia and New Zealand (10.3); rates are lowest in Middle Africa (1.0) [Ref. 5.4: 06C8B2]. More than one-third of incident cases occur in Europe, with nearly 139,000 incident cases expected in 2020 [Ref. 5.4: 06C86L]. In the US, kidney cancer incidence is anticipated to be 16.4 per 100,000, yielding roughly 76,000 new cases in 2021 [Ref. 5.4: 06C86L]. Approximately 85% of all primary kidney cancers are comprised by RCC, and approximately 70% have clear cell histology, [Ref. 5.4: 05DMDM]. RCC incidence has increased over the last few decades in the US and most of Europe [Ref. 5.4: 05BFQQ, 04BN0K]. The 5-year prevalence of kidney cancer in Europe is 405,983 cases [Ref. 5.4: 06C8B2].

Demographics of the indication population and risk factors for the disease:

Kidney cancer incidence increases by age with a median age at diagnosis of 64 years in the US [Ref. 5.4: 06CL4Y]. In most countries, it is roughly twice as common among males compared to females [Ref. 5.4: 06C8B2]. Established risk factors for RCC include obesity, smoking, hypertension, and chronic kidney disease; other probable risk factors include low physical activity, diabetes, occupational chemical exposure, radiation exposure, and analgesic use [Ref. 5.4: 05DPRK, 053GXR, 053GXZ, 053GY2].

The main existing treatment options:

Surgical resection is the primary treatment for localized disease [Ref. 5.4: 05DMDM]. There is little evidence of benefit with adjuvant therapy for most patients with locally advanced disease. Approved and emerging targeted therapies for advanced or relapsed RCC include checkpoint inhibitors, anti-vascular endothelial growth factor (VEGF) antibodies, tyrosine kinase inhibitors (TKI), and mammalian target of rapamycin (mTOR) protein inhibitors.

Natural history of the indicated condition, including mortality and morbidity:

Worldwide, kidney cancer age-standardized mortality rates (per 100,000) are highest in Central/Eastern Europe (3.4) and Western Europe (2.8); 54,000 deaths occurred in Europe during 2020 [Ref. 5.4: 06C8B2]. Prognosis has improved significantly in the US and Europe, due in part to the advent of TKI therapy [Ref. 5.4: 053TKZ, 05BFQQ]. The majority (65%) of kidney cancers diagnosed in the US is localized and 16% of tumors are metastatic [Ref. 5.4: 06CL4Y]. The overall 5-year survival in Europe and the US is 60% and 76%, respectively [Ref. 5.4: 06CL4Y, 04D76G]. Clear cell histology, accounting for the majority of RCC, is associated with a better prognosis than non-clear cell RCC [Ref. 5.4: 053GY0].

Important co-morbidities:

Cardiovascular or cerebrovascular diseases, hypertension, chronic obstructive pulmonary disease, diabetes, and other prevalent comorbidities among elderly populations are frequently observed in cancer patients [Ref. 5.4: 04VW4L].

Microsatellite Instability-High (MSI-H)/deficient Mismatch Repair (dMMR) Colorectal Cancer (CRC)

Incidence and prevalence:

The prevalence of MSI-H/dMMR status varies widely by tumor type and disease stage [Ref. 5.4: 04TCGC, 04TCK9]. MSI-H/dMMR is present in approximately 15% of all colorectal cancer [Ref. 5.4: 04TC0P]. MSI-H/dMMR is more common among stage II (~20%) than stage III (~12%) colorectal cancer and is least frequent in stage IV (~4%) disease [Ref. 5.4: 04TC0P].

Demographics of the indication population and risk factors for the disease:

Lynch Syndrome results from germline mutations in mismatch repair (MMR) genes and accounts for a small portion of MSI-H/dMMR colorectal cancers; the majority of MSI-H/dMMR tumors arise in the sporadic form [Ref. 5.4: 04TC0P]. Few consistent differences have been observed for the association between MSI status and lifestyle factors among colorectal cancer patients [Ref. 5.4: 05FB0L]. However, some studies report a higher proportion of MSI-H/dMMR tumors for current smokers [Ref. 5.4: 04W005, 05FB0L] and those with a family history of colorectal cancer [Ref. 5.4: 04VYPZ, 04W02X]. BRAF mutations are more common in MSI-H/dMMR colorectal patients [Ref. 5.4: 04VWL2].

The main existing treatment options:

Pembrolizumab was approved in the US to treat MSI-H or MMR deficient colorectal cancer following progression on a fluoropyrimidine, oxaliplatin, and irinotecan treatment. Prior to the approval of pembrolizumab, there were no FDA-approved therapies specifically for the treatment of MSI-H/dMMR cancers and the standard of care consisted of chemotherapy regimens and palliative care if response was not achieved. Nivolumab (as a single agent or in combination with ipilimumab) is approved for MSI-H or dMMR metastatic CRC that has progressed following treatment with a fluoropyrimidine, oxaliplatin, and irinotecan [Ref. 5.4: 05DLL7]. Routine testing for MMR deficiency or MSI status is recommended for all patients with colorectal cancers [Ref. 5.4: 05DLL7].

Natural history of the indicated condition, including mortality and morbidity:

Sporadic MSI-H/dMMR colorectal cancer is generally associated with lower stage at diagnosis, larger tumor size, proximal colon predominance, wildtype p53 status, poor differentiation, mucinous histology, and less metastasis [Ref. 5.4: 04VZ0J, 04VWL8]. MSI-H/dMMR colorectal cancer patients have a survival advantage limited to those with localized disease [Ref. 5.4: 04VZ0J, 043SX8]. In contrast, mCRC patients with MSI-H/dMMR had worse survival and poor prognosis compared to microsatellite stable (MSS) mCRC [Ref. 5.4: 043VF3].

Important co-morbidities:

Co-morbidity prevalence among MSI-H/dMMR cancer patients is unknown. However, co-morbidities are common among cancer patients, particularly with older adults for whom cancer is most common [Ref. 5.4: 04VW4L]. Secondary tumors or multiple primary have been reported among MMR deficient and MSI-H colorectal cancer patients [Ref. 5.4: 04W654, 04VZ03].

Esophageal Cancer

Incidence and prevalence:

Over 77% of the 572,034 new esophageal cancer cases diagnosed worldwide occur in Asia [Ref. 5.4: 05DXHB]. The highest age-standardized incidence rates (per 100,000) are found in Eastern Asia (12.2) and the lowest are in Central America (0.96) [Ref. 5.4: 05DXHB]. In

Europe, nearly 53,000 new cases were diagnosed in 2018; incidence rates range from 3.0 in Central/Eastern countries to 5.5 in Northern Europe [Ref. 5.4: 05DXHB]. In the US, incidence is 4.3 per 100,000, with 17,650 new cases in 2019 [Ref. 5.4: 05DMHK]. Squamous cell carcinoma is the predominant histologic subtype worldwide; however, adenocarcinoma is most frequently diagnosed in the US and much of Europe [Ref. 5.4: 054HBS]. The 5-year prevalence in Europe is approximately 56,000 cases [Ref. 5.4: 05DXHB].

Demographics of the indication population and risk factors for the disease:

Esophageal cancer incidence increases with age, with a median age at diagnosis of 68 years in the US [Ref. 5.4: 05DMHK]. In most regions, including the US and Europe, males have a 3- to 5-fold higher incidence compared to females [Ref. 5.4: 05DXHB]. Smoking is the primary risk factor for the squamous cell carcinoma subtype, but alcohol use, poor nutrition, and infectious agents have also been implicated [Ref. 5.4: 054HBS]. Gastroesophageal reflux disease occurs in 10-20% of adults in Western countries and is a major risk factor for adenocarcinoma, either directly or through an intermediate pre-malignant lesion, Barrett's esophagus [Ref. 5.4: 054HBS].

The main existing treatment options:

Surgical resection is the standard treatment for early-stage disease. Locally advanced disease is generally managed by multimodal approaches including, perioperative chemotherapy, neoadjuvant chemoradiation with or without adjuvant immunotherapy (Nivolumab), definitive chemoradiation (dCRT) or dCRT with salvage surgery. Patients with esophageal squamous cell carcinoma (ESCC) are treated with chemoradiotherapy (CRT) followed by surgery or dCRT with salvage surgery while patients with esophageal adenocarcinoma (EAC) receive neoadjuvant CRT or perioperative chemotherapy followed by surgery. Treatment with dCRT is recommended for ESCC or EAC patients who are unresectable or refuse surgery. In the metastatic setting, patients with EAC are managed similarly to patients with advanced gastric cancer. In general, platin/fluoropyrimidine-based systemic chemotherapy with or without immunotherapy is the current standard of care. For patients with EAC, a trastuzumab-containing regimen may be indicated for HER2-positive tumors. In the second and later lines, approvals of immunotherapy for ESCC has provided additional treatment options to chemotherapy [Ref. 5.4: 08DSV2].

Natural history of the indicated condition, including mortality and morbidity:

In Europe, mortality rates (per 100,000) range from 1.5 in Southern Europe to 4.1 in Northern Europe [Ref. 5.4: 08JK5].[Ref. 5.4: 05DXHB]The overall 5-year survival in the US and Europe is 21.6% and 5.4-23.9%, respectively [Ref. 5.4: 08KZXV][Ref. 5.4: 05DMHK, 054HM8][Ref. 5.4: 08QPK8]. Most cases are diagnosed at regional (32%) or distant (39%) stage, contributing to the poor overall survival [Ref. 5.4: 08KZXV] [Ref. 5.4: 05DMHK].

Important co-morbidities:

Most esophageal cancers occur among adults over age 60 years who have an increased risk of cardiovascular or cerebrovascular diseases, diabetes, and second primary malignancies, such as cancers of the oropharynx, lung and stomach [Ref. 5.4: 054HS0].

Triple Negative Breast Cancer (TNBC)

Incidence and prevalence:

Triple negative breast cancer (TNBC) is an aggressive subtype of breast cancer (BC) defined by an absence of expression of the estrogen receptor and progesterone receptor, and by absence of overexpression or amplification of the human epidermal growth factor receptor 2 [Ref. 5.4: 059X3Q]. TNBC comprises 15-20% of all BC, with infiltrating ductal carcinoma (not otherwise specified) accounting for approximately 92% of TNBC cases [Ref. 5.4: 059PJJ]. Data on the incidence and prevalence rates of TNBC are scarce. Based on overall BC incidence (per 100,000) for the world, US and Europe, the estimated number of annual TNBC cases is roughly 365,000, 91,000, and 47,000, respectively [Ref. 5.4: 05DXHB] [Ref. 5.4: 059RH3].

Demographics of the indication population and risk factors for the disease:

TNBC is most common in young, pre-menopausal patients; although BC incidence generally increases with age, the proportion of BC cases categorized with the TNBC subtype decreases with age [Ref. 5.4: 05DNTY]. In the US, women of African ancestry are twice as likely to have the TNBC subtype compared to other women with BC [Ref. 5.4: 05DNTY]. TNBC etiology is not well understood. Compared to other BC subtypes, TNBC has a higher prevalence of BRCA1 or BRCA2 germline mutations. The reproductive factors generally associated with increased BC risk, such as increased parity, younger age at first birth, older age at menarche, and oral contraceptive use, are not strongly linked to TNBC risk [Ref. 5.4: 05DQP0].

The main existing treatment options:

The standard of care for TNBC continues to be conventional chemotherapy and radiation as with BC in general; optimal regimens for TNBC have yet to be established. For women with metastatic breast cancer, including TNBC, who have germline BRCA mutations, PARP inhibitors may be considered. For PD-L1 expressing metastatic TNBC, PD-L1 inhibitors in combination with chemotherapy are newer options.

Management of early-stage breast cancer has been based on use of systemic chemotherapy given prior to (neoadjuvant therapy, NAC) or after (adjuvant therapy) definitive surgery according to NCCN and ESMO guidelines [Ref. 5.4: 05MDND, 05B2QG]. Patients with high-risk, early-stage TNBC who are eligible for NAC are primarily treated with taxane- and anthracycline-based chemotherapy regimens. Until recently, radiation therapy was the only adjuvant treatment option, if clinically indicated, for patients who received chemotherapy prior to surgery. Capecitabine has been added as a recommended adjuvant treatment for patients not achieving pathological complete response (pCR) following NAC.

Natural history of the indicated condition, including mortality and morbidity:

Important characteristics of TNBC are high proliferation and frequent metastasis to visceral organs and brain; TNBC tumors have the highest recurrence rates compared to other BC subtypes [Ref. 5.4: 05DNRG]. Compared to other hormonal and HER2 subtypes, TNBC has

the poorest survival and prognosis is consistently lower at each disease stage; TNBC 3-year survival ranges from 94% for stage IA disease to 20% for stage IV [Ref. 5.4: 05DNNW].

Important co-morbidities:

Data describing specific co-morbidities among TNBC patients are scarce. A recent study of more than 50,000 TNBC patients reported 82% with no comorbid conditions [Ref. 5.4: 05DRW2].

Endometrial Carcinoma

Incidence and prevalence:

The age-standardized incidence of carcinoma of the uterine corpus, often referred to as endometrial carcinoma, is highest in North America (21.1 per 100,000) and lowest in Middle Africa (2.3 per 100,000) [Ref. 5.4: 08G0HL]. In Europe, the age-standardized incidence ranges from 12.9 per 100,000 in Western countries to 20.2 per 100,000 in the Central/Eastern region, yielding roughly 130,000 new cases in 2020 [Ref. 5.4: 08G0HL]. The age-adjusted incidence in the US is 27.6 per 100,000 women with 66,200 new cases expected in 2023 [Ref. 5.4: 087X38]. The estimated 5-year prevalence in Europe is 482,952 cases [Ref. 5.4: 08G0HL].

Demographics of the indication population and risk factors for the disease:

Endometrial carcinoma is most frequently diagnosed among women aged 55-64 years (63% of cases) with a median age at diagnosis of 63 years in the US [Ref. 5.4: 087X38]. In Europe, the median age at diagnosis is between 64-69 years [Ref. 5.4: 05Q3MX]. In the US, non-Hispanic black women have the highest incidence and mortality compared to women of other races/ethnicities [Ref. 5.4: 087X38]. The main risk factors for endometrial carcinoma are related to increased estrogen, including excess weight and abdominal fat, early age at menarche, late menopause, nulliparity, and diabetes [Ref. 5.4: 04SXYB] [Ref. 5.4: 08G0JV]. Genetic factors account for approximately 5% of endometrial carcinoma [Ref. 5.4: 0874ZG]. Cancer predisposition syndromes, such as Lynch Syndrome, are associated with an increased lifetime risk of developing endometrial carcinoma [Ref. 5.4: 08G0HC].

The main existing treatment options:

The main categories for delineating treatment include disease limited to the uterus, involvement of the cervix, and suspected extrauterine disease [Ref. 5.4: 0874ZG]. Minimally invasive surgery is preferred for disease limited to the uterus, whereas total or radical hysterectomy with bilateral salpingo-oophorectomy is recommended for tumors with cervical involvement. Radiotherapies or systemic therapies may be considered for uterine-confined disease. Multi-agent chemotherapy regimens are preferred in the primary/adjuvant setting for extrauterine or recurrent disease and high-risk histology types. Immunotherapy, such as pembrolizumab, is effective for treating MSI-H/dMMR tumors after prior systemic therapy. When combined with lenvatinib, pembrolizumab can be an option for patients who have received prior systemic therapy. Emergent therapies currently under investigation target CTLA-4, CDK4/6, PI3K/mTOR, PARP, and AMP kinase [Ref. 5.4: 08G0HQ].

Natural history of the indicated condition, including mortality and morbidity:

In Europe, the age-standardized mortality of endometrial carcinoma varies regionally, being highest in Central/Eastern countries (3.7 per 100,000) and lowest in the Western region (2.0 per 100,000) [Ref. 5.4: 08G0HL]. In the US, the age-adjusted death rate was 5.1 per 100,000 women per year during 2016-2020 [Ref. 5.4: 087X38]. A majority (67%) of endometrial cancer cases in the US are diagnosed as localized tumors with 19% having regional disease and 10% diagnosed at distant stage; the corresponding 5-year survival rates were 95%, 70% and 18%, respectively [Ref. 5.4: 087X38]. Despite early detection, approximately 13% of all endometrial carcinomas still experience recurrence [Ref. 5.4: 04SJ3G]. **Important comorbidities:**

Co-morbidities are common among patients with cancer, particularly in older adults [Ref. 5.4: 057XQG]. Most endometrial carcinoma occurs among women over age 55 for whom comorbidities, such as hypertension, diabetes, and obesity are common [Ref. 5.4: 057XM4]. The presence of comorbidities has been associated with higher risk of aggressive tumor features, recurrence, and death [Ref. 5.4: 08G0H3] [Ref. 5.4: 08G0J9].

MSI-H/ dMMR Solid tumors

Incidence and prevalence:

The prevalence of MSI-H/dMMR status varies widely by tumor type and disease stage [Ref. 5.4: 04TCGC] [Ref. 5.4: 04TCK9]. The highest prevalence of MSI-H/dMMR is found in endometrial (20-30%) and gastric (20%) cancers [Ref. 5.4: 04TCGC] [Ref. 5.4: 04TCK9] [Ref. 5.4: 04Q8R4]. MSI-H/dMMR prevalence in small intestinal cancer is about 10% [Ref. 5.4: 04SRJK] [Ref. 5.4: 05PJBW] [Ref. 5.4: 05P545] [Ref. 5.4: 05PBJJ]. For most other cancers, the prevalence of MSI-H/dMMR is well below 5% [Ref. 5.4: 04TCGC] [Ref. 5.4: 04TCK9] [Ref. 5.4: 04Q8R4]. Both pancreatic cancer and biliary cancer have a very low MSI-H/dMMR prevalence of <3% [Ref. 5.4: 05PBJJ] [Ref. 5.4: 04TCGC] [Ref. 5.4: 04SRJK] [Ref. 5.4: 04SRJK] [Ref. 5.4: 05P545] [Ref. 5.4: 05PJ9H] [Ref. 5.4: 05PJ0F]. MSI-H/dMMR prevalence is generally lower for more advanced stage tumors compared to low stage disease [Ref. 5.4: 04Q8R4].

Demographics of the indication population and risk factors for the disease:

Among patients with solid tumors, MSI-H/dMMR frequency generally increases with age and is higher for females [Ref. 5.4: 04VW8W]. Lynch Syndrome results from germline mutations in mismatch repair genes and accounts for a small portion of MSI-H/dMMR cancers; the majority of MSI-H/dMMR tumors arise in the sporadic form [Ref. 5.4: 04VWCZ]. Few consistent differences are observed for MSI-H/dMMR cancer patients compared to those with low MSI or MSS tumors with regard to common solid tumor risk factors [Ref. 5.4: 04VWCZ] [Ref. 5.4: 04VW73].

The main existing treatment options:

Pembrolizumab was approved in the US to treat unresectable or metastatic MSI-H and dMMR solid tumors that have progressed after prior treatment and for patients with no satisfactory

alternative treatment options. Otherwise, MSI-H/dMMR solid tumors are managed with cancer-specific standard-of-care chemotherapeutic treatments regardless of MSI status [Ref. 5.4: 05DMLF]. Routine testing for MMR deficiency or MSI status is recommended for all patients with endometrial (epithelial only) cancers or patients with other select solid tumors who may be candidates for anti-PD-1 treatment [Ref. 5.4: 05DMDS].

Natural history of the indicated condition, including mortality and morbidity:

The prognostic effect of MSI-H/dMMR status varies by tumor type. MSI-H/dMMR gastric cancer patients generally have an overall favorable prognosis including those with advanced disease, possibly due to these tumors having a lower ability to invade serosal layers and spread through the lymphatic system [Ref. 5.4: 04VWCZ]. In contrast, the relationship between MSI/MMR status and disease stage is less consistent for endometrial tumors; MSI-H/dMMR endometrial tumors are often associated with poorer survival [Ref. 5.4: 04FR9W].

Important co-morbidities:

Co-morbidity prevalence among MSI-H/dMMR cancer patients is unknown. However, co-morbidities are common among cancer patients, particularly with older adults for whom cancer is most common [Ref. 5.4: 04VW4L]. Secondary tumors or multiple primary tumors have been reported among MMR deficient and MSI-H endometrial cancer patients [Ref. 5.4: 04W654] [Ref. 5.4: 04VZ03].

Cervical Cancer

Incidence and Prevalence:

Over 85% of the world's 604,000 new cervical cancer cases in 2020 occurred in developing countries [Ref. 5.4: 085XNF, 085XN9]. In Europe, incidence rate (per 100,000 women) ranges from 7.0 in the Western region to 14.5 in Central/Eastern countries, yielding roughly 58,000 new cases in 2020 [Ref. 5.4: 085XNF]. In the US, incidence is 7.7 per 100,000 females, corresponding to 13,960 new cases in 2023 [Ref. 5.4: 08CJK3]. The 5-year prevalence in Europe is 172,721 cases [Ref. 5.4: 085XNF].

Demographics of the indication population and risk factors for the disease:

In Europe, 57% of cases occur before age 55 years compared to 60% in the US [Ref. 5.4: 086YC8] [Ref. 5.4: 086YC9] [Ref. 5.4: 085XN3]. The primary risk factor is persistent infection with certain types of human papillomavirus (HPV) [Ref. 5.4: 04YDD6]. In most developed countries, incidence has significantly decreased (as much as 65%) over the last four decades with the widespread use of cervical Papanicolaou (Pap) screening and the decline is expected to continue with the recent widespread implementation of the preventive HPV vaccine [Ref. 5.4: 085XN9]. In contrast, cervical cancer burden remains high in many low and middle-income countries, including parts of Central and Eastern Europe, without established vaccination, screening, and treatment programs [Ref. 5.4: 04YDDS] [Ref. 5.4: 086YCC].

The main existing treatment options:

The primary treatment of early-stage cervical cancer is surgery or radiotherapy [Ref. 5.4: 08BPP2]. Concurrent chemo-radiotherapy with cisplatin as chemosensitizer and external beam radiotherapy (EBRT) plus brachytherapy boost is the standard of care for locally advanced disease. For advanced disease, primary treatment is usually platinum-containing chemotherapy. Targeted therapies, including pembrolizumab (in the US) and bevacizumab (in the EU and US), have an established role for selected metastatic cases. Emerging therapies under investigation for recurrent disease include immunotherapy, antibody-drug conjugates and vaccines, and those targeting PIK3CA [Ref. 5.4: 085XMZ].

Natural history of the indicated condition, including mortality and morbidity:

Cervical cancer mortality is highly preventable with access to early screening and treatment [Ref. 5.4: 085XN9]. In Europe, mortality rates (per 100,000) ranges from 2.0 (Western) to 6.1 (Central/Eastern) with nearly 26,000 deaths in 2020 [Ref. 5.4: 085XNF]. The overall cervical cancer 5-year survival in the US and Europe is 67% and 62%, respectively [Ref. 5.4: 08CJK3] [Ref. 5.4: 04QRV7].

Important co-morbidities:

The prevalence of co-morbidities among cervical cancer patients is relatively low, possibly due to relatively earlier diagnosis and younger average age at diagnosis compared to other cancers. Cohort studies in Europe reported 80-90% of patients having no recorded comorbidity [Ref. 5.4: 04YVTB] [Ref. 5.4: 04YVTL].

Relevant adverse events in target populations:

Relevant adverse events to be anticipated in the aforementioned target populations exposed to immunotherapy are summarized below. The adverse events selected for inclusion are aligned with the Summary of Product Characteristics (SmPC) section 4.8.

Biliary Tract Carcinoma (BTC)

Incidence and Prevalence:

BTC arises from the epithelial lining of the biliary tree and is comprised by cancers of the intrahepatic or extrahepatic bile ducts (or cholangiocarcinoma, CCA), and gallbladder. Globally, the prevalence of BTC has risen by 35% from 2007 to 2017, resulting in 235,900 prevalent cases [Ref. 5.4: 05FVMP]. In 2017, the global incidence rate for BTC was 2.71 per 100,000 population, with approximately 210,878 new cases [Ref. 5.4: 07Z66G]. Incidence (per 100,000) is highest in Chile (10.38) and generally lowest (<1) in African or Middle Eastern countries. In Europe, incidence (per 100,000) is higher in Eastern (6.3), Southern (5.4), and Central (4.3) regions compared to Northern regions (3.4) [Ref. 5.4: 07Z66C]. Intrahepatic CCA is on the rise in the US and Europe (partly due to changes in diagnosis and classification), while gallbladder cancer is declining [Ref. 5.4: 05PFKM, 07Z66H].

Demographics of the indication population and risk factors for the disease:

In Europe and the US, BTC is more common in women; incidence is higher for men in some regions of Asia [Ref. 5.4: 07Z667, 07Z66C, 07Z66G]. Many risk factors have been identified, though the relative strength of these factors varies somewhat by BTC subtype. These include bile duct cysts, cholangitis, cholelithiasis, cirrhosis, HBV/HCV, nonalcoholic fatty liver disease (NAFLD)/nonalcoholic steatohepatitis (NASH), chronic pancreatitis, inflammatory bowel disease, *H pylori* infection, obesity, liver fluke infections, and some occupational exposures or genetic factors; however, roughly half of cases are diagnosed without any identifiable risk factors [Ref. 5.4: 05PFKM, 07Z66H].

The main existing treatment options:

Surgical procedures or radiation may be options for localized BTC [Ref. 5.4: 07Z655]. Liver transplantation is potentially curative for select patients with locally advanced disease. Treatment options for advanced BTC include a combination of durvalumab plus gemcitabine and cisplatin-based chemotherapy, gemcitabine- or fluoropyrimidine-based chemotherapy, pembrolizumab for MSI-H/dMMR tumors, FGFR2 inhibitors, IDH inhibitors fluoropyrimidine-based chemoradiation, or radiation alone. Potential molecular targets for emerging therapies to treat BTC subtypes include NRAS, BAP1, ARID1A, EGFR, NTRK, BRAF V600E, KRAS, HER2, PIK3C2G, STK11, TP53, and TGFBR2 [Ref. 5.4: 0885WD, 0885WH, 0885WN, 0885WP].

Natural history of the indicated condition, including mortality and morbidity:

Globally, mortality rates (per 100,000) vary widely from 0.8 in South Africa to 21.2 in Chile [Ref. 5.4: 05H5SD]. Survival is similarly poor for CCA and gallbladder cancer with a median survival under 1 year and 5-year survival rates of 10%-20% in Europe and the US [Ref. 5.4: 07Z66C, 07Z653]. Most BTC is diagnosed at a late stage when treatment is less effective [Ref. 5.4: 07Z667].

Important co-morbidities:

Comorbidities are common as most patients are elderly. A study of 86,134 BTC cases found cholelithiasis, liver disease, or diabetes in 18%, 19%, and 33%, respectively [Ref. 5.4: 07Z669].

Gastric Cancer

Incidence and prevalence:

Approximately 70% of the world's 968,784 annual new gastric cancer (GC) cases occur in Asia [Ref. 5.4: 06D898][Ref. 5.4: 08JJK6]. Incidence varies globally, with the highest age-standardized incidence rates (per 100,000) found in Eastern Asia (16.1); rates are lowest in Southern Africa (3.4)[Ref. 5.4: 08JJK6]. In Europe, incidence (per 100,000 PY) ranges from 4.3 in Northern countries to 7.2 in the Eastern region, resulting in an estimated 135,610 incident cases in 2022 [Ref. 5.4: 08JJK6]. In the US, age-standardized incidence is 7.0 per 100,000,

corresponding to 26,890 new cases expected in 2024[Ref. 5.4: 08389Q][Ref. 5.4: 08KSQF]. The 5-year prevalence in Europe is 188,578 cases[Ref. 5.4: 08JJK6].

Demographics of the indication population and risk factors for the disease:

Incidence increases with age, with incidence peaking in those over age 70 years [Ref. 5.4: 08097V]. Males have a roughly 2--fold higher incidence compared to females in the US and Europe[Ref. 5.4: 08KSQF][Ref. 5.4: 08JJK6]. The primary risk factor for non-cardia (distal) GC (comprising a majority of cases) is *Helicobacter pylori* gastritis; the reduction in *H. pylori* prevalence through eradication efforts is significantly contributing to the decline in GC incidence in the US and Europe [Ref. 5.4: 06D8FH]. The less common cardia (proximal) subtype is suspected to be related to obesity, smoking, and gastro-esophageal reflux; incidence is increasing in the US and Europe [Ref. 5.4: 06D8FH].

The main existing treatment options:

Surgery is the primary treatment option for localized disease[Ref. 5.4: 08LJMC]. Combined modality therapy is recommended for locoregional disease; options include peri- or post-operative chemotherapy and pre- or post-operative chemoradiation. For unresectable, locally advanced and metastatic disease, first-line systemic therapy regimens with two cytotoxic drugs are preferred. A fluoropyrimidine combined with oxaliplatin is recommended for most patients who are HER2-negative, whereas trastuzumab plus chemotherapy is indicated for HER2-positive metastatic disease. With recent options to add immunotherapy to either regimen based on approvals of nivolumab and pembrolizumab in HER2-negative disease and pembrolizumab in HER2-positive disease, subsequent lines of therapy may include ramucirumab with or without paclitaxel; single-agent chemotherapy; dostarlimab (dMMR tumors); pembrolizumab (MSI-H/dMMR or TMB-high tumors); entrectinib or larotrectinib (NTRK gene fusion-positive tumors); or trifluridine and tipiracil (recurrent/metastatic tumors).

Natural history of the indicated condition, including mortality and morbidity:

Mortality rates (per 100,000) are highest in Eastern Asia (9.2) and lowest in North America (1.7); rates in Europe range from 2.8 in the North to 7.8 in the Eastern region[Ref. 5.4: 08JJK6]. In Europe, overall 5-year survival ranges from 18.3% to 35.4% [Ref. 5.4: 08097L]. Most US cases are diagnosed at regional (25%) or distant (37%) stage, contributing to a relatively poor 5-year survival (36.4%)[Ref. 5.4: 08KSQF].

Important co-morbidities:

A large study of 12,612 GC patients reported a high prevalence of co-morbidities prior to diagnosis; the most common chronic conditions were hypertension (60%), coronary artery disease (30%), diabetes (23.6%), COPD (17%), congestive heart failure (16.4%), and osteoarthritis (13.6%) [Ref. 5.4: 04KCJW].

Malignant Pleural Mesothelioma (MPM)

Incidence and prevalence:

MPM is a rare, aggressive tumor that arises in the mesothelial lining of the pleura and accounts for roughly 85%-90% of all mesotheliomas [Ref. 5.4: 04LJ0D, 088TP2]. Incidence is widely considered to be under-estimated and countries with the highest incidence are those with a long tradition of studying the disease epidemiology and patterns of asbestos use [Ref. 5.4: 088RW8]. Reported annual incidence (per 100,000 population) for all mesothelioma varied between 0.15 in East Asia to 2.13 in Australasia; in the Central, Eastern, and Western regions of Europe, incidence was 0.42, 0.38, and 1.31, respectively [Ref. 5.4: 088RWP]. In the U.S., MPM incidence is 0.6 per 100,000 population, resulting in over 2,000 incident cases annually [Ref. 5.4: 088TMY]. In the pediatric population, MPM is extremely rare and has only been anecdotally described in the literature. The age-adjusted incidence rate of mesothelioma is 0.00 per 100,000 in the UK, the EU-27 and worldwide with a total of 0 estimated incident pediatric mesothelioma cases in 2022 for the UK, 5 estimated incident cases for the EU-27, and 49 estimated incident cases worldwide [Ref. 5.4: 08HXKC, 08HXK6, 08HXK0].

Demographics of the indication population and risk factors for the disease:

More than 70% of MPM cases arise in men [Ref. 5.4: 088RWP]. In the US, incidence is higher for Whites and Hispanics compared to African Americans or Asian Americans [Ref. 5.4: 088TRN]. Roughly 86% of cases worldwide occur in patients 50 years or older and 45% are over age 70 years [Ref. 5.4: 088RWP]. Asbestos is the main risk factor, most commonly through occupational exposure involved with manufacturing or installation of asbestos products [Ref. 5.4: 088RW8]. MPM is associated with a long latency period (20-60 years) following asbestos exposure [Ref. 5.4: 088R34]. Although asbestos is banned in more than 60 countries (including most of Europe), leading to modest MPM declines in some countries around two decades following a ban, MPM continues to occur due to the late disease onset and remaining exposure to asbestos in some older buildings [Ref. 5.4: 088RWP, 088R4G]. Furthermore, several countries continue exporting some asbestos products and use is increasing in many low- or middle-income countries, contributing to the global MPM burden [Ref. 5.4: 088R4G]. Other risk factors include radiation exposure, polio vaccine contamination with Simian virus 40 between 1950-1970, and specific genetic mutations [Ref. 5.4: 088R4G, 088R69].

The main existing treatment options:

A trimodality regimen with chemotherapy, surgery and hemi-thoracic radiotherapy is recommended for operable MPM [Ref. 5.4: 088TP2]. For inoperable MPM, recommended first-line treatments include nivolumab plus ipilimumab, and cisplatin (or carboplatin) plus pemetrexed (with or without bevacizumab). Subsequent treatment may include vinorelbine, gemcitabine or Nivolumab +/- ipilimumab (if not given in first line). Emerging treatments under investigation include additional immune checkpoint inhibitors, oncolytic viruses, cancer vaccines, and adoptive T cell therapy [Ref. 5.4: 088R5X].

Based on small case series studies, a therapeutic approach similar to adults has been adopted for children with mesothelioma, which may include surgery, chemotherapy with cisplatin and pemetrexed, and hyperthermic intraperitoneal chemotherapy (HIPEC) [Ref. 5.4: 0880L0, 08J9LT, 08J9N7].

Natural history of the indicated condition, including mortality and morbidity:

MPM is highly fatal as most patients are diagnosed with advanced stage [Ref. 5.4: 088RW8]. In the US and Europe, the median survival is less than one year following diagnosis with a 5-year survival of 10% [Ref. 5.4: 088TP2, 088R4G]. Non-epithelioid histology, increasing age, and male gender are independent risk factors for poor prognosis [Ref. 5.4: 088R4G]. A WHO analysis estimated that global mesothelioma deaths were approximately 38,400 per year [Ref. 5.4: 088R4G].

Important co-morbidities:

Data regarding the prevalence of comorbidities among MPM patients is scarce. Studies of MPM cohorts in the US found 30-45% of patients to have a Charlson Comorbidity Index score of one or higher [Ref. 5.4: 05DPTW, 088R5H].

Table SI.1: Background Frequencies of Relevant Adverse Events in Target Populations

Adverse Event	Frequency*
Immune-mediated Pneumonitis	Pneumonitis occurred in 3.1% of 1994 patients treated with single-agent nivolumab, 6.4% of 1291 patients treated with combined ipilimumab and nivolumab, 3% of 2616 patients treated with atezolizumab, 2% of 1414 patients without prior radiation or 16.6% in 475 patients with radiation and treated with durvalumab, 1.2% of 1738 patients treated with avelumab, 3.2% of 810 patients treated with cemiplimab, and 1.1% of 444 patients receiving dostarlimab.
Immune-mediated Colitis	Colitis occurred in 2.9% of 1994 patients treated with single-agent nivolumab, 21.1% of 982 patients treated with single-agent ipilimumab, 9% of 1291 patients treated with combined nivolumab and IPI, 1.6% of 1889 patients treated with durvalumab, 1.5% of 1738 patients treated with avelumab, and 1% of 2616 patients treated with atezolizumab, and 2.2% of 810 patients treated with cemiplimab, and 1.4% of 444 patients receiving dostarlimab.
Immune-mediated Hepatitis	Hepatitis occurred in 1.8% of 1994 patients treated with single-agent nivolumab, 9.3% of 982 patients treated with single-agent ipilimumab 7.9% in 715 patients treated with combined nivolumab and IPI, 1% of 1889 patients treated with durvalumab, 0.9% of 1738 patients treated with avelumab, 1.8% of 2616 patients treated with atezolizumab, 2.0% of 810 patients treated with cemiplimab, and 0.2% of 444 patients receiving dostarlimab.
Immune-mediated Nephritis	Nephritis and renal dysfunction occurred in 1.2% of 1994 patients treated with single-agent nivolumab, 0.1% of 1738 patients treated with avelumab, 0.3% of 1889 patients treated with durvalumab, <1% of 666 patients treated with ipilimumab (IPI), and 0.6% of 810 patients treated with cemiplimab, and 0.5% of 444 patients receiving dostarlimab.

Table SI.1: Background Frequencies of Relevant Adverse Events in Target Populations

Adverse Event	Frequency*
Immune-mediated Endocrinopathies- Hypophysitis (including hypopituitarism)	Hypophysitis occurred in 0.6% of 1994 patients treated with single-agent nivolumab; hypophysitis occurred in 4.4% of 715 patients with combined ipilimumab and nivolumab; hypophysitis occurred in <0.1% of 2616 patients treated with atezolizumab; hypopituitarism occurred in 0.1% of 1738 patients treated with avelumab, and 0.4% in 810 patients treated with cemiplimab.
Immune-mediated Endocrinopathies- Adrenal insufficiency (primary and secondary)	Adrenal insufficiency occurred in 1% of 1994 patients treated with single-agent nivolumab; adrenal insufficiency occurred in 7.8% of 715 patients treated with combined ipilimumab and nivolumab; adrenal insufficiency was reported in 0.4% of 2616 patients treated with atezolizumab; adrenal insufficiency occurred in 0.5% of 1738 patients treated with avelumab; adrenal insufficiency occurred in 0.7% of 1889 patients treated with durvalumab; adrenal insufficiency occurred in 0.4% in 810 patients treated with cemiplimab; and in 0.9% of 444 patients receiving dostarlimab.
Immune-mediated Endocrinopathies- Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)	Hypothyroidism, hyperthyroidism, and thyroiditis occurred in 8%, 2.7%, and 0.6%, respectively, of 1994 patients treated with single-agent nivolumab; 18.3%, 11.9%, and 2.7% of 715 patients treated with combined IPI and nivolumab; 7.3%, 1.4%, and 0.4%, respectively, of 1889 patients treated with durvalumab; 4.9%, 0.8%, and 0.2%, respectively, of 2616 patients treated with atezolizumab; 5%, 0.4%, and 0.2% respectively, of 1738 patients treated with avelumab; 7%, 3.2%, and 0.6%, respectively, of 810 patients treated with cemiplimab; 5.6%, 1.8%, and 0.5%, respectively, of 444 patients treated with dostarlimab.
Immune-mediated Endocrinopathies- Type 1 Diabetes Mellitus	Diabetes occurred in 0.9% of 1994 patients treated with single-agent nivolumab, 2.7% of 666 patients treated with combined IPI and nivolumab, 0.1% of 1738 patients treated with avelumab; <0.1% of 1889 patients treated with durvalumab, 0.3% of 2616 patients treated with atezolizumab, and 0.1% in 810 patients treated with cemiplimab.
For hematologic malignancies: increased risk of severe complications of allogeneic stem cell transplantation (SCT)	<p>Graft versus host disease (GVHD)</p> <ul style="list-style-type: none"> In 18 studies of patients treated with allogeneic transplant, without PD-1 inhibitors, the median and range of incidences (overall or subgroup-specific) reported for Grades 2-4 acute GVHD, Grades 3-4 acute GVHD, and chronic GVHD were 38.2% (11.5%-61%), 15.9% (8%-22%), 41.5% (5%-56%), respectively [Ref. 5.4: 04FF3L, 04FH09, 04FH0G, 04FG37, 04FJTb, 04FF00, 04FF0J, 04FFX4, 04FG2L, 04FF27, 04FGZW, 04FJ33, 04FGZP, 04FG2J, 04FJSZ, 04FJT7, 04FJTD, 04FG08]. <p>Veno-occlusive Disease (VOD)</p> <ul style="list-style-type: none"> 8.8% VOD was seen among 843 allogeneic Hematopoietic Stem Cell Transplant (HSCT) procedures in 763 lymphoma/leukemia patients in Spain [Ref. 5.4: 04FKV7]. Hepatic VOD was diagnosed in 2.9% of 1326 patients who received allogeneic HSCT and were included in a nationwide HSCT registry in Taiwan from 2009-14 [Ref. 5.4: 05G7NG]. 1.6% of 1583 patients undergoing reduced-intensity conditioning allogeneic HSCT had developed VOD/sinusoidal obstruction syndrome from 2007-17 at the Dana Farber Cancer Institute [Ref. 5.4: 05G7NH].
Graft versus host disease (GVHD) in patients with a history of allogeneic stem cell transplant (SCT)	A comprehensive literature review of 13 case reports and 11 original manuscripts described GVHD (acute and chronic) in 283 patients with hematologic malignancies who received checkpoint inhibitors (CPI), including ipilimumab, nivolumab or pembrolizumab, either before or after having failed allogeneic hematopoietic stem cell transplantation (allo-HSCT) [Ref. 5.4: 05G88Y]. Among the 150 patients receiving a CPI after undergoing allo-HSCT (in the original articles), 62 received nivolumab, 85 had

Table SI.1: Background Frequencies of Relevant Adverse Events in Target Populations

Adverse Event	Frequency*
	<p>ipilimumab, and 4 had pembrolizumab. Of these patients who received CPI post-allo-HSCT, acute GVHD and chronic GVHD developed in 13% and 11%, respectively; 49% had a prior history of GVHD. Among the 26 patients receiving CPI post-allo-HSCT (as part of the case reports and case series), 12 received pembrolizumab and 14 had nivolumab. 19% of these patients developed acute GVHD, and 4% had a flare of existing GVHD.</p>
<p>* Except where otherwise noted, frequencies for the immune-mediated AEs are based on US Prescribing information [Ref. 5.4: 05DMN4, 05DMN5, 05DMN6, 05DNDC, 05DP3H] [Ref. 5.4: 06CLG4] [Ref. 5.4: 06CLG5] for similar class agents.</p>	

PART II: MODULE SII - NON-CLINICAL PART OF THE SAFETY SPECIFICATION

Key safety findings from non-clinical studies and relevance to human usage:

Table SII.1: Summary of Important Safety Findings from Non-clinical Studies

Key Safety Findings (from non-clinical studies)	Relevance to Human Usage
<p>Repeat-dose toxicity: None</p> <p>Pembrolizumab (MK-3475) was well-tolerated when administered intravenously (IV) to Cynomolgus monkeys at doses of 6, 40 and 200 mg/kg during the 1- and 6-month toxicology studies. There were no findings of toxicological significance and the NOAEL was 200 mg/kg in both studies. The exposure multiple based on a predicted AUC_{0-tau} of 913.2 day* µg./mL at the clinical dose of 200 mg every 3 weeks (Q3W) and/or 400 mg every 6 weeks (Q6W) was 74-fold at the dose of 200 mg/kg based on the 6-month study.</p>	<p>No safety concerns for repeated IV administration.</p>
<p>Pregnancy: Based upon data from the literature on the role of the PD-1/PD-L1 pathway in maintaining pregnancy and on experimental results in murine models of allogeneic pregnancy a potential risk with the administration of pembrolizumab to women of childbearing potential has been clearly identified. Because the PD-1/PD-L1 pathway plays a fundamental role in maintaining immune tolerance to the fetus, it is anticipated that the inhibition of PD-1 during pregnancy by treatment with pembrolizumab will have adverse effects on the fetus, including increased risks of abortion and stillbirth [Ref. 5.4: 00VMR9, 02CFKR, 02CFKT, 00VMXH].</p>	<p>Due to its mechanism of action, pembrolizumab may increase the risk of abortion and stillbirths if administered during pregnancy.</p>
<p>Immune-mediated disease: Based on the mechanism of action of pembrolizumab a safety risk for immune-mediated adverse events has been identified. The PD-1/PD-L1 pathway plays an important role in mediating peripheral immune tolerance to self, therefore its inhibition may contribute to the development of immune-mediated adverse events. [Ref. 5.4: 02CFKV, 02CFKW, 02CFKX, 02CFKQ]. However in a T cell-dependent Antibody Response (TDAR) study in mice, neither immune-mediated toxicity nor effects on the magnitude and kinetics of the primary and recall antibody responses to Hepatitis B vaccine were observed during treatment with anti-PD-1 antibody in comparison to untreated control group [Ref. 4.2.3.7.2: TT151105FIN].</p>	<p>The potential development of immune-mediated adverse events is a known risk with pembrolizumab and is discussed in Section SVII.3.1.</p>

Table SII.1: Summary of Important Safety Findings from Non-clinical Studies

Key Safety Findings (from non-clinical studies)	Relevance to Human Usage
<p>Local Tolerance: None</p> <p>MK-3475A (containing pembrolizumab and berahyaluronidase alfa, a permeation enhancer) was well tolerated when administered subcutaneously (SC) to Cynomolgus monkeys at a dose of 50 mg/kg pembrolizumab and 0.0041 mg/kg (approximately 600 U/kg) berahyaluronidase alfa in a 4-week repeat-dose tolerability study. MK-3475A showed acceptable local tolerability at the SC injection sites and draining lymph nodes. The exposure multiples for pembrolizumab administered SC as MK-3475A are 6-fold, based on a predicted AUC_{0-3weeks} at steady state of 1439 day*µg/mL at the anticipated Q3W clinical dose of 395 mg pembrolizumab and 0.03 mg berahyaluronidase alfa [4800 U], and 3-fold, based on AUC_{0-6weeks} at steady state of 2962 day*µg/mL at the anticipated Q6W clinical dose of 790 mg pembrolizumab + 0.06 mg berahyaluronidase alfa [9600 U]. Berahyaluronidase alfa had negligible systemic exposure.</p>	<p>No safety concerns for repeated SC administration.</p>

Conclusions on Non-clinical Data

Nonclinical toxicology studies in monkeys of up to 6-months in duration with pembrolizumab demonstrated that there were no adverse effects. The exposure multiple between the NOAEL and a human IV dose of 200 mg Q3W or 400 mg Q6W was 74. Repeat-dose local tolerance study in monkeys up to 4-weeks in duration with pembrolizumab administered SC as MK-3475A, which contains berahyaluronidase alfa as a permeation enhancer, showed MK-3475A was well tolerated with acceptable local tolerability at the SC injection sites at exposure multiples of 6 and 3 times the exposures at a human dose of 395 mg pembrolizumab and 0.03 mg (4800 U) berahyaluronidase alfa administered Q3W and 790 mg pembrolizumab and 0.06 mg (9600 U) berahyaluronidase alfa administered Q6W, respectively. No reproductive toxicity studies were conducted with pembrolizumab or MK-3475A based upon the potential risk to pregnancy that is clearly identified from experimental results in the literature. Based on the mechanism of action of pembrolizumab, there is a potential risk of developing immune-mediated adverse events.

PART II: MODULE SIII - CLINICAL TRIAL EXPOSURE

The clinical trial exposure is presented only for the clinical trial datasets/indications that are included in this RMP. The clinical trials listed in the RMP consist of trials which utilized pembrolizumab administered intravenously and subcutaneously. All studies were conducted with administration of pembrolizumab IV, unless noted as MK-3475A subcutaneous.

Clinical Trial Exposure by Duration of Exposure

KEYNOTE-001 (KN001) is a multi-cohort clinical trial evaluating the safety, tolerability, pharmacokinetics (PK) and pharmacodynamics (PD), and anti-tumor activity of pembrolizumab in patients with melanoma (naïve to or previously treated with IPI (IPI)) and Non-Small Cell Lung Cancer (NSCLC). The “All Melanoma” population consisted of cohorts B1 (IPI naïve or IPI treated), B2 (IPI refractory), B3 (IPI naïve or IPI treated), and D (IPI naïve). The NSCLC population consisted of cohort C (2 prior systemic therapy regimens) and cohort F (naïve to treatment, or, had one or more prior systemic treatment regimens). Study visits were every 2 or 3 weeks, dependent on dosing cycles, and patients remained on drug until progressive disease or other protocol defined stopping criteria occurred (e.g., as intercurrent illness that prevented further administration of treatment; or the patient withdrew consent).

KEYNOTE-002 (KN002) is a randomized, Phase II study of pembrolizumab compared to chemotherapy in patients with advanced melanoma. Study visits were every 2 or 3 weeks, dependent on dosing cycles, and patients remained on drug until progressive disease or other protocol defined stopping criteria occurred.

KEYNOTE-006 (KN006) is a multicenter, randomized, controlled, open-label, three-arm, Phase III trial designed to evaluate the safety and efficacy of two dosing regimens of pembrolizumab compared to IPI in patients with unresectable or metastatic advanced melanoma who are IPI-naïve. Subjects randomized to either of the pembrolizumab treatment arms received treatment for up to 24 months or until disease progression or other protocol defined stopping criteria occur.

KEYNOTE-671 (KN671) is a multi-center Phase III, randomized, double-blind study of platinum-containing chemotherapy with or without pembrolizumab as neoadjuvant/adjuvant therapy for participants with resectable (American Joint Committee on Cancer [AJCC] 8th edition) Stage II, IIIA, and resectable IIIB (T3-N2) NSCLC. Participants were randomized to receive concomitant neoadjuvant platinum-containing chemotherapy plus pembrolizumab (Q3W × 4 cycles) followed by surgery and adjuvant pembrolizumab (Q3W × 13 cycles) or concomitant neoadjuvant platinum containing chemotherapy plus placebo (Q3W × 4 cycles) followed by surgery and adjuvant placebo (Q3W × 13 cycles), and followed up until death, disease recurrence/progression.

KEYNOTE-091 (KN091) is a multi-center, randomized double blind, placebo-controlled Phase III trial of pembrolizumab monotherapy following resection of stages IB-IIIa non-small cell lung cancer (NSCLC) American Joint Committee on Cancer (AJCC) 7th edition. Adjuvant chemotherapy is not mandatory but considered for patients with stage IB (T ≥ 4 cm) and

strongly recommended for stage II and IIIA, according to national and local guidelines. Participants are randomized to receive 18 cycles of pembrolizumab or placebo every 3 weeks and followed until death, lung cancer recurrence, occurrence of a second primary lung cancer or occurrence of a second primary cancer. KEYNOTE-010 (KN010) is a multi-center, worldwide, randomized, adaptively designed Phase II/III trial of pembrolizumab at two dosing schedules versus docetaxel in subjects with NSCLC with PD-L1 positive tumors who have experienced disease progression after platinum-containing systemic therapy. Subjects randomized to either of the pembrolizumab treatment arms or docetaxel received treatment for up to 24 months or until disease progression or other protocol defined stopping criteria occur.

KEYNOTE-021 (KN021) is a multi-center, open-label, Phase I/II study of pembrolizumab in combination with chemotherapy or immunotherapy in subjects with advanced/metastatic NSCLC. The study consists of two parts with multiple cohorts; cohort A of Part 1 of the study evaluated pembrolizumab in combination with carboplatin-paclitaxel in subjects with metastatic squamous NSCLC who have not previously received systemic therapy for metastatic disease. Subjects were randomized 1:1 to either receive 2mg/kg of pembrolizumab or 10mg/kg combined with carboplatin and paclitaxel. All subjects received up to a maximum of 4 cycles of carboplatin and paclitaxel. Treatment with pembrolizumab continued until a total of 35 treatment administrations or until disease progression or other protocol defined stopping criteria. Cohort C of Part 1 of the study determined the recommended Phase II dosage (RP2D) of pembrolizumab in combination with carboplatin and pemetrexed [standard of care (SOC)]. Part 2 of the study included a randomized comparison of carboplatin and pemetrexed with or without pembrolizumab (Cohort G). Subjects randomized to the combination received 4 cycles of carboplatin and continued maintenance pemetrexed with pembrolizumab for up to 35 cycles or until disease progression or other protocol defined stopping criteria. Subjects randomized to SOC received 4 cycles of carboplatin and continued maintenance pemetrexed until disease progression or other protocol defined stopping criteria.

KEYNOTE-189 (KN189) is a worldwide, multi-site, randomized, double-blind, parallel-group, placebo-controlled study to evaluate intravenous pembrolizumab combined with pemetrexed/platinum chemotherapy versus saline placebo combined with pemetrexed/platinum chemotherapy in subjects with advanced or metastatic non-squamous NSCLC who have not previously received systemic therapy for advanced/metastatic disease. Subjects were randomized 2:1 to either receive pembrolizumab at 200 mg combined with pemetrexed and platinum (investigators choice of cisplatin or carboplatin), or saline placebo, pemetrexed and platinum (investigators choice of cisplatin or carboplatin). Subjects randomized to the combination received 4 cycles of chemotherapy and continued maintenance pemetrexed with pembrolizumab for up to 35 cycles or until disease progression or other protocol defined stopping criteria. Subjects randomized to SOC received 4 cycles of chemotherapy and continued maintenance pemetrexed until disease progression or other protocol defined stopping criteria.

KEYNOTE-024 (KN024) is a multi-center, worldwide, randomized, open label, controlled trial of intravenous (IV) pembrolizumab monotherapy versus the choice of multiple standard of care (SOC) platinum-based chemotherapies in subjects previously untreated for their stage IV, PD-L1 strong, NSCLC. Subjects randomized to pembrolizumab received up to 35 treatments or until disease progression or other protocol defined stopping criteria. Subjects

randomized to SOC received the recommended SOC treatments for the selected platinum-based chemotherapies or until disease progression or other protocol defined stopping criteria.

KEYNOTE-042 (KN042) is a Phase 3 multi-center, international, randomized, open-label, controlled clinical study of pembrolizumab monotherapy versus platinum-based chemotherapy in previously untreated adult subjects with locally advanced or metastatic Tumor Proportion Score (TPS) $\geq 1\%$ NSCLC, with no EGFR or ALK genomic tumor aberrations. Subjects randomized to pembrolizumab received up to 35 treatments or until disease progression or other protocol defined stopping criteria. Subjects randomized to platinum-based chemotherapy received the investigator's choice of carboplatin/paclitaxel or carboplatin/pemetrexed for up to 6 cycles followed by optional pemetrexed maintenance, or until disease progression or other protocol defined stopping criteria. While pemetrexed maintenance was optional, it was strongly recommended for subjects with nonsquamous histologies only.

KEYNOTE-407 (KN407) is a worldwide, multi-site, randomized, double-blind, parallel-group, placebo-controlled study to evaluate intravenous pembrolizumab combined with carboplatin-paclitaxel/nab-paclitaxel chemotherapy versus saline placebo combined with carboplatin-paclitaxel/nab-paclitaxel chemotherapy in subjects with metastatic squamous NSCLC who have not previously received systemic therapy for metastatic disease. Subjects were randomized 1:1 to either receive pembrolizumab combined with carboplatin and investigators choice of paclitaxel or nab-paclitaxel, or saline placebo combined with carboplatin and investigators choice of paclitaxel or nab-paclitaxel. All subjects received up to a maximum of 4 cycles of carboplatin, paclitaxel or nab-paclitaxel and pembrolizumab or saline placebo. Treatment with pembrolizumab or saline placebo continued until a total of 35 treatment administrations or until disease progression or other protocol defined stopping criteria.

MK-3475A-D77 is an open-label, multi-center, Phase 3 randomized study that evaluates the pharmacokinetics (PK) and safety of MK-3475A [pembrolizumab with berahyaluronidase alfa), a permeation enhancer] administered subcutaneously compared to intravenous (IV) pembrolizumab, both administered with chemotherapy for the first-line treatment of adult participants with metastatic non-small cell lung cancer (NSCLC). Subjects were randomized (2:1) to receive either pembrolizumab administered as MK-3475A 790 mg subcutaneously every 6 weeks with platinum doublet chemotherapy or pembrolizumab 400 mg intravenously every 6 weeks with platinum doublet chemotherapy. Treatment with subcutaneous MK-3475A or intravenous pembrolizumab continued until RECIST 1.1-defined progression of disease as determined by the investigator, unacceptable toxicity, or a maximum of 18 cycles (approximately 24 months). Treatment with subcutaneous MK-3475A or intravenous pembrolizumab could be reinitiated for subsequent disease progression and administered for up to an additional 9 cycles (approximately 1 year).

KEYNOTE-013 (KN013) is a multi-center, multi-cohort, nonrandomized trial of pembrolizumab in hematologic malignancies. Cohort 3 included subjects with relapsed/refractory nodular sclerosing or mixed cellularity Hodgkin lymphoma (HL) that have failed, are ineligible for, or refused a stem cell transplant and have relapsed after treatment with or failed to respond to brentuximab vedotin. Subjects in Cohort 3 received pembrolizumab 10 mg/kg administered every 2 weeks. Subjects were evaluated at Week 12, then every 8

weeks (± 7 days) to assess response to treatment according to the Revised Response Criteria for Malignant Lymphoma [Ref. 5.4: 03RC2F].

KEYNOTE-087 (KN087) is a multicenter, single arm, multi-cohort, nonrandomized trial of pembrolizumab in subjects with relapsed or refractory classical Hodgkin lymphoma who have failed to achieve a response or progressed after autologous stem cell transplant (ASCT) and have relapsed after treatment with, or failed to respond to, brentuximab vedotin post ASCT (Cohort 1); who were unable to achieve a complete response (CR) or partial response (PR) to salvage chemotherapy and did not receive ASCT, but have relapsed after treatment with, or failed to respond to, brentuximab vedotin (Cohort 2); and subjects who have failed to respond to, or progressed after, ASCT and have not received brentuximab vedotin post ASCT. These subjects may or may not have received brentuximab vedotin as part of primary or salvage treatment (Cohort 3). Subjects received pembrolizumab 200mg every 3 weeks. Subjects were evaluated every 12 weeks (± 7 days) to assess response to treatment according to the Revised Response Criteria for Malignant Lymphoma [Ref. 5.4: 03RC2F].

KEYNOTE-204 (KN204) is a Phase III, randomized, open-label, multi-national, clinical trial of pembrolizumab compared with brentuximab vedotin (BV) in subjects with relapsed or refractory classical Hodgkin lymphoma (HL). This study enrolled subjects with relapsed or refractory classical HL, who had received at least 1 prior multi-agent chemotherapy regimen. Prior BV or a BV-containing regimen was allowed, provided the subjects responded (partial or complete response) to the BV or BV-containing regimen. Patients received study medication (200mg pembrolizumab or 1.8 mg/kg BV) every 3 weeks. Patients were evaluated every 12 weeks (± 7 days) to assess response to treatment according to the Revised Response Criteria for Malignant Lymphoma [Ref. 5.4: 03RC2F].

KEYNOTE-051 (KN051) is a Phase 1/2, nonrandomized, open-label, single-arm, multicenter study to evaluate the pharmacokinetics, pharmacodynamics, toxicity, safety, and antitumor activity of pembrolizumab in pediatric participants aged 6 months to less than 18 years of age with either advanced melanoma, advanced relapsed or refractory PD-L1 positive malignant solid tumor or other lymphoma, relapsed or refractory classical HL (3 years to 18 years of age), or advanced relapsed or refractory MSI-H solid tumors. Part I objectives are safety, PK, PD, toxicity, and preliminary efficacy in pediatric participants with advanced melanoma or PD-L1-positive advanced, relapsed or refractory solid tumor or other lymphoma. Part 2 objectives are to evaluate safety and efficacy at the established RP2D in pediatric participants within the indications of advanced melanoma; PD-L1-positive advanced, relapsed or refractory solid tumors or other lymphoma; relapsed or refractory classical Hodgkin lymphoma (rrcHL); or advanced, relapsed or refractory microsatellite instability high (MSI-H) solid tumors. Standard RECIST 1.1 criteria were used for assessment of the primary efficacy endpoint in patients with solid tumors and other lymphoma assessments: (every 8 weeks ± 7 days through Week 24 and once every 12 weeks ± 7 days thereafter) or IWG Revised Response Criteria for patients in the rrcHL Cohort (assessments: every 12 weeks ± 7 days) [Ref. 5.4: 03QY3L].

KEYNOTE-045 (KN045) is a randomized, active-controlled, multi-site, open-label trial of pembrolizumab monotherapy versus investigator's choice of paclitaxel, docetaxel or vinflunine in subjects with locally advanced or metastatic urothelial carcinoma who have received platinum-containing chemotherapy. Subjects randomized to pembrolizumab received

up to 24 months of study treatment or until disease progression or other protocol defined stopping criteria. Subjects randomized to chemotherapy received the recommended standard treatment for selected chemotherapy until disease progression or other protocol defined stopping criteria.

KEYNOTE-052 (KN052) is an open-label, single-arm, multi-site trial of pembrolizumab monotherapy in subjects with locally advanced or metastatic UC who have not received prior systemic chemotherapy and who are not eligible to receive cisplatin. Subjects may remain on treatment for 24 months, until disease progression or another protocol-specified reason for discontinuation from therapy. After discontinuation from treatment, the subject is followed for survival.

KEYNOTE-869/EV-103 (KN869) is a multi-center, multi-cohort, open-label, Phase 1b/2 trial. Three cohorts of this study are dedicated to investigating enfortumab vedotin (EV) in combination with pembrolizumab in participants with locally advanced or metastatic urothelial cancer (la/mUC) who are not eligible for cisplatin-based chemotherapy. These include a dose-escalation cohort, Cohort A (expansion cohort), and Cohort K (randomized treatment cohort). All participants in the dose escalation cohort and Cohort A received EV in combination with pembrolizumab. Participants in Cohort K were randomized in a 1:1 ratio to receive either the combination or EV monotherapy. Initial tolerability and DLT assessments of the combination were completed in the dose escalation cohort among participants who received study treatment in the first-line or second-line setting. Pembrolizumab dosing was continued up to 35 treatments (approximately 2 years); however, treatment with EV could be continued beyond 2 years until protocol defined discontinuation criteria are met.

KEYNOTE-A39/EV-302 (KNA39) is a multi-center, open-label, randomized, controlled Phase 3 study investigating EV in combination with pembrolizumab versus platinum-based chemotherapy alone in previously untreated locally advanced or metastatic urothelial cancer. Pembrolizumab dosing was continued up to 35 treatments (approximately 2 years); however, treatment with EV could be continued beyond 2 years until protocol defined discontinuation criteria are met.

KEYNOTE-040 (KN040) is a randomized, active-controlled, multicenter, open-label Phase III clinical trial to examine the efficacy and safety of pembrolizumab versus the choice of 3 different standard treatment options in subjects with recurrent or metastatic (R/M) head and neck squamous cell carcinoma (HNSCC) whose disease has progressed on or after prior platinum-containing chemotherapy. Subjects may remain on treatment for 24 months, until disease progression or another protocol-specified reason for discontinuation from therapy. After discontinuation from treatment, the subject is followed for survival.

KEYNOTE-048 (KN048) is a Phase III, randomized, active-controlled, multi-site, open-label trial of pembrolizumab, or pembrolizumab plus chemotherapy (platinum plus 5-FU) versus cetuximab plus chemotherapy in subjects with first line recurrent or metastatic (R/M) head and neck squamous cell carcinoma (HNSCC). Subjects are randomized in a 1:1:1 ratio to one of the three treatment arms in the trial. Randomization is stratified by PD-L1 tumor expression (strongly positive versus not strongly positive) defined by TPS 50% cutpoint, HPV status for oropharyngeal cancer (positive versus negative), and ECOG Performance Scale (0 versus 1).

Subjects may remain on treatment for up to 24 months (for those randomized to pembrolizumab arms), until disease progression or another protocol-specified reason for discontinuation from therapy. After discontinuation from treatment, the subject is followed for survival.

KEYNOTE-689 (KN689) is an open-label, Phase 3 randomized, active-controlled, multisite study that evaluated pembrolizumab as Neoadjuvant Therapy and in Combination With Standard of Care as Adjuvant Therapy for Stage III-IVA Resectable Locoregionally Advanced Head and Neck Squamous Cell Carcinoma (LA HNSCC). Participants were randomized in 1:1 ratio to one of the following arms: Arm A - Pembrolizumab (2 cycles) as neoadjuvant therapy followed by surgical resection followed by adjuvant therapy with pembrolizumab (15 cycles) plus SoC RT ± cisplatin or Arm B - No neoadjuvant therapy prior to surgical resection followed by adjuvant therapy with SoC RT ± cisplatin. KEYNOTE-012 (KN012) is a Phase Ib, nonrandomized, open-label, multicenter, multicohort trial of pembrolizumab in subjects with advanced solid tumors. Subjects were enrolled into Cohort A for triple negative breast cancer (TNBC), Cohort B for the initial head and neck (H/N) cancer cohort and Cohort B2 for the H/N cancer expansion cohort, Cohort C for urothelial tract cancer, or Cohort D for gastric cancer. Only subjects with PD-L1 positive tumors were enrolled in Cohorts A, B, C and D. Subjects in Cohort B2 were enrolled regardless of PD-L1 status. Subjects may remain on treatment for 24 months, until disease progression or another protocol-specified reason for discontinuation from therapy. After discontinuation from treatment, the subject is followed for survival.

KEYNOTE-055 (KN055) is a Phase II, nonrandomized, multicenter, single-cohort trial of pembrolizumab in subjects with recurrent and/or metastatic head and neck squamous cell carcinoma who have progressed on platinum and cetuximab therapy. Subjects may remain on treatment for 24 months, until disease progression or another protocol-specified reason for discontinuation from therapy. After discontinuation from treatment, the subject is followed for survival.

KEYNOTE-054 (KN054) is an international, double-blind, placebo-controlled randomized Phase III trial of pembrolizumab adjuvant treatment in subjects with high-risk stage III melanoma following complete surgical resection compared to placebo. Eligible subjects were randomized into two treatment groups to receive pembrolizumab 200 mg fixed dose or placebo every 3 weeks for one year or up to 18 doses. Subjects were stratified by stage (IIIA [> 1 mm metastasis] versus IIIB versus IIIC 1 to 3 positive lymph nodes versus IIIC >4 positive lymph nodes) and geographical region (North America, European countries, Australia and other countries as designated). After first disease recurrence, subjects have the option to participate in Part 2 of the study; a single treatment group to receive pembrolizumab 200 mg fixed dose every 3 weeks for up to 2 years or until occurrence of a withdrawal criterion. Subjects assigned to the placebo group in Part 1 who experience disease recurrence are offered to crossover to pembrolizumab in Part 2. Subjects randomized to pembrolizumab in Part 1 are considered for re-treatment if their disease recurs >6 months after completing 1 year of adjuvant treatment. Re-challenge is optional and at the discretion of the Investigator.

KEYNOTE-716 (KN716) is a Phase III, randomized, placebo-controlled, multicenter study of adjuvant pembrolizumab in participants 12 years of age and older with resected Stage IIB or

IIC cutaneous melanoma per AJCC eighth edition guidelines. Eligible participants were randomized into two treatment groups to receive pembrolizumab (Adult: 200 mg IV Q3W or Pediatric: 2 mg/kg IV Q3W up to a maximum of 200 mg Q3W) or placebo every 3 weeks for 17 cycles. Participants were stratified as follows: one stratum for pediatric participants (≥ 12 years of age and < 18 years of age) and 3 strata for adult participants (≥ 18 years of age) based on T-stage tumor thickness and ulceration (T3b, T4a, T4b). After first disease recurrence, participants have the option to participate in Part 2 of the study; a single treatment group to receive up to 17 or 35 cycles of pembrolizumab therapy Q3W per Part 2 eligibility guidelines.

KEYNOTE-426 (KN426) is a Phase III randomized, open-label study to evaluate efficacy and safety of pembrolizumab in combination with axitinib versus sunitinib monotherapy as a first-line treatment for locally advanced or metastatic renal cell carcinoma (mRCC). Eligible subjects were stratified by the International Metastatic Renal Cell Carcinoma Database Consortium (IMDC) risk categories (favorable versus intermediate versus poor) and geographic region (North America versus Western Europe versus “Rest of the World”). Subjects were randomized in a 1:1 ratio to one of the 2 treatment arms in the trial. Study treatments are continued until progressive disease (PD) is BICR-verified or further confirmed by the investigator, unacceptable adverse events (AEs) or intercurrent illness prevents further administration of treatment, death or withdrawal of consent. Following verification or confirmation of PD, all subjects are followed for survival (by phone contact or clinic visit) until death, withdrawal of consent, loss to follow-up, or until the study is concluded or terminated early, whichever comes first.

KEYNOTE-581 (KN581) is a Phase III randomized, open-label study to evaluate efficacy and safety of pembrolizumab plus lenvatinib and lenvatinib plus everolimus versus sunitinib monotherapy as a first-line treatment for advanced or mRCC. Eligible participants were stratified by the Memorial Sloan-Kettering Cancer Center (MSKCC) risk score (favorable, intermediate versus poor risk groups) and geographic region (North America and Western Europe versus Other). Participants were randomized in a 1:1:1 ratio to one of the 3 treatment arms in the trial. Study treatments are continued until confirmed PD by independent review, development of unacceptable toxicity, participant request, withdrawal of consent or study termination by the sponsor. Participants will then enter the follow-up period. The follow up period begins the day after the Off-Treatment Visit and continues as long as the participants is alive, unless the participant withdraws consent, is lost to follow-up, or the sponsor terminates the study.

KEYNOTE-564 (KN564) is a Phase III, randomized, double-blind, placebo-controlled, multicenter, global study to evaluate the efficacy and safety of pembrolizumab in participants with RCC completely resected by nephrectomy and/or metastasectomy with intermediate-high or high risk of recurrence. Eligible patients were randomly assigned in a 1:1 ratio to receive pembrolizumab or placebo as adjuvant therapy following nephrectomy/metastasectomy. Randomization was stratified by metastatic status by investigator review (M0 versus M1 no evidence of disease [NED]). Within the M0 group, randomization was further stratified by ECOG performance status (0 versus 1) and geographic location (United States versus outside of the United States). The study treatment was administered for up to approximately 1 year, or until disease recurrence, unacceptable toxicity, intercurrent illness preventing further administration of study treatment, investigator’s decision, protocol noncompliance, or

administrative reasons requiring discontinuation of study treatment. Participants will then enter the follow-up period. After the end of treatment, participants will be followed for the occurrence of AEs. Participants who discontinue for reasons other than disease recurrence will have posttreatment imaging follow-up for disease status until confirmation of disease recurrence, initiating a new anticancer treatment, withdrawing consent, or becoming lost to follow-up. All participants will be followed for OS until death, withdrawal of consent, or the end of the study.

KEYNOTE-177 (KN177) is a 2-arm, multicenter, international, randomized, open-label, Phase III study evaluating the efficacy and safety of pembrolizumab monotherapy versus globally accepted SOC therapies for CRC in participants with locally confirmed dMMR or MSI-H unresectable or metastatic CRC who have not received prior chemotherapy for unresectable or metastatic CRC. Participants were randomized 1:1 to receive pembrolizumab (200 mg Q3W) or investigator's choice of SOC therapies for CRC (mFOLFOX6 or FOLFIRI with or without combination with bevacizumab or cetuximab). Treatment with pembrolizumab was administered until documented disease progression, patient has received 35 trial treatments (for pembrolizumab), unacceptable toxicity, or other protocol defined stopping criteria. Those subjects that received investigator's choice chemotherapy and have centrally verified progression can crossover and receive pembrolizumab therapy for up to 35 cycles.

KEYNOTE-590 (KN590) is a randomized, double-blind, placebo-controlled, worldwide, Phase III trial to evaluate the efficacy and safety of pembrolizumab in combination with cisplatin and 5-fluorouracil (5-FU) versus placebo in combination with cisplatin and 5-FU as first-line treatment in subjects with locally advanced unresectable or metastatic adenocarcinoma or squamous cell carcinoma of the esophagus or advanced/metastatic Siewert type 1 adenocarcinoma of the esophagogastric junction (GEJ). Subjects were randomized 1:1 to receive pembrolizumab 200 mg intravenous (IV) or saline placebo once every three weeks or chemotherapy (5-fluorouracil 800 mg/m² IV on days 1-5 plus cisplatin 80 mg/m² IV [for a maximum of 6 cycles]) every 3 weeks. Randomization was stratified by geographic region (Asia vs non-Asia), histology (esophageal squamous cell carcinoma versus adenocarcinoma), and ECOG performance status (0 vs 1). Treatment was continued until confirmed disease progression, completion of 24 months of treatment, or other protocol defined stopping criteria.

KEYNOTE-355 (KN355) is a randomized, double-blind, multi-center, placebo-controlled, Phase III study evaluating the efficacy and safety of pembrolizumab in combination with chemotherapy (paclitaxel, or nab-paclitaxel, or gemcitabine and carboplatin) in subjects with locally recurrent unresectable or metastatic TNBC, regardless of tumor PD-L1 expression, and not previously treated with chemotherapy. Subjects were randomized (2:1) to one of the two treatment arms (pembrolizumab + chemotherapy or placebo + chemotherapy) stratified by chemotherapy treatment (taxane vs. gemcitabine/carboplatin), tumor PD-L1 expression (CPS ≥ 1 vs. CPS < 1), and prior treatment with the same class of chemotherapy in the neoadjuvant setting (yes vs. no). Treatment with pembrolizumab or placebo was administered until confirmed disease progression, unacceptable toxicity, or a maximum of 24 months. Administration with pembrolizumab was permitted beyond disease progression if the subject was clinically stable and considered to be deriving clinical benefit.

KEYNOTE-522 (KN522) is a randomized, double-blind, multi-center, placebo-controlled, Phase III study evaluating pembrolizumab plus chemotherapy (paclitaxel and carboplatin followed by doxorubicin, or epirubicin, and cyclophosphamide) versus placebo plus chemotherapy as neoadjuvant therapy and pembrolizumab versus placebo as adjuvant therapy in patients with previously untreated locally advanced TNBC. Subjects were randomized (2:1) to one of the two treatment arms (pembrolizumab + chemotherapy or placebo + chemotherapy), and stratified by nodal status (positive vs. negative), tumor size (T1/T2 vs. T3/T4) and choice of carboplatin (Q3W vs. weekly). Treatment was continued until completion of study treatment (17 cycles of pembrolizumab/placebo), disease progression in the neoadjuvant phase or recurrence (local or distance) after surgery, or other protocol defined stopping criteria.

KEYNOTE-146 (KN146) is a multi-center, multi-cohort, open-label, Phase Ib/II trial of the combination of lenvatinib and pembrolizumab in patients with advanced/metastatic NSCLC, RCC, endometrial carcinoma, urothelial carcinoma, HNSCC, or melanoma. In KN146, lenvatinib dose is 20 mg orally once daily (the maximum tolerated dose/recommended phase II dose) in combination with pembrolizumab 200 mg administered intravenously every 3 weeks. Patients received study treatment until unacceptable toxicity or disease progression as determined by the investigators. Administration of lenvatinib and pembrolizumab was permitted beyond RECIST-defined disease progression if the patient was clinically stable and considered to be deriving clinical benefit by the investigator. Pembrolizumab dosing was continued up to 35 treatments (approximately 2 years); however, treatment with lenvatinib could be continued beyond 2 years. The primary safety data from KN146 includes all patients with histologically confirmed endometrial carcinoma from Study KN146 who received at least 1 dose of lenvatinib 20 mg/day with pembrolizumab 200 mg Q3W as starting dose, who received 1 or more prior systemic anticancer therapy(ies), who have a median follow-up time of at least 12 months and minimum 6 months follow-up after objective response assessment per investigator (ie, enrolled prior to 01-JUL-2018), and whose tumors are not MSI-H or dMMR.

KEYNOTE-775 (KN775) is a multicenter, randomized, open-label, Phase 3 study to compare the efficacy and safety of lenvatinib in combination with pembrolizumab versus treatment of physician's choice (TPC) in participants with advanced endometrial carcinoma who have been treated with at least one prior platinum-based chemotherapy regimen. In KN775, participants were randomized to one of two treatment arms - Arm A, lenvatinib 20 mg orally once daily plus pembrolizumab 200 mg intravenously every 3 weeks. Arm B, TPC consisting of either doxorubicin 60 mg/m² by IV bolus injection, 1-hour infusion, or per institutional guidelines every 3 weeks or paclitaxel 80 mg/m² by 1-hour IV infusion or per institutional guidelines given weekly, 3 weeks on/1 week off. Crossover between treatment arms is not permitted. Prior to randomization, investigators selected and recorded the TPC option in the event the participant was assigned to that arm. Randomization followed a predefined randomization scheme based on specific stratification factors. Each eligible participant is assigned to receive study treatment until disease progression is radiographically documented and verified by blinded independent central review (BICR) per RECIST 1.1, unacceptable adverse event(s), withdrawal of consent, until the participant has received a maximum 35 administrations of pembrolizumab (approximately 2 years), or the participant has received a lifetime cumulative dose of 500 mg/m² of doxorubicin. Participants who stop study treatment after receiving 35

administrations of pembrolizumab for reasons other than disease progression or intolerability, or participants who attain a complete response (CR) and stop study treatment may be eligible for up to 1 year of treatment with pembrolizumab (17 cycles) ± lenvatinib upon experiencing disease progression (Second Course Phase). Participants who complete treatment with pembrolizumab after 35 cycles or CR will continue to receive lenvatinib alone until disease progression is confirmed by BICR, development of unacceptable toxicity, or withdrawal of consent. Participants are permitted to continue study treatment beyond RECIST 1.1-defined disease progression per investigator's assessment as long as the maximum dose of the study drugs have not been reached, if the treating investigator considers that the participant may experience clinical benefit with continued treatment, and the participant is tolerating study treatment.

KEYNOTE-868 (KN868) is a multicenter, randomized, placebo-controlled, double blind Phase 3 study to compare the efficacy and safety of pembrolizumab versus placebo, each in combination with paclitaxel and carboplatin in female participants ≥ 18 years of age with newly diagnosed, measurable Stage III or IVA EC, or Stage IVB or recurrent EC, each with or without measurable disease. In KN868, participants were randomized into two cohorts according to whether they were pMMR or dMMR. In each cohort, participants were randomized to one of two treatment arms - Arm 1, Placebo (Q3W) in combination with chemotherapy (carboplatin AUC 5 and paclitaxel 175 mg/m² Q3W) for 6 cycles (Stage 1) followed by placebo alone (Q6W) for up to 14 cycles (Stage 2) Arm 2, Pembrolizumab (200mg Q3W) in combination with chemotherapy (carboplatin AUC 5 and paclitaxel 175 mg/m² Q3W) for 6 cycles (Stage 1) followed by pembrolizumab alone (400mg Q6W) for up to 14 cycles (Stage 2). Crossover between treatment arms was not permitted. Randomization followed a predefined randomization scheme based on specific stratification factors. Each eligible participant is assigned to receive study treatment until disease progression is radiographically assessed by the investigator per RECIST 1.1, unacceptable adverse event(s), withdrawal of consent or until the participant has received a maximum 20 administrations of pembrolizumab or placebo (approximately 2 years).

KEYNOTE-158 (KN158) is a multicenter, global, open-label Phase II study of pembrolizumab monotherapy in multiple types of advanced (unresectable and/or metastatic) rare cancer. All patients enrolled in this study must have a histologically or cytologically documented, advanced solid tumor that is incurable and for which prior standard first-line treatment has failed. Patients must have progressed on or be intolerant to therapies that are known to provide clinical benefit. Treatment with pembrolizumab is administered until documented disease progression, patient has received 35 trial treatments (for pembrolizumab), unacceptable toxicity, or other protocol defined stopping criteria. This multi-histology clinical study has been designed to enroll a "basket" of rare tumor types. Patients with any of the following solid tumor types have been enrolled:

- (A) Anal squamous cell carcinoma,
- (B) Biliary adenocarcinoma (gallbladder or biliary tree (intrahepatic or extrahepatic cholangiocarcinoma) except ampulla of Vater cancers,

- (C) Neuroendocrine tumors (well- and moderately-differentiated), of the lung, appendix, small intestine, colon, rectum, or pancreas,
- (D) Endometrial carcinoma (sarcomas and mesenchymal tumors are excluded),
- (E) Cervical squamous cell carcinoma,
- (F) Vulvar squamous cell carcinoma,
- (G) Small cell lung carcinoma,
- (H) Mesothelioma (malignant pleural mesothelioma),
- (I) Thyroid carcinoma (papillary or follicular subtypes),
- (J) Salivary gland carcinoma (sarcomas and mesenchymal tumors are excluded)

OR

- (K) Any advanced solid tumor (except CRC) that is MSI-H.
- (L) Any advanced solid tumor (including CRC) which is dMMR/MSI-H in subjects from mainland China who are of Chinese descent,
- (M) Any advanced solid tumor that has failed at least one systemic line of therapy and is Tumor Mutational Burden-High (TMB-H) (≥ 10 mut/Mb, F1CDx assay), excluding dMMR/MSI-H tumors.

KEYNOTE-164 (KN164) is a single-arm, multisite, two cohort, Phase II study of pembrolizumab that was designed to evaluate the efficacy of pembrolizumab in patients with locally advanced unresectable or metastatic (Stage IV) MSI-H/dMMR CRC. Treatment with pembrolizumab is administered until documented disease progression, patient has received 35 trial treatments (for pembrolizumab), unacceptable toxicity, or other protocol defined stopping criteria. Patients who met eligibility criteria have been enrolled in the following 2 cohorts:

- Cohort A: Participants with locally advanced unresectable or metastatic dMMR or MSI-H CRC who had been previously treated with at least 2 lines of standard of care therapies, which must have included fluoropyrimidine, oxaliplatin, and irinotecan.
- Cohort B: Participants with locally advanced unresectable or metastatic dMMR or MSI-H CRC who had been previously treated with at least 1 line of systemic standard of care therapy (fluoropyrimidine + oxaliplatin or fluoropyrimidine + irinotecan \pm antivascular endothelial growth factor [anti-VEGF]/ epidermal growth factor receptor [EGFR] monoclonal antibody [mAb]).

KEYNOTE-826 (KN826) is a Phase III, randomized, double-blind, placebo-controlled trial of pembrolizumab plus chemotherapy versus chemotherapy alone in women with persistent, recurrent, or metastatic cervical cancer who are not eligible for treatment with curative intent

(such as with surgery and/or radiation) and who have not previously been treated with systemic chemotherapy, with the exception of chemotherapy used as a radio sensitizing agent. Patients were randomized 1:1 to receive pembrolizumab with carboplatin/cisplatin + paclitaxel or placebo with carboplatin/cisplatin + paclitaxel. Bevacizumab was also permitted by investigator choice. Participants received study treatments until disease progression, unacceptable AEs, or other protocol defined stopping criteria.

KEYNOTE-A18 (KNA18) is an ongoing, randomized, Phase III placebo-controlled, multisite, double-blind study of chemoradiotherapy with or without pembrolizumab in participants with high-risk LACC (FIGO 2014 Stage IB2-IIB [with node-positive disease] or Stage III-IVA [either node-positive or node-negative disease]). Patients were randomized in a 1:1 ratio to receive concurrent chemoradiotherapy in combination with pembrolizumab (200 mg Q3W for 5 infusions) followed by pembrolizumab monotherapy (400 mg Q6W for 15 infusions) or concurrent chemoradiotherapy in combination with placebo (Q3W for 5 infusions) followed by placebo alone (Q6W for 15 infusions). Standard of care chemoradiotherapy was given during the Q3W dosing period of pembrolizumab (Arm 1) or placebo (Arm 2) and included: Cisplatin, 5 infusions given IV QW at 40 mg/m² (an optional, sixth infusion of cisplatin could be administered according to local practice) plus EBRT over 40 days, followed by brachytherapy. Participants received study treatments until radiographic disease progression, unacceptable AE, or other protocol defined stopping criteria.

KEYNOTE-811 (KN811) is a global, randomized, double-blind, Phase III study of pembrolizumab or placebo in combination with trastuzumab and the investigator's choice of chemotherapy with 5-fluorouracil and cisplatin or capecitabine and oxaliplatin in participants with previously untreated unresectable or metastatic, HER2-positive gastric or gastroesophageal junction adenocarcinoma. Patients may remain on treatment for 24 months, until disease progression or another protocol-specified reason for discontinuation from therapy. Patients who complete 35 administrations of pembrolizumab/placebo have the option to receive up to 1 additional year of trastuzumab +capecitabine or 5-fluorouracil (5-FU) or S-1 (tegafur, gimeracil, oteracil) at the discretion of the investigator. After discontinuation from treatment, patients are followed for survival.

KEYNOTE-859 (KN859) is a global, randomized, double-blind, Phase III study of pembrolizumab or placebo in combination with the investigator's choice of chemotherapy with 5-fluorouracil and cisplatin or capecitabine and oxaliplatin in participants with previously untreated unresectable or metastatic, HER2-negative gastric or gastro-oesophageal junction adenocarcinoma. Patients were stratified by geographical region (Europe/Israel/North America/Australia, Asia or Rest of the World), PD-L1 tumor expression status (CPS<1, ≥1) and chemotherapy regimen (FP or CAPOX). Patients may remain on treatment for 24 months, until disease progression or another protocol-specified reason for discontinuation from therapy.

KEYNOTE-966 (KN966) is a global, randomized, double-blinded, multi-site Phase 3 trial of pembrolizumab plus gemcitabine/cisplatin versus placebo plus gemcitabine/cisplatin in patients with locally advanced unresectable or metastatic BTC who have not previously been treated with systemic chemotherapy, with the exception of adjuvant or neoadjuvant therapy. Prior therapy with an anti-PD-1, anti-PD-L1, or anti-PD-L2 agent or with an agent directed to

another stimulatory or coinhibitory T-cell receptor (eg, CTLA-4, OX-40, CD137) was exclusionary. The trial includes patients with BTC diagnosis confirmed by histology, or cytology (small cell cancer, neuroendocrine tumors, lymphoma, sarcoma, and/or mucinous cystic neoplasms are not eligible). To be eligible, patients must have disease not amenable to a curative treatment approach. Patients must also have measurable disease based on RECIST 1.1, as determined by the site investigator, an Eastern Cooperative Oncology Group (ECOG) performance score of 0 or 1 and predicted life expectancy of greater than 3 months. Patients are enrolled regardless of tissue PD-L1 biomarker status. Patients are required to provide a tumor tissue sample for biomarker analysis. Per protocol defined criteria patients with controlled HBV infection on antiviral therapy, as well as patients with past or ongoing HCV infection were eligible for the trial. Participants received study treatments until disease progression, unacceptable AEs, or other protocol defined stopping criteria.

KEYNOTE-483 (KN483) is an open-label, multi-center, Phase 2/3 randomized study that evaluated pembrolizumab in combination with platinum and pemetrexed for the treatment of patients with previously untreated unresectable advanced and/or metastatic MPM. In the phase 2 part, participants were randomized in 1:1:1 ratio to platinum and pemetrexed, platinum and pemetrexed in combination with pembrolizumab, or pembrolizumab monotherapy. In the phase 3 part, participants were randomized in 1:1 ratio to platinum and pemetrexed or platinum and pemetrexed in combination with pembrolizumab. Participants received study treatments until disease progression, unacceptable AEs, or other protocol defined stopping criteria. To be eligible, patients must be ≥ 18 years of age, must have histologically confirmed unresectable advanced and/or metastatic MPM incurable by standard therapy, must be eligible to receive standard chemotherapy, must have available tissue sample, must have measurable disease by mRECIST or RECIST 1.1, and must have ECOG PS of 0 or 1. Participants were not eligible if they received prior chemotherapy for any stage of advanced/metastatic disease; received targeted small molecule therapy, immunotherapies and viral therapies, biologic therapies and angiogenesis inhibitors for advanced/metastatic disease; received any prior immunotherapy for any stage of disease; had active autoimmune disease that required systemic treatment within the previous 3 years; had known active CNS metastases and/or carcinomatous meningitis; had untreated and/or uncontrolled cardiovascular conditions and/or symptomatic cardiac dysfunction, or had unstable angina, congestive heart failure, or myocardial infarction within the previous year. Patients who received previous (neo)adjuvant cisplatin-based systemic chemotherapy must have received the last dose of chemotherapy at least 12 months before registration.

KEYNOTE-A17 (KNA17) was a multicenter, open-label, non-randomized, phase 1b study to evaluate the safety, tolerability and preliminary efficacy of pembrolizumab in combination with cisplatin and pemetrexed in treatment-naïve, Japanese participants with a histologically confirmed diagnosis of advanced/unresectable MPM. To be eligible, patients must be ≥ 20 years of age, must have confirmed diagnosis of advanced/unresectable MPM, must have systemic therapy naïve disease measurable per mRECIST by investigator, and must have ECOG PS of 0 or 1. Participants were not eligible if they received prior systemic anti-cancer therapy for advanced/metastatic MPM; received prior therapy with an anti-PD-1, anti-PD-L1, or anti PD-L2 agent or with an agent directed to another stimulatory or co-inhibitory T-cell receptor; had active autoimmune disease that required systemic treatment within the previous 2 years; had known active CNS metastases and/or carcinomatous meningitis; Patients who

received previous (neo)adjuvant systemic chemotherapy must have received the last dose of chemotherapy at least 6 months before registration.

Clinical Trial Summary of Drug Exposure

Table SIII.1 displays duration of exposure for all patients with advanced melanoma who received pembrolizumab monotherapy in the pooled KN001, KN002 and KN006 all patients as treated (APaT) population.

Table SIII.2 displays duration of exposure for all patients with melanoma who received pembrolizumab monotherapy in KN054 and KN716 APaT population. Of note, the median exposure to pembrolizumab in KN054 and KN716 is more than twice as long as the reference safety dataset (8.78 months vs 4.18 months, respectively).

Table SIII.3 displays duration of exposure for all patients with NSCLC who received pembrolizumab monotherapy in the pooled KN001, KN010, KN024, and KN042 APaT population.

Table SIII.4 displays duration of exposure to pembrolizumab for all patients with NSCLC who received pembrolizumab combination therapy in the pooled KN021 (Cohort A, C and G), KN189 and KN407 APaT population.

Table SIII.5 displays duration of exposure for all patients with NSCLC who received pembrolizumab monotherapy in KN091 APaT population. Of note, the median exposure to pembrolizumab in KN091 is more than twice as long as the reference safety dataset (11.7 months vs 4.18 months, respectively).

Table SIII.6 displays duration of exposure to pembrolizumab for all patients with NSCLC who received pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy in KN671 APaT population.

Table SIII.7 displays duration of exposure for all patients with NSCLC who received pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy in MK-3475A-D77 APaT population.

Table SIII.8 displays duration of exposure for all patients with HL who received pembrolizumab monotherapy in the pooled KN013 (Cohort 3), KN087, and KN204 APaT population.

Table SIII.9 displays duration of exposure for all patients with UC who received pembrolizumab monotherapy in the pooled KN045 and KN052 APaT population.

Table SIII.10 displays duration of exposure for all patients with UC who received pembrolizumab combination therapy in the pooled KNA39 and KN869 APaT population.

Table SIII.11 displays duration of exposure for all patients with HNSCC who received pembrolizumab monotherapy in the pooled KN012 (B+B2 Cohorts), KN040, KN055, and KN048 APaT population.

Table SIII.12 displays duration of exposure to pembrolizumab for all patients with HNSCC who received pembrolizumab combination therapy in KN048 ASaT Population.

Table SIII.13 displays duration of exposure to pembrolizumab for all patients with HNSCC who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy in KN689 APaT Population.

Table SIII.14 displays duration of exposure to pembrolizumab for all patients with RCC who received pembrolizumab combination therapy in the KN426 ASaT population.

Table SIII.15 displays duration of exposure to pembrolizumab for all patients with RCC who received pembrolizumab combination therapy in the KN581 APaT population.

Table SIII.16 displays duration of exposure for all patients with RCC who received pembrolizumab monotherapy in the KN564 APaT population.

Table SIII.17 displays duration of exposure for the patients with CRC who received pembrolizumab monotherapy in the KN177 ASaT population.

Table SIII.18 displays duration of exposure to pembrolizumab for all patients with esophageal cancer who received pembrolizumab combination therapy in KN590 ASaT population.

Table SIII.19 displays duration of exposure to pembrolizumab for all patients with TNBC who received pembrolizumab combination therapy in KN355 ASaT population.

Table SIII.20 displays duration of exposure to pembrolizumab for all patients with TNBC who received pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy in the KN522 APaT population.

Table SIII.21 displays duration of exposure to pembrolizumab for all patients with endometrial carcinoma who received pembrolizumab combination therapy in the KN146 and KN775 APaT population.

Table SIII.22 displays duration of exposure to pembrolizumab for all patients with endometrial carcinoma who received pembrolizumab combination therapy in the KN868 APaT population.

Table SIII.23 displays duration of exposure for all patients with MSI-H who received pembrolizumab monotherapy in the KN158-Cohort K and KN164-Cohorts A and B ASaT population.

Table SIII.24 displays duration of exposure to pembrolizumab for all patients with cervical cancer who received pembrolizumab combination therapy in the KNA18 and KN826 APaT population.

Table SIII.25 displays duration of exposure to pembrolizumab for all patients with gastric cancer who received pembrolizumab combination therapy in the KN811 APaT population. Of note, the median exposure to pembrolizumab in KN811 is more than twice as long as the

pembrolizumab monotherapy reference safety dataset (9.10 months vs 4.18 months, respectively).

Table SIII.26 displays duration of exposure to pembrolizumab for all patients with gastric cancer who received pembrolizumab combination therapy in the KN859 APaT population.

Table SIII.27 displays duration of exposure to pembrolizumab for all patients with BTC who received pembrolizumab combination therapy in the KN966 APaT population.

Table SIII.28 displays duration of exposure to pembrolizumab for all patients with mesothelioma who received pembrolizumab combination therapy in the KN483 and KNA17 APaT population.

Table SIII.1: Summary of Drug Exposure Melanoma Subjects KN001, KN002 and KN006 Monotherapy (APaT Population)

	Pembrolizumab 2 mg/kg Q3W	Pembrolizumab 10 mg/kg Q3W	Pembrolizumab 10 mg/kg Q2W	Total
	N=340	N=769	N=458	N=1567
Study Days On-Therapy (days)				
Mean	230.04	224.66	231.36	227.79
Median	129.00	148.00	183.00	155.00
SD	204.06	189.74	185.88	191.74
Range	1.00 to 729.00	1.00 to 759.00	1.00 to 862.00	1.00 to 862.00
Number of Administrations				
Mean	11.43	11.26	16.61	12.86
Median	7.00	8.00	13.00	9.00
SD	9.31	8.69	12.46	10.34
Range	1.00 to 34.00	1.00 to 36.00	1.00 to 59.00	1.00 to 59.00
(pembrolizumab KN001 Database Cutoff Date: 18APR2014). (pembrolizumab KN002 Database Cutoff Date: 28FEB2015). (pembrolizumab KN006 Database Cutoff Date: 03MAR2015).				

Data on File

Table SIII.2: Summary of Drug Exposure Melanoma Participants KN054 and KN716 Monotherapy (APaT Population)

	KN716 + KN054 Pembrolizumab Q3W (N=992)
Study Days On-Therapy (Months)	
Mean	8.78
Median	11.07
SD	3.89
Range	0.03 to 15.70
Number of Administrations	
Mean	13.20
Median	17.00
SD	5.47
Range	1.00 to 18.00
Each participant is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. Database cutoff date for Melanoma (KN054: 02OCT2017, KN716: 04DEC2020)	

Data on File

Table SIII.3: Summary of Drug Exposure NSCLC Subjects KN001, KN010, KN024, and KN042 Monotherapy (APaT Population)

	Pembrolizumab 2 mg/kg Q3W (N=400)	Pembrolizumab 10 mg/kg Q3W (N=630)	Pembrolizumab 10 mg/kg Q2W (N=202)	Pembrolizumab 200 mg Q3W (N=790)	Total (N=2022)
Study Days On-Therapy (days)					
Mean	145.01	164.77	176.32	269.07	202.77
Median	106.00	106.00	113.00	176.00	127.00
SD	137.66	165.27	166.20	249.20	205.26
Range	1.00 to 681.00	1.00 to 925.00	1.00 to 601.00	1.00 to 988.00	1.00 to 988.00
Number of Administrations					
Mean	7.56	8.41	12.91	13.23	10.57
Median	6.00	6.00	9.00	9.00	7.00
SD	6.30	7.52	11.38	11.46	9.80
Range	1.00 to 30.00	1.00 to 45.00	1.00 to 42.00	1.00 to 48.00	1.00 to 48.00
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. (pembrolizumab KN001 Database Cutoff Date for Lung: 23JAN2015). (pembrolizumab KN010 Database Cutoff Date: 30SEP2015). (pembrolizumab KN024 Database Cutoff Date: 10JUL2017). (pembrolizumab KN042 Database Cutoff Date: 26FEB2018).					

Data on File

Table SIII.4: Summary of Drug Exposure to Pembrolizumab NSCLC Subjects KN021 (Cohorts A, C and G), KN189 and KN407 Combination Therapy (APaT Population)

	Pembrolizumab 10mg/kg Combo (N=24)	Pembrolizumab 2mg/kg Combo (N=25)	Pembrolizumab 200mg Combo (N=742)	Total (N=791)
Study Days On-Therapy (days)				
Mean	245.29	386.64	222.62	228.49
Median	172.50	302.00	198.00	199.00
SD	226.91	267.05	151.76	161.55
Range	1.00 to 862.00	29.00 to 785.00	1.00 to 750.00	1.00 to 862.00
Number of Administrations				
Mean	11.21	17.52	10.72	10.95
Median	9.00	15.00	10.00	10.00
SD	9.50	12.18	6.87	7.27
Range	1.00 to 35.00	2.00 to 35.00	1.00 to 35.00	1.00 to 35.00
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. MK-3475 Database Cutoff Date for Lung Combination Therapy (KN021 CohortA: 07NOV2016, Cohort G/C: 31MAY2017, KN189: 8NOV2017, KN047: 03APR2018)				

Data on File

Table SIII.5: Summary of Drug Exposure NSCLC Participants KN091 Monotherapy (APaT Population)

	Pembrolizumab 200 mg Q3W (N=580)
Study Days On-Therapy (Months)	
Mean	8.66
Median	11.70
SD	4.50
Range	0.03 to 18.86
Number of Administrations	
Mean	12.85
Median	17.00
SD	6.22
Range	1.00 to 18.00
Each participant is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. KN091 Database Cutoff Date: 20SEP2021	

Data on File

Table SIII.6: Summary of Drug Exposure NSCLC Participants in KN671 Combination Therapy (APaT Population)

	Pembrolizumab 200 mg Q3W (N=396)
Study Days On-Therapy (Months)	
Mean	9.32
Median	10.91
SD	5.35
Range	0.03 to 18.63
Number of Administrations	
Mean	10.92
Median	12.00
SD	5.88
Range	1.00 to 17.00
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. Database Cutoff Date: 29JUL2022	

Data on File

Table SIII.7: Summary of Drug Exposure NSCLC Participants in MK-3475A-D77 Combination Therapy (APaT Population)

	Pembrolizumab SC 790 mg Q6W (N=251)
Duration of Pembrolizumab Exposure (months)	
Mean	6.4
Median	6.9
SD	3.3
Range	0.03 to 13.17
Number of Pembrolizumab Administrations	
Mean	5.2
Median	5.0
SD	2.4
Range	1.00 to 10.00
Each participant is counted once on each applicable duration category row. Duration of exposure (months) is calculated as (last dose date - first dose date + 1) / 30.4367 for pembrolizumab. Database Cutoff Date for Lung (MK-3475A-D77: 12JUL2024).	

Data on File

Table SIII.8: Summary of Drug Exposure HL Subjects KN013, KN087, and KN204 Monotherapy (APaT Population)

	Pembrolizumab 10 mg/kg Q2W (N=31)	Pembrolizumab 200 mg Q3W (N=358)	Total (N=389)
Study Days On-Therapy (months)			
Mean	10.74	12.67	12.52
Median	7.85	11.07	10.65
SD	8.97	8.18	8.25
Range	0.03 to 27.93	0.03 to 27.60	0.03 to 27.93
Number of Administrations			
Mean	22.84	18.67	19.01
Median	18.00	16.00	16.00
SD	17.73	11.53	12.16
Range	1.00 to 52.00	1.00 to 36.00	1.00 to 52.00
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. Database Cutoff Date (KN013-Cohort 3: 28SEP2018, KN087: 21MAR2019, KN204: 16JAN2020)			

Data on File

Table SIII.9: Summary of Drug Exposure UC Subjects KN045 and KN052 Monotherapy (APaT Population)

	Pembrolizumab 200 mg Q3W N=636	Total N=636
Time on Therapy (days)		
Mean	135.64	135.64
Median	85.00	85.00
SD	131.72	131.72
Range	1.00 to 610.00	1.00 to 610.00
Number of Administrations		
Mean	7.16	7.16
Median	5.00	5.00
SD	6.09	6.09
Range	1.00 to 30.00	1.00 to 30.00
Includes all subjects with urothelial carcinoma who received at least one dose of Pembrolizumab in KN045 and KN052. Pembrolizumab KN045 Database Cutoff Date: 07SEP2016. Pembrolizumab KN052 Database Cutoff Date: 01SEP2016.		

Data on File

Table SIII.10: Summary of Drug Exposure to Pembrolizumab Urothelial Cancer Participants in KNA39 and KN869 Pembrolizumab + EV Combination Therapy (APaT Population)

	Pembrolizumab (200 mg Q3W) + Enfortumab Vedotin (N=564)
Duration of Pembrolizumab Exposure (months)	
Mean	9.01
Median	7.61
SD	7.07
Range	0.03 to 31.67
Number of Pembrolizumab Administrations	
Mean	12.6
Median	10.0
SD	9.4
Range	1.00 to 35.00
<p>Each participant is counted once on each applicable duration category row. Duration of exposure is calculated as last dose date - first dose date + 1 for pembrolizumab in the combination. Pembrolizumab KN-869 Database Cutoff Date: 13Mar2023, KN-A39 Database Cutoff Date: 08Aug2023</p>	

Table SIII.11: Summary of Drug Exposure HNSCC Subjects KN012 B+B2 Cohorts, KN040, KN055 and KN048 Monotherapy (ASaT Population)

	Head and Neck Cancer (MK3475 10mg/kg Q2W) (N=60)	Head and Neck Cancer (MK3475 200mg Q3W) (N=849)	Total KN012 B+B2/KN040/KN055/KN048 (N=909)
Study Days On-Therapy (days)			
Mean	208.22	160.18	163.35
Median	100.50	92.00	92.00
SD	236.24	175.01	179.92
Range	1.00 to 778.00	1.00 to 816.00	1.00 to 816.00
Number of Administrations			
Mean	14.87	8.20	8.64
Median	8.00	5.00	5.00
SD	15.63	7.97	8.83
Range	1.00 to 52.00	1.00 to 40.00	1.00 to 52.00
<p>Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. MK-3475 Database Cutoff Date for HNSCC (KN012-HNSCC: 26APR2016, KN040: 15MAY2017, KN048: 13JUN2018, KN055: 22APR2016)</p>			

Data on File

Table SIII.12: Summary of Drug Exposure to Pembrolizumab HNSCC Subjects KN048 Combination Therapy (ASaT Population)

	Head and Neck Cancer (MK3475 200mg Q3W) (N=276)
Study Days On-Therapy Pembro (days)	
Mean	223.16
Median	176.00
SD	200.98
Range	1.00 to 736.00
Number of Pembro Administrations	
Mean	10.60
Median	8.00
SD	9.01
Range	1.00 to 35.00
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. MK-3475 Database Cutoff Date for KN048: 13JUN2018.	

Data on File

Table SIII.13: Summary of Drug Exposure to Pembrolizumab HNSCC Participants in KN689 Monotherapy or Combination Therapy (APaT Population)

	Pembrolizumab 200mg Q3W (N=361)
Duration of Pembrolizumab Exposure (month)	
Mean	7.59
Median	9.10
SD	5.48
Range	0.03 to 22.34
Number of Pembrolizumab Administrations	
Mean	10.02
Median	11.00
SD	6.78
Range	1.00 to 18.00
Each participant is counted once on each applicable duration category row. Duration of exposure is (months) calculated as (last dose date - first dose date + 1) /30.4367 for pembrolizumab Database cutoff date for KN689: 25JUL2024	

Data on File

Table SIII.14: Summary of Drug Exposure to Pembrolizumab RCC Subjects KN426 Combination Therapy (ASaT Population)

	KN426 Data for Pembrolizumab + Axitinib (N=429)
Study Days On-Therapy Pembrolizumab (days)	
Mean	287.04
Median	279.00
SD	175.03
Range	1.00 to 646.00
Number of Pembrolizumab Administrations	
Mean	13.78
Median	14.00
SD	8.04
Range	1.00 to 31.00
Each subject is counted once on each applicable duration category row. Study Days On-Therapy is calculated as last dose date - first dose date + 1 for pembrolizumab in the combo. MK-3475 Database Cutoff Date for KN426: 24AUG2018.	

Data on File

Table SIII.15: Summary of Drug Exposure to Pembrolizumab RCC Participants KN581 Combination Therapy (APaT Population)

	Pembrolizumab + Lenvatinib (N=352)
Study Days On-Therapy (days)	
Mean	439.8
Median	459.0
SD	260.60
Range	1 to 901
Number of Pembrolizumab Administrations	
Mean	20.7
Median	22.0
SD	11.86
Range	1 to 39
Study Days On-Therapy is calculated as last dose date - first dose date + 1 for pembrolizumab in the combo. MK-3475 Database Cutoff Date for KN581: 28AUG2020.	

Data on File

Table SIII.16: Summary of Drug Exposure RCC Participants KN564 Monotherapy (APaT Population)

	Pembrolizumab 200 mg Q3W (N=488)
Duration on therapy (months)	
Mean	9.0
Median	11.1
SD	3.70
Range	0.0 to 14.3
Number of Administrations	
Mean	13.5
Median	17.0
SD	5.19
Range	1.0 to 17.0
Each participant is counted once on each applicable duration category row. Duration of Exposure (months) is calculated as (last dose date - first dose date + 1)/30.4367. Includes all participants who received at least one dose of pembrolizumab in KN564. Database cutoff date for RCC (KN564: 14DEC2020)	

Data on File

Table SIII.17: Summary of Drug Exposure CRC Subjects KN177 Monotherapy (ASaT Population)

	KN177 Data for Pembrolizumab ^{††} (N=153)
Study Days On-Therapy (Months)	
Mean	13.3
Median	11.1
SD	10.2
Range	0.0 to 30.6
Number of Administrations	
Mean	19.0
Median	16.0
SD	14.1
Range	1.0 to 35.0
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. ^{††} Includes all subjects who received at least one dose of pembrolizumab in KN177. Database cutoff date for CRC (KN177: 19FEB2020)	

Data on File

Table SIII.18: Summary of Drug Exposure to Pembrolizumab Esophageal Cancer Subjects KN590 Combination Therapy (ASaT Population)

	KN590 Data for Pembrolizumab (N=370)
Duration On Therapy (month)	
Mean	7.5
Median	5.52
SD	6.86
Range	0.03 to 25.89
Number of Administrations	
Mean	10.8
Median	8.00
SD	9.34
Range	1.00 to 35.00
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as (last dose date - first dose date +1)/30.4367 (months). Database Cutoff date for Esophageal (KN590: 02JUL2020)	

Data on File

Table SIII.19: Summary of Drug Exposure to Pembrolizumab TNBC Subjects KN355 Combination Therapy (ASaT Population)

	Pembrolizumab (200 mg Q3W) + Chemotherapy (N=596)
Duration of Therapy Pembrolizumab (months)	
Mean	8.2
Median	5.7
SD	7.1
Range	0.0 to 33.0
Number of Pembrolizumab Administrations	
Mean	11.9
Median	9.0
SD	9.5
Range	1.0 to 35.0
Each subject is counted once on each applicable duration category row. Duration on-therapy is calculated as (last dose date - first dose date +1)/30.4367 (months) for pembrolizumab in the combo. Database Cutoff Date for KN355: 11DEC2019.	

Data on File

Table SIII.20: Summary of Drug Exposure to Pembrolizumab TNBC Subjects KN522 Combination Therapy (APaT Population)

	Pembrolizumab 200 mg Q3W (N=783)
Duration of Pembrolizumab Exposure (month)	
Mean	10.9
Median	13.31
SD	5.27
Range	0.03 to 21.91
Number of Pembrolizumab Administrations	
Mean	13.1
Median	17.00
SD	5.41
Range	1.00 to 17.00
Each participant is counted once on each applicable duration category row. Duration of exposure is the time from the first dose date to the last dose date for pembrolizumab. Database cutoff date for TNBC: (KN522: 23MAR2021)	

Data on File

Table SIII.21: Summary of Drug Exposure to Pembrolizumab Endometrial Carcinoma Participants KN146 and KN775 Combination Therapy (APaT Population)

	Pembrolizumab 200mg Q3W (N=530)
Duration of Exposure (day)	
Mean	261.7
Median	211.0
SD	209.40
Range	1.0 to 885.0
Number of Administrations	
Mean	12.6
Median	10.0
SD	9.49
Range	1.0 to 36.0
Duration of exposure (day) is calculated as last dose date - first dose date + 1. Database cutoff date for endometrial carcinoma (KN146: 18AUG2020, KN775: 26OCT2020)	

Data on File

Table SIII.22: Summary of Drug Exposure to Pembrolizumab Endometrial Carcinoma Participants Subjects KN868 Combination Therapy (APaT Population)

	Pembrolizumab* (N=382)
Duration of Pembrolizumab Exposure (Days)	
Mean	212.3
Median	170.0
SD	167.75
Range	1.0 to 729.0
Number of Pembrolizumab Administrations	
Mean	8.5
Median	8.0
SD	5.20
Range	1.0 to 29.0
Each participant is counted once on each applicable duration category row Duration of exposure is (days) calculated as last dose date - first dose date + 1 for pembrolizumab. *Number of administrations (Pembrolizumab): 200mg Q3W in Stage 1 and 400mg Q6W in Stage 2. Database cutoff date (KN868: 16DEC2022 for dMMR participants and 06DEC2022 for pMMR participants)	

Data on file

Table SIII.23: Summary of Drug Exposure MSI-H Subjects KN158 Cohort K and KN164 Cohorts A and B Monotherapy (ASaT Population)

	Pembrolizumab 200 mg Q3W (N=475)
Study Days On-Therapy (Months)	
Mean	9.7
Median	5.49
SD	9.30
Range	0.03 to 38.01
Number of Administrations	
Mean	14.3
Median	8.00
SD	12.83
Range	1.00 to 35.00
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. Database cutoff date for MSI-H (KN158-cohort K: 05OCT2020) Database cutoff date for Colorectal (KN164-cohorts A and B: 09SEP2019)	

Data on File

Table SIII.24: Summary of Drug Exposure to Pembrolizumab Cervical Cancer Participants in KNA18 + KN826 Combination Therapy (APaT Population)

	Pembrolizumab* (N=835)
Duration of Exposure (month)	
Mean	11.82
Median	11.11
SD	7.71
Range	0.03 to 27.01
Number of Pembrolizumab Administrations	
Mean	13.21
Median	12.00
SD	8.25
Range	1.00 to 35.00
Each participant is counted once on each applicable duration category row. *In KNA18, participants received 5 infusions of pembrolizumab 200 mg or placebo at Q3W before proceeding to pembrolizumab 400 mg or placebo Q6W dosing. In KN826, participants received pembrolizumab 200 mg or placebo at Q3W. Duration of exposure is calculated as last dose date - first dose date + 1 for pembrolizumab in the combo. Database cutoff date for Cervical Cancer (KNA18: 09JAN2023, KN826: 03MAY2021).	

Data on File

Table SIII.25: Summary of Drug Exposure Gastric Cancer Participants in KN811 Combination Therapy (APaT Population)

	Pembrolizumab 200 mg Q3W (N=350)
Study Days On-Therapy (Months)	
Mean	10.80
Median	9.10
SD	7.88
Range	0.03 to 30.29
Number of Administrations	
Mean	15.08
Median	13.00
SD	10.50
Range	1.00 to 35.00
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. Database Cutoff Date: 25MAY2022	

Data on file

Table SIII.26: Summary of Drug Exposure to Pembrolizumab Gastric Cancer Participants in KN859 Combination Therapy (APaT Population)

	Pembrolizumab 200 mg Q3W (N=785)
Duration of Exposure (days)	
Mean	258.5
Median	190.0
SD	227.9
Range	1 to 1025
Number of Pembrolizumab Administrations	
Mean	12.2
Median	9.0
SD	10.1
Range	1 to 35
Each participant is counted once on each applicable duration category row. Duration of exposure is calculated as last dose date - first dose date + 1. Database cutoff date for Gastric (KN859: 03OCT2022)	

Data on File

Table SIII.27: Summary of Drug Exposure to Pembrolizumab BTC Participants in KN966 Combination Therapy (APaT Population)

	KN966 Pembrolizumab (200mg Q3W) + Chemotherapy (N=529)
Duration of Exposure (Days)	
Mean	233.9
Median	184.0
SD	202.86
Range	1.0 to 853.0
Number of Administrations	
Mean	11.1
Median	9.0
SD	8.94
Range	1.0 to 35.0
Each participant is counted once on each applicable duration category row Duration of exposure is calculated as last dose date - first dose date + 1 for pembrolizumab in the combo. (pembrolizumab KN966 Database Cutoff Date: 15DEC2022)	

Data on File

Table SIII.28: Summary of Drug Exposure to Pembrolizumab Mesothelioma Participants in KN483 and KNA17 Combination Therapy (APaT Population)

	Pembrolizumab 200 mg Q3W (N=260)
Study Days On-Therapy (Months)	
Mean	9.2
Median	7.0
SD	7.1
Range	0.03 to 25.79
Number of Administrations	
Mean	13.3
Median	10.0
SD	9.8
Range	1.00 to 36.00
Each participant is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. Database Cutoff Date for Thoracic Malignancy (KN483: 16SEP2022, KNA17: 21SEP2022).	

Data on File

Clinical Trial Exposure by Dose and Duration

Table SIII.29 displays the duration of exposure to pembrolizumab monotherapy in the pooled KN001, KN002 and KN006 melanoma population with respect to time and dose.

Table SIII.30 displays the duration of exposure to pembrolizumab monotherapy in the KN054 and KN716 melanoma population with respect to time and dose.

Table SIII.31 displays the duration of exposure to pembrolizumab monotherapy in the pooled KN001, KN010, KN024, and KN042 NSCLC population with respect to time and dose.

Table SIII.32 displays the duration of exposure to pembrolizumab in the pooled KN021 (Cohorts A, C and G), KN189, and KN407 NSCLC population who received pembrolizumab combination therapy with respect to time and dose.

Table SIII.33 displays the duration of exposure to pembrolizumab monotherapy in KN091 NSCLC population with respect to time and dose.

Table SIII.34 displays the duration of exposure to pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy in the KN671 NSCLC population with respect to time and dose.

Table SIII.35 displays duration of exposure to pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy in MK-3475A-D77 NSCLC population with respect to time and dose.

Table SIII.36 displays the duration of exposure to pembrolizumab monotherapy in the pooled KN013, KN087, and KN204 HL population with respect to time and dose.

Table SIII.37 displays the duration of exposure to pembrolizumab monotherapy in the pooled KN045 and KN052 UC population with respect to time and dose.

Table SIII.38 displays the duration of exposure to pembrolizumab in the pooled KNA39 and KN869 UC population who received pembrolizumab combination therapy with respect to time and dose.

Table SIII.39 displays the duration of exposure to pembrolizumab monotherapy in the pooled KN040, KN012, KN055 and KN048 HNSCC population with respect to time and dose.

Table SIII.40 displays the duration of exposure to pembrolizumab in the KN048 HNSCC population who received pembrolizumab combination therapy with respect to time and dose.

Table SIII.41 displays the duration of exposure to pembrolizumab in the KN689 HNSCC population who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy with respect to time and dose.

Table SIII.42 displays the duration of exposure to pembrolizumab in the KN426 RCC population who received pembrolizumab combination therapy with respect to time and dose.

Table SIII.43 displays the duration of exposure to pembrolizumab in the KN581 RCC population who received pembrolizumab combination therapy with respect to time and dose.

Table SIII.44 displays the duration of exposure to pembrolizumab monotherapy in the KN564 RCC population with respect to time and dose.

Table SIII.45 displays the duration of exposure to pembrolizumab monotherapy in the KN177 CRC population with respect to time and dose.

Table SIII.46 displays the duration of exposure to pembrolizumab in the KN590 esophageal cancer population who received pembrolizumab combination therapy with respect to time and dose.

Table SIII.47 displays the duration of exposure to pembrolizumab in the KN355 TNBC population who received pembrolizumab combination therapy with respect to time and dose.

Table SIII.48 displays the duration of exposure to pembrolizumab in KN522 TNBC population who received pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy with respect to time and dose.

Table SIII.49 displays the duration of exposure to pembrolizumab in the KN146 and KN775 endometrial carcinoma population who received pembrolizumab combination therapy with respect to time and dose.

Table SIII.50 displays the duration of exposure to pembrolizumab in KN868 endometrial carcinoma population who received pembrolizumab combination therapy with respect to time and dose.

Table SIII.51 displays the duration of exposure to pembrolizumab monotherapy in the KN158 Cohort K and KN164 Cohorts A and B MSI-H population with respect to time and dose.

Table SIII.52 displays the duration of exposure to pembrolizumab in the KNA18 and KN826 cervical cancer population who received pembrolizumab combination therapy with respect to time and dose.

Table SIII.53 displays the duration of exposure to pembrolizumab in the KN811 gastric cancer population who received pembrolizumab combination therapy with respect to time and dose.

Table SIII.54 displays the duration of exposure to pembrolizumab in the KN859 gastric cancer population who received pembrolizumab combination therapy with respect to time and dose.

Table SIII.55 displays the duration of exposure to pembrolizumab in the KN966 BTC population who received pembrolizumab combination therapy with respect to time and dose.

Table SIII.56 displays the duration of exposure to pembrolizumab in KN483 and KNA17 mesothelioma population who received pembrolizumab combination therapy with respect to time and dose.

Table SIII. 29: Clinical Trial Exposure by Dose and Duration Melanoma Subjects KN001, KN002 and KN006 Monotherapy (APaT Population)

Duration of Exposure	Pembrolizumab 2 mg/kg Q3W (N=340)		Pembrolizumab 10 mg/kg Q3W (N=769)		Pembrolizumab 10 mg/kg Q2W (N=458)		Total (N=1567)	
	n	Patient Years	n	Patient Years	n	Patient Years	n	Patient Years
> 0 m	340	214.1	769	473.0	458	290.1	1567	977.2
≥ 1 m	303	213.1	674	469.9	408	288.0	1385	971.0
≥ 3 m	206	196.1	492	437.2	310	270.9	1008	904.2
≥ 6 m	149	176.1	356	388.6	230	241.6	735	806.3
≥ 12 m	96	137.0	215	286.0	124	160.7	435	583.7

Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date +1.
 (pembrolizumab KN001 Database Cutoff Date: 18APR2014).
 (pembrolizumab KN002 Database Cutoff Date: 28FEB2015).
 (pembrolizumab KN006 Database Cutoff Date: 03MAR2015).

Data on File

Table SIII.30: Clinical Trial Exposure by Dose and Duration Melanoma Participants in KN054 and KN716 Monotherapy (APaT Population)

	KN716 + KN054 Pembrolizumab Q3W (N=992)		
	n	(%)	Person-years
Duration of Exposure			
>0 m	992	(100.0)	(725.4)
>=1 m	944	(95.2)	(723.5)
>=3 m	862	(86.9)	(708.7)
>=6m	724	(73.0)	(656.6)
>=12m	98	(9.9)	(104.2)
Each participant is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. Database cutoff date for Melanoma (KN054: 02OCT2017, KN716: 04DEC2020)			

Data on File

Table SIII.31: Clinical Trial Exposure by Dose and Duration NSCLC Subjects KN001, KN010, KN024 and KN042 Monotherapy (APaT Population)

Duration of Exposure	Pembrolizumab 2 mg/kg Q3W (N=400)			Pembrolizumab 10 mg/kg Q3W (N=630)			Pembrolizumab 10 mg/kg Q2W (N=202)			Pembrolizumab 200 mg Q3W (N=790)			Total (N=2022)		
	n	(%)	Person-years	n	(%)	Person-years	n	(%)	Person-years	n	(%)	Person-years	n	(%)	Person-years
>0 m	400	(100.0)	(158.8)	630	(100.0)	(284.2)	202	(100.0)	(97.5)	790	(100.0)	(582.0)	2022	(100.0)	(1122.5)
>=1 m	326	(81.5)	(156.1)	513	(81.4)	(280.1)	170	(84.2)	(96.5)	671	(84.9)	(578.8)	1680	(83.1)	(1111.5)
>=3 m	217	(54.3)	(139.1)	325	(51.6)	(250.1)	106	(52.5)	(86.2)	521	(65.9)	(555.5)	1169	(57.8)	(1030.9)
>=6m	123	(30.8)	(105.2)	220	(34.9)	(211.6)	75	(37.1)	(74.8)	393	(49.7)	(508.4)	811	(40.1)	(900.0)
>=12m	37	(9.3)	(48.3)	91	(14.4)	(120.9)	37	(18.3)	(47.3)	250	(31.6)	(406.3)	415	(20.5)	(622.8)
Each patient is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. (pembrolizumab KN001 Database Cutoff Date for Lung: 23JAN2015). (pembrolizumab KN010 Database Cutoff Date: 30SEP2015). (pembrolizumab KN024 Database Cutoff Date: 10JUL2017). (pembrolizumab KN042 Database Cutoff Date: 26FEB2018).															

Data on File

Table SIII.32: Clinical Trial Exposure to Pembrolizumab by Dose and Duration NSCLC Subjects KN021 (Cohorts A, C and G), KN189 and KN407 Combination Therapy (APaT Population)

	Pembrolizumab 10mg/kg Combo (N=24)			Pembrolizumab 2mg/kg Combo (N=25)			Pembrolizumab 200mg Combo (N=742)			Total (N=791)		
	n	(%)	Person-years	n	(%)	Person-years	n	(%)	Person-years	n	(%)	Person-years
Duration of Exposure												
>0 m	24	(100.0)	(16.1)	25	(100.0)	(26.5)	742	(100.0)	(452.2)	791	(100.0)	(494.8)
>=1 m	20	(83.3)	(15.9)	24	(96.0)	(26.4)	669	(90.2)	(449.6)	713	(90.1)	(491.9)
>=3 m	17	(70.8)	(15.5)	22	(88.0)	(26.1)	585	(78.8)	(434.7)	624	(78.9)	(476.2)
>=6m	11	(45.8)	(13.1)	17	(68.0)	(24.3)	406	(54.7)	(367.8)	434	(54.9)	(405.3)
>=12m	6	(25.0)	(9.2)	11	(44.0)	(20)	120	(16.2)	(156.8)	137	(17.3)	(186.0)
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. MK-3475 Database Cutoff Date for Lung Combination Therapy (KN021 Cohort A: 07NOV2016, Cohort G/C: 31MAY2017, KN189: 8NOV2017, KN407: 03APR2018)												

Data on File

Table SIII.33: Clinical Trial Exposure by Dose and Duration NSCLC Participants KN091 Monotherapy (APaT Population)

	Pembrolizumab 200 mg Q3W (N=580)		
	n	(%)	Person-Years
Duration of Exposure			
>0 m	580	(100.0)	(418.7)
>=1 m	538	(92.8)	(417.4)
>=3 m	461	(79.5)	(403.9)
>=6m	397	(68.4)	(380.1)
>=12m	114	(19.7)	(122.3)
Each participant is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. KN091 Database Cutoff Date: 20SEP2021			

Data on File

Table SIII. 34: Clinical Trial Exposure by Dose and Duration NSCLC Participants in KN671 Combination Therapy (APaT Population)

	Pembrolizumab 200 mg Q3W (N=396)		
	n	(%)	Person-Years
Duration of Exposure			
>0 m	396	(100.0)	(307.4)
>=1 m	372	(93.9)	(306.5)
>=3 m	292	(73.7)	(293.1)
>=6m	281	(71.0)	(288.7)
>=12m	185	(46.7)	(218.2)
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. Database Cutoff Date: 29JUL2022			

Data on File

Table SIII.35: Clinical Trial Exposure by Dose and Duration NSCLC Participants in MK-3475A-D77 Combination Therapy (APaT Population)

	Pembrolizumab SC 790 mg Q6W (N=251)		
	n	(%)	Person-Years
Duration of Pembrolizumab Exposure			
>0 m	251	(100.0)	(307.4)
>=1 m	227	(90.4)	(306.5)
>=3 m	203	(80.9)	(293.1)
>=6m	146	(58.2)	(288.7)
>=12m	8	(3.2)	(218.2)
Each participant is counted once on each applicable duration category row. Duration of exposure (months) is calculated as (last dose date - first dose date + 1) / 30.4367 for pembrolizumab. Database Cutoff Date for Lung (MK-3475A-D77: 12JUL2024).			

Data on File

Table SIII.36: Clinical Trial Exposure by Dose and Duration HL Subjects in KN013, KN087, and KN204 Monotherapy (APaT Population)

	Pembrolizumab 10 mg/kg Q2W (N=31)			Pembrolizumab 200 mg Q3W (N=358)			Total (N=389)		
	n	(%)	Person-years	n	(%)	Person-years	n	(%)	Person-years
Duration of Exposure									
>0 m	31	(100.0)	(27.7)	358	(100.0)	(378.1)	389	(100.0)	(405.8)
>=1 m	29	(93.5)	(27.7)	353	(98.6)	(378.0)	382	(98.2)	(405.7)
>=3 m	25	(80.6)	(27.1)	322	(89.9)	(372.2)	347	(89.2)	(399.3)
>=6m	18	(58.1)	(24.8)	253	(70.7)	(346.5)	271	(69.7)	(371.3)
>=12m	11	(35.5)	(19.7)	171	(47.8)	(288.5)	182	(46.8)	(308.2)
>=18m	8	(25.8)	(16.0)	111	(31.0)	(216.2)	119	(30.6)	(232.2)

Each subject is counted once on each applicable duration category row.
Duration of Exposure is calculated as last dose date - first dose date + 1.
Database Cutoff Date (KN013-Cohort 3: 28SEP2018, KN087: 21MAR2019, KN204: 16JAN2020)

Data on File

Table SIII.37: Clinical Trial Exposure by Dose and Duration UC Subjects KN045 and KN052 Monotherapy (APaT Population)

Duration of Exposure	Pembrolizumab 200 mg Q3W (N=636)	
	n	Patient Years
> 0 m	636	236.2
>= 1 m	510	232.0
>= 3 m	296	195.3
>= 6 m	167	148.0
>= 12 m	52	65.1

Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date +1.
Includes all subjects with urothelial carcinoma who received at least one dose of Pembrolizumab in KN045 and KN052.
Pembrolizumab KN045 Database Cutoff Date: 07SEP2016.
Pembrolizumab KN052 Database Cutoff Date: 01SEP2016.

Data on File

Table SIII.38: Clinical Trial Exposure to Pembrolizumab by Dose and Duration Urothelial Cancer Participants in KNA39 and KN869 Pembrolizumab + EV Combination Therapy (APaT Population)

Duration of Exposure	Pembrolizumab (200 mg Q3W) + Enfortumab Vedotin (N=564)		
	n	(%)	Person-Years
>0 m	564	(100.0)	423.5
>=1 m	498	(88.3)	421.9
>=3 m	432	(76.6)	410.6
>=6m	327	(58.0)	371.2
>=12m	168	(29.8)	254.9

Each participant is counted once on each applicable duration category row.
Duration of exposure is calculated as last dose date - first dose date + 1 for pembrolizumab in the combination.
Pembrolizumab KN-869 Database Cutoff Date: 13Mar2023, KN-A39 Database Cutoff Date: 08Aug2023

Data on File

Table SIII.39: Clinical Trial Exposure by Dose and Duration HNSCC Subjects KN012 B+B2 Cohorts, KN040, KN055 and KN048 Monotherapy (ASaT Population)

	Pembrolizumab 10mg/kg Q2W (N=60)			Pembrolizumab 200mg Q3W (N=849)			Total (N=909)		
	n	(%)	Person-years	n	(%)	Person-years	n	(%)	Person-years
Duration of Exposure									
>0 m	60	(100.0)	(34.2)	849	(100.0)	(372.3)	909	(100.0)	(406.5)
≥1 m	53	(88.3)	(34.0)	681	(80.2)	(366.9)	734	(80.7)	(400.9)
≥3 m	34	(56.7)	(31.0)	425	(50.1)	(324.6)	459	(50.5)	(355.5)
≥6m	19	(31.7)	(25.9)	262	(30.9)	(266.0)	281	(30.9)	(291.8)
≥12m	12	(20.0)	(21.1)	106	(12.5)	(161.0)	118	(13.0)	(182.1)
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. MK-3475 Database Cutoff Date for HNSCC (KN012-HNSCC: 26APR2016, KN040: 15MAY2017, KN048: 13JUN2018, KN055: 22APR2016)									

Data on File

Table SIII.40: Clinical Trial Exposure to Pembrolizumab by Dose and Duration HNSCC Subjects in KN048 Combination Therapy (ASaT Population)

	Pembrolizumab 200mg Q3W (N=276)		
	n	(%)	Person-years
Duration of Exposure			
>0 m	276	(100.0)	(170.0)
≥1 m	239	(86.6)	(168.9)
≥3 m	198	(71.7)	(161.8)
≥6m	129	(46.7)	(135.2)
≥12m	51	(18.5)	(81.0)
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. MK-3475 Database Cutoff Date for KN048: 13JUN2018.			

Data on File

Table SIII.41: Clinical Trial Exposure to Pembrolizumab by Dose and Duration HNSCC Participants in KN689 Monotherapy or Combination Therapy (APaT Population)

	Pembrolizumab 200mg Q3W (N=361)		
	n	(%)	Person-years
Duration of Exposure			
>0 m	361	(100.00)	(228.20)
≥1 m	255	(70.64)	(222.54)
≥3 m	245	(67.87)	(220.49)
≥6 m	212	(58.73)	(209.24)
≥12 m	142	(39.34)	(155.04)
Each participant is counted once on each applicable duration category row. Duration of exposure is (months) calculated as (last dose date - first dose date + 1) /30.4367 for pembrolizumab Database cutoff date for KN689: 25JUL2024			

Table SIII.42: Clinical Trial Exposure to Pembrolizumab by Dose and Duration RCC Subjects KN426 Combination Therapy (ASaT Population)

	KN426 Data for Pembrolizumab + Axitinib (N=429)		
	n	(%)	Person-years
Duration of Exposure			
>0 m	429	(100.0)	(337.1)
>=1 m	395	(92.1)	(335.7)
>=3 m	346	(80.7)	(327.7)
>=6m	300	(69.9)	(311.0)
>=12m	152	(35.4)	(200.4)
>=18m	32	(7.5)	(50.9)

Each subject is counted once on each applicable duration category row.
Duration of Exposure is calculated as (last dose date - first dose date +1)/30.4367 (months).
Duration of Exposure is calculated as last dose date - first dose date + 1 for pembrolizumab in the combo.
MK-3475 Database Cutoff Date for KN426: 24AUG2018.

Data on File

Table SIII.43: Clinical Trial Exposure to Pembrolizumab by Dose and Duration RCC Participants KN581 Combination Therapy (APaT Population)

Duration of Exposure	Pembrolizumab + Lenvatinib(N=352)		
	n	(%)	Person-years
>0 m	352	(100.0)	(423.9)
>=1 m	326	(92.6)	(423.2)
>=3 m	301	(85.5)	(418.9)
>=6 m	272	(77.3)	(407.7)
>=12 m	213	(60.5)	(363.8)
>=18 m	140	(39.8)	(272.2)

Each participant is counted once on each applicable duration category row.
Duration of Exposure is calculated as last dose date - first dose date + 1 for pembrolizumab in the combo.MK-3475
Database Cutoff Date for KN581: 28AUG2020.

Data on File

Table SIII.44: Clinical Trial Exposure by Dose and Duration RCC Participants KN564 Monotherapy (APaT Population)

	Pembrolizumab 200 mg Q3W (N=488)		
	n	(%)	Person-years
Duration of Exposure			
>0 m	488	(100.0)	(364.3)
>=1 m	463	(94.9)	(363.1)
>=3 m	428	(87.7)	(357.0)
>=6 m	370	(75.8)	(334.4)
>=9 m	325	(66.6)	(306.7)
>=12 m	32	(6.6)	(34.6)
Each participant is counted once on each applicable duration category row. Duration of Exposure (months) is calculated as (last dose date - first dose date + 1)/30.4367. Includes all participants who received at least one dose of pembrolizumab in KN564. Database cutoff date for RCC (KN564: 14DEC2020)			

Data on File

Table SIII.45: Clinical Trial Exposure by Dose and Duration CRC Subjects KN177 Monotherapy (ASaT Population)

	KN177 Data for Pembrolizumab ^{††} (N=153)		
	n	(%)	Person-years
Duration of Exposure			
>0 m	153	(100.0)	(169.0)
>=1 m	134	(87.6)	(168.5)
>=3 m	112	(73.2)	(165.0)
>=6m	96	(62.7)	(158.7)
>=12m	73	(47.7)	(141.7)
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. ^{††} Includes all subjects who received at least one dose of pembrolizumab in KN177. Database cutoff date for CRC (KN177: 19FEB2020)			

Data on File

Table SIII.46: Clinical Trial Exposure to Pembrolizumab by Dose and Duration Esophageal Cancer Subjects KN590 Combination Therapy (ASaT Population)

	KN590 Data for Pembrolizumab (N=370)		
	n	(%)	Person-years
Duration of Exposure			
>0 m	370	(100.0)	(231.5)
>=1 m	323	(87.3)	(230.3)
>=3 m	266	(71.9)	(220.6)
>=6m	161	(43.5)	(180.9)
>=12m	76	(20.5)	(122.3)
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as (last dose date - first dose date +1)/30.4367 (months). Database Cutoff date for Esophageal (KN590: 02JUL2020)			

Data on File

Table SIII.47: Clinical Trial Exposure to Pembrolizumab by Dose and Duration TNBC Subjects KN355 Combination Therapy (ASaT Population)

	Pembrolizumab (200 mg Q3W) + Chemotherapy (N=596)		
	n	(%)	Person-years
Duration of Pembrolizumab Exposure			
>0 m	596	(100.0)	(405.3)
>=1 m	550	(92.3)	(403.9)
>=3 m	443	(74.3)	(385.8)
>=6 m	291	(48.8)	(331.3)
>=12 m	146	(24.5)	(229.6)
Each subject is counted once on each applicable duration category row. Duration of exposure is calculated as (last dose date - first dose date +1)/30.4367 (months) for pembrolizumab in the combo. Database Cutoff Date for KN355: 11DEC2019.			

Data on File

Table SIII.48: Clinical Trial Exposure to Pembrolizumab by Dose and Duration TNBC Subjects KN522 Combination Therapy (APaT Population)

	Pembrolizumab 200 mg Q3W (N=783)		
	n	(%)	Person-years
Duration of Pembrolizumab Exposure			
>0 m	778	(99.4)	(709.7)
>=1 m	741	(94.6)	(708.2)
>=3 m	669	(85.4)	(695.2)
>=6 m	560	(71.5)	(653.5)
>=12 m	503	(64.2)	(607.7)
Each participant is counted once on each applicable duration category row. Duration of exposure is the time from the first dose date to the last dose date for pembrolizumab. Database cutoff date for TNBC: (KN522: 23MAR2021)			

Data on File

Table SIII.49: Clinical Trial Exposure to Pembrolizumab by Dose and Duration Endometrial Carcinoma Participants KN146 and KN775 Combination Therapy (APaT Population)

	Pembrolizumab 200mg Q3W (N=530)		
	n	(%)	Person-years
Duration of Exposure (month)			
> 0 m	530	(100.0)	379.8
≥ 1 m	472	(89.1)	378.1
≥ 3 m	394	(74.3)	364.4
≥ 6 m	301	(56.8)	330.6
≥ 12 m	141	(26.6)	214.6
≥ 18 m	70	(13.2)	127.4
≥ 24 m	8	(1.5)	16.8
Each participant is counted once on each applicable duration category row. Duration of exposure (month) is calculated as (last dose date - first dose date + 1) / 30.4367. Database cutoff date for endometrial carcinoma (KN146: 18AUG2020, KN775: 26OCT2020)			

Data on File

Table SIII.50: Clinical Trial Exposure to Pembrolizumab by Dose and Duration Endometrial Carcinoma Participants in KN868 Combination Therapy (APaT Population)

	KN868 Pembrolizumab + Chemotherapy		
	(N=382)	(%)	Person-years
Duration of Pembrolizumab Exposure			
> 0 m	382	(100.0)	(222.0)
≥ 1 m	341	(89.3)	(221.0)
≥ 3 m	288	(75.4)	(211.9)
≥ 6 m	176	(46.1)	(169.5)
≥ 12 m	66	(17.3)	(92.3)
≥ 18 m	25	(6.5)	(43.3)
Each Participants is counted once on each applicable duration category row. Duration of Exposure is calculated as (last dose date - first dose date + 1)/30.4367. Dose (Pembrolizumab): 200mg Q3W in Stage 1 and 400mg Q6W in Stage 2. Database cutoff date (KN868: 16DEC2022 for dMMR participants and 06DEC2022 for pMMR participants)			

Data on File

Table SIII.51: Clinical Trial Exposure by Dose and Duration MSI-H Subjects KN158 Cohort K and KN164 Cohorts A and B Monotherapy (ASaT Population)

	Pembrolizumab 200 mg Q3W (N=475)		
	n	(%)	Person-years
Duration of Exposure			
>0 m	475	(100.0)	(384.2)
≥1 m	405	(85.3)	(382.3)
≥3 m	305	(64.2)	(365.5)
≥6m	227	(47.8)	(337.9)
≥12m	161	(33.9)	(290.9)
≥18m	121	(25.5)	(241.1)
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. Database cutoff date for MSI-H (KN158-cohort K: 05OCT2020) Database cutoff date for Colorectal (KN164-cohorts A and B: 09SEP2019)			

Data on File

Table SIII.52: Clinical Trial Exposure to Pembrolizumab by Dose and Duration Cervical Cancer Participants in KNA18 + KN826 Combination Therapy (APaT Population)

	Pembrolizumab 200 mg Q3W (N=835)		
	n	(%)	Person-years
Duration of Exposure			
>0 m	835	(100.00)	(822.28)
>=1 m	775	(92.81)	(820.37)
>=3 m	709	(84.91)	(809.33)
>=6 m	589	(70.54)	(765.61)
>=12 m	389	(46.59)	(621.23)
Each participant is counted once on each applicable duration category row. Duration of exposure is calculated as last dose date - first dose date + 1 for pembrolizumab in the combo. Database cutoff date for Cervical Cancer (KNA18: 09JAN2023, KN826: 03MAY2021).			

Data on File

Table SIII.53: Clinical Trial Exposure to Pembrolizumab by Dose and Duration Gastric Cancer Participants KN811 Combination Therapy (APaT Population)

	Pembrolizumab 200 mg Q3W (N=350)		
	n	(%)	Person-Years
Duration of Exposure			
>0 m	350	(100.0)	(315.0)
>=1 m	323	(92.3)	(314.0)
>=3 m	286	(81.7)	(307.3)
>=6m	231	(66.0)	(286.4)
>=12m	133	(38.0)	(214.1)
Each subject is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. Database Cutoff Date: 25MAY2022			

Data on file

Table SIII.54: Clinical Trial Exposure to Pembrolizumab by Dose and Duration Gastric Cancer Participants KN859 Combination Therapy (APaT Population)

	Pembrolizumab 200 mg Q3W (N=785)		
	n	(%)	Person-years
Duration of exposure (month)			
>0	785	(100.0)	555.6
>=1	688	(87.6)	552.9
>=3	589	(75.0)	535.7
>=6	405	(51.6)	467.4
>=12	186	(23.7)	314.0

Each participant is counted once on each applicable duration category row.
Duration of exposure (month) is calculated as (last dose date - first dose date + 1) / 30.4367
Database cutoff date for Gastric (KN859: 03OCT2022)

Data on File

Table SIII.55: Clinical Trial Exposure to Pembrolizumab by Dose and Duration BTC Participants in KN966 Combination Therapy (APaT Population)

	KN966 Pembrolizumab (200mg Q3W) + Chemotherapy (N=529)		
	n	(%)	Person-Years
Duration of Exposure			
> 0 m	529	(100.0)	(338.7)
≥ 1 m	464	(87.7)	(337.0)
≥ 3 m	376	(71.1)	(322.2)
≥ 6 m	267	(50.5)	(282.5)
≥ 12 m	113	(21.4)	(172.9)
≥ 18 m	57	(10.8)	(106.0)
≥ 24 m	20	(3.8)	(41.9)

Each participant is counted once on each applicable duration category row.
Duration of exposure is calculated as last dose date - first dose date + 1
(pembrolizumab KN966 Database Cutoff Date: 15DEC2022)

Data on File

Table SIII.56: Clinical Trial Exposure to Pembrolizumab by Dose and Duration Mesothelioma Participants in KN483 and KNA17 Combination Therapy (APaT Population)

	Pembrolizumab 200 mg Q3W (N=260)		
	n	(%)	Person-years
Duration of Exposure			
>0 m	260	(100.0)	(199.7)
>=1 m	246	(94.6)	(199.2)
>=3 m	206	(79.2)	(191.9)
>=6m	157	(60.4)	(173.4)
>=12m	74	(28.5)	(117.6)
Each participant is counted once on each applicable duration category row. Duration of Exposure is calculated as last dose date - first dose date + 1. Database Cutoff Date for Thoracic Malignancy (KN483: 16SEP2022, KNA17: 21SEP2022).			

Data on File

Clinical Trial Exposure by Age and Gender

Clinical trial exposure to pembrolizumab monotherapy by age and gender for all melanoma patients in the pooled KN001, KN002 and KN006 population are summarized in Table SIII.57. Overall, a greater number of patients less than 65 years have been treated compared to 65 years or older (906 vs. 661). Across the age groups, more males have been treated compared to females (951 vs. 616).

Clinical trial exposure to pembrolizumab monotherapy by age and gender for all melanoma patients in KN054 and KN716 is summarized in Table SIII.58. Overall, a greater number of patients less than 65 years (one patient ≥ 12 and ≤ 18 years of age) have been treated compared to 65 years or older (684 vs. 308). Across the age groups, more males have been treated compared to females (617 vs. 375).

Clinical trial exposure to pembrolizumab monotherapy by age and gender for all NSCLC patients in KN001, KN010, KN024 and KN042 is summarized in Table SIII.59. Overall, a greater number of patients less than 65 years have been treated compared to 65 years or older (1,116 vs. 906). Across the age groups, more males have been treated compared to females (1,249 vs. 773).

Clinical trial exposure to pembrolizumab by age and gender for all NSCLC patients who received pembrolizumab in combination with chemotherapy in KN189, KN021 (Cohorts A, C and G) and KN407 is summarized in Table SIII.60. Overall, a similar number of patients less than 65 years have been treated compared to 65 years or older (382 vs. 409). Across the age groups, more males have been treated compared to females (516 vs. 275).

Clinical trial exposure to pembrolizumab monotherapy by age and gender for NSCLC patients in KN091 is summarized in Table SIII.61. Overall, a similar number of patients less than 65 years have been treated compared to 65 years or older (281 vs. 299). Across the age groups, more males have been treated compared to females (394 vs. 186).

Clinical trial exposure to pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy by age and gender for all NSCLC patients in KN671 is summarized in Table SIII.62. Overall, a greater number of patients less than 65 years have been treated compared to 65 years or older (221 vs 175). Across the age groups, more males have been treated compared to females (278 vs 118).

Clinical trial exposure to pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy by age and gender for all NSCLC patients in MK-3475A-D77 is summarized in Table SIII.63. Overall, a similar number of patients less than 65 years have been treated compared to 65 years or older (119 vs 132). Across the age groups, more males have been treated compared to females (182 vs 69).

Clinical trial exposure to pembrolizumab monotherapy by age and gender for all HL patients in KN013, KN087, and KN204 is summarized in Table SIII.64. Overall, a greater number of patients less than 65 years have been treated compared to 65 years or older (343 vs. 46). Across the age groups, more males have been treated compared to females (212 vs. 177).

Clinical trial exposure to pembrolizumab monotherapy by age and gender for patients under 18 years in KN051, of which 22 were HL patients, is summarized in Table SIII.65. Of the age groups, there were 3 patients 6 months to less than 2 years, 23 patients 2 to 5 years, 25 patients 6 to 9 years, 36 patients 10 to 13 years, and 74 patients 14 to 17 years. Across the age groups, a similar number of males and females have been treated (82 vs. 79).

Clinical trial exposure to pembrolizumab monotherapy by age and gender for all UC patients in KN045 and KN052 is summarized in Table SIII.66. Overall, a greater number of patients 65 years or older have been treated compared to patients less than 65 years (465 vs. 171). Across the age groups, more males have been treated compared to females (484 vs. 152).

Clinical trial exposure to pembrolizumab by age and gender for all UC patients in KNA39 and KN869 who received pembrolizumab + EV combination therapy is summarized in Table SIII.67. Overall, a greater number of patients 65 years or older have been treated compared to patients less than 65 years (391 vs. 173). Across the age groups, more males have been treated compared to females (432 vs. 132).

Clinical trial exposure to pembrolizumab monotherapy by age and gender for all HNSCC patients in KN012 B+B2 Cohorts, KN040, KN055, and KN048 is summarized in Table SIII.68. Overall, a greater number of patients less than 65 years have been treated compared to 65 years or older (586 vs.323). Across the age groups, more males have been treated compared to females (752 vs.157).

Clinical trial exposure to pembrolizumab by age and gender for HNSCC patients in KN048 who received pembrolizumab combination therapy is summarized in Table SIII.69. Overall, a greater number of patients less than 65 years have been treated compared to 65 years or older (176 vs.100). Across the age groups, more males have been treated compared to females (220 vs. 56).

Clinical trial exposure to pembrolizumab by age and gender for HNSCC patients in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy is summarized in Table SIII.70. Overall, a greater number of patients less than 65 years have been treated compared to 65 years or older (247 vs.114). Across the age groups, more males have been treated compared to females (284 vs.77)

Clinical trial exposure to pembrolizumab by age and gender for all RCC patients in KN426 who received pembrolizumab combination therapy is summarized in Table SIII.71. Overall, a greater number of patients less than 65 years have been treated compared to 65 years or older (257 vs. 172). Across the age groups, more males have been treated compared to females (306 vs. 123).

Clinical trial exposure to pembrolizumab by age and gender for all RCC patients in KN581 who received pembrolizumab combination therapy is summarized in Table SIII.72. Overall, a greater number of patients less than 65 years have been treated compared to 65 years or older (193 vs. 159). Across the age groups, more males have been treated compared to females (252 vs. 100).

Clinical trial exposure to pembrolizumab monotherapy by age and gender for all RCC patients in KN564 is summarized in Table SIII.73. Overall, a greater number of patients less than 65 years have been treated compared to 65 years or older (333 vs. 155). Across the age groups, more males have been treated compared to females (340 vs. 148).

Clinical trial exposure to pembrolizumab monotherapy by age and gender for all CRC patients in KN177 is summarized in Table SIII.74. Overall, a greater number of patients 65 years or older have been treated compared to patients less than 65 years (80 vs. 73). Across the age groups, more females have been treated compared to males (82 vs. 71). Of note, the absolute difference is small as compared to a number of other datasets.

Clinical trial exposure to pembrolizumab by age and gender for all esophageal cancer patients in KN590 who received pembrolizumab combination therapy is summarized in Table SIII.75. Overall, a greater number of patients less than 65 years have been treated compared to patients 65 years or older (201 vs. 169). Across the age groups, more males have been treated compared to females (305 vs. 65).

Clinical trial exposure to pembrolizumab by age and gender for all TNBC patients in KN355 who received pembrolizumab combination therapy is summarized in Table SIII.76. Overall, a greater number of patients less than 65 years have been treated compared to 65 years or older (459 vs. 137). Only female patients were included in KN355.

Clinical trial exposure to pembrolizumab by age and gender for the TNBC patients in KN522 who received pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy is summarized in Table SIII.77. Overall, a greater number of patients less than 65 years have been treated compared to 65 years or older (695 vs. 83). Across the age groups, more females have been treated compared to males (777 vs. 1).

Clinical trial exposure to pembrolizumab by age for all endometrial carcinoma patients in KN146 and KN775 who received pembrolizumab combination therapy is summarized in Table SIII.78. Overall, a similar number of patients less than 65 years have been treated compared to 65 years or older (252 vs. 278). Only female patients were included in KN146 and KN775.

Clinical trial exposure to pembrolizumab by age for all endometrial carcinoma patients in KN868 who received pembrolizumab combination therapy is summarized in Table SIII.79. Overall, a greater number of patients 65 years or older have been treated compared to patients less than 65 years (210 vs. 172). Only female patients were included in KN868.

Clinical trial exposure to pembrolizumab monotherapy by age and gender for all MSI-H patients in KN158 Cohort K and KN164 Cohorts A and B is summarized in Table SIII.80. Overall, a greater number of patients less than 65 years have been treated compared to patients 65 years or older (305 vs. 170). Across the age groups, more females have been treated compared to males (262 vs. 213).

Clinical trial exposure to pembrolizumab by age and gender for the cervical cancer patients in KNA18 and KN826 who received pembrolizumab combination therapy is summarized in Table SIII.81. Overall, a greater number of patients less than 65 years have been treated compared to patients 65 years or older (731 vs. 104). Only female patients were included in KNA18 and KN826.

Clinical trial exposure to pembrolizumab by age and gender for gastric cancer patients in KN811 who received pembrolizumab combination therapy is summarized in Table SIII.82. Overall, a greater number of patients less than 65 years have been treated compared to 65 years or older (205 vs. 145). Across the age groups, more males have been treated compared to females (284 vs. 66).

Clinical trial exposure to pembrolizumab by age and gender for gastric cancer patients in KN859 who received pembrolizumab combination therapy is summarized in Table SIII.83. Overall, a greater number of patients less than 65 years have been treated compared to 65 years or older (483 vs. 302). Across the age groups, more males have been treated compared to females (522 vs. 263).

Clinical trial exposure to pembrolizumab by age and gender for all BTC patients who received pembrolizumab combination therapy in KN966 is summarized in Table SIII.84. Overall, a similar number of patients less than 65 years and patients 65 years or older were treated (266 vs. 263). Across the age groups, more males have been treated compared to females (279 vs. 250).

Clinical trial exposure to pembrolizumab by age and gender for all mesothelioma patients who received pembrolizumab combination therapy in KN483 and KNA17 is summarized in Table SIII.85. Overall, a greater number of patients 65 years or older were treated compared to patients less than 65 years (205 vs. 55). Across the age groups, more males have been treated compared to females (196 vs. 64).

Table SIII.57: Clinical Trial Exposure by Age and Gender Melanoma Subjects KN001, KN002 and KN006 Monotherapy (APaT Population)

Age Category (years)	Subjects			Mean Duration (days)			Subject Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	516	390	906	236.5	213.4	226.6	334.1	227.8	562.0
≥65	435	226	661	242.2	204.8	229.4	288.4	126.7	415.2
Total	951	616	1567	239.1	210.2	227.8	622.6	354.6	977.2

Duration of exposure is calculated assuming one day of dosing = 7 days of exposure.
Duration of Exposure is calculated as last dose date - first dose date +1.
(pembrolizumab KN001 Database Cutoff Date: 18APR2014).
(pembrolizumab KN002 Database Cutoff Date: 28FEB2015).
(pembrolizumab KN006 Database Cutoff Date: 03MAR2015).

Data on File

Table SIII.58: Clinical Trial Exposure by Age and Gender Melanoma Participants KN054 and KN716 Monotherapy (APaT Population)

Age Category (Years)	Participants			Mean Duration (days)			Participant Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65 *	412	272	684	271.3	279.1	274.4	306.0	207.8	513.8
≥65	205	103	308	250.9	250.8	250.9	140.8	70.7	211.6
Total	617	375	992	264.5	271.3	267.1	446.8	278.6	725.4

Duration of Exposure is calculated as last dose date - first dose date + 1.
*one patient in KN716 ≥12 and ≤ 18 years of age
Database cutoff date for Melanoma (KN054: 02OCT2017, KN716: 04DEC2020)

Data on File

Table SIII.59: Clinical Trial Exposure by Age and Gender NSCLC Subjects KN001, KN010, KN024 and KN042 Monotherapy (APaT Population)

Age Category (Years)	Subjects			Mean Duration (days)			Subject Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	662	454	1116	207.2	192.1	201.0	375.5	238.7	614.2
≥65	587	319	906	218.9	179.3	204.9	351.7	156.6	508.3
Total	1249	773	2022	212.7	186.8	202.8	727.2	395.3	1122.5

Duration of Exposure is calculated as last dose date - first dose date +1.
(pembrolizumab KN001 Database Cutoff Date for Lung: 23JAN2015).
(pembrolizumab KN010 Database Cutoff Date: 30SEP2015).
(pembrolizumab KN024 Database Cutoff Date: 10JUL2017).
(pembrolizumab KN042 Database Cutoff Date: 26FEB2018).

Data on File

Table SIII.60: Clinical Trial Exposure to Pembrolizumab by Age and Gender NSCLC Subjects KN021 (Cohorts A, C and G), KN189 and KN407 Combination Therapy (APaT Population)

Age Category (Years)	Subjects			Mean Duration (days)			Subject Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	231	151	382	207.6	264.7	230.2	131.3	109.4	240.7
≥65	285	124	409	216.1	251.8	226.9	168.6	85.5	254.1
Total	516	275	791	212.3	258.9	228.5	299.9	194.9	494.8

Duration of Exposure is calculated as last dose date - first dose date + 1.
 MK-3475 Database Cutoff Date for Lung Combination Therapy (KN021 Cohort A: 07NOV2016, Cohort G/C: 31MAY2017, KN189: 8NOV2017, KN407: 03APR2018)

Data on File

Table SIII.61: Clinical Trial Exposure by Age and Gender NSCLC Participants KN091 Monotherapy (APaT Population)

Age Category (Years)	Patients			Mean Duration (days)			Patient Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	191	90	281	272.3	270.0	271.5	142.4	66.5	208.9
≥65	203	96	299	255.2	258.7	256.3	141.8	68.0	209.8
Total	394	186	580	263.5	264.1	263.7	284.2	134.5	418.7

Duration of Exposure is calculated as last dose date - first dose date + 1.
 KN091 Database Cutoff Date: 20SEP2021

Data on File

Table SIII.62: Clinical Trial Exposure to Pembrolizumab by Age and Gender NSCLC Participants in KN671 Combination Therapy (APaT Population)

Age Category (Years)	Patients			Mean Duration (days)			Patient Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	150	71	221	285.3	303.7	291.3	117.2	59.0	176.2
≥65	128	47	175	272.9	276.4	273.8	95.6	35.6	131.2
Total	278	118	396	279.6	292.9	283.5	212.8	94.6	307.4

Duration of Exposure is calculated as last dose date - first dose date + 1.
 Database Cutoff Date: 29JUL2022

Data on File

Table SIII.63: Clinical Trial Exposure to Pembrolizumab SC by Age and Gender NSCLC Participants in MK-3475A-D77 Combination Therapy (APaT Population)

Age Category (Years)	Patients			Mean Duration (days)			Patient Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	86	33	119	194.1	185.1	191.6	45.7	16.7	62.4
≥65	96	36	132	191.4	209.2	196.3	50.3	20.6	70.9
Total	182	69	251	192.7	197.7	194.1	96.0	37.3	133.4

Duration of Exposure (days) is calculated as last dose date - first dose date +1 for pembrolizumab.
Database Cutoff Date for Lung (MK-3475A-D77: 12JUL2024).

Data on File

Table SIII.64: Clinical Trial Exposure by Age and Gender HL Subjects KN013, KN087, and KN204 Monotherapy (APaT Population)

Age Category (Years)	Subjects			Mean Duration (days)			Subject Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	187	156	343	359.9	427.0	390.4	184.3	182.4	366.7
≥65	25	21	46	302.0	321.9	311.1	20.7	18.5	39.2
Total	212	177	389	353.1	414.6	381.1	204.9	200.9	405.8

Duration of Exposure is calculated as last dose date - first dose date + 1.
Database Cutoff Date (KN013-Cohort 3: 28SEP2018, KN087: 21MAR2019, KN204: 16JAN2020)

Data on File

Table SIII.65: Clinical Trial Exposure by Age and Gender KN051 Parts I and II Monotherapy (APaT Population)

Age Category (years)	Persons*		Person Time (years)	
	Male	Female	Male	Female
6 months - <2 years	1	2	0.1	0.2
2 - 5 years	10	13	2.8	4.7
6 - 9 years	9	16	2.8	9.2
10 - 13 years	21	15	6.4	3.3
14 - 17 years	41	33	22.0	19.8

(Database Cutoff Date: 10JAN2020).
* HL n=22

Data on File

Table SIII.66: Clinical Trial Exposure by Age and Gender UC Subjects KN045 and KN052 Monotherapy (APaT Population)

Age Category (years)	Subjects			Mean Duration (days)			Subject Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	133	38	171	131.7	143.9	134.4	47.9	15.0	62.9
≥65	351	114	465	136.9	133.6	136.1	131.6	41.7	173.3
Total	484	152	636	135.5	136.2	135.6	179.5	56.7	236.2

Duration of exposure is calculated assuming one day of dosing = 7 days of exposure.
Duration of Exposure is calculated as last dose date - first dose date +1.
Includes all subjects with urothelial carcinoma who received at least one dose of Pembrolizumab in KN045 and KN052.
Pembrolizumab KN045 Database Cutoff Date: 07SEP2016.
Pembrolizumab KN052 Database Cutoff Date: 01SEP2016.

Data on File

Table SIII.67: Clinical Trial Exposure to Pembrolizumab by Age and Gender UC Participants KNA39 and KN869 Pembrolizumab + EV Combination Therapy (APaT Population)

Age Category (years)	Participants			Mean Duration (days)			Participant Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	141	32	173	10.16	9.80	10.09	119.38	26.14	145.52
≥65	291	100	391	8.57	8.42	8.53	207.80	70.20	278.01
Total	432	132	564	9.09	8.76	9.01	327.19	96.34	423.53

Duration of exposure is calculated as last dose date - first dose date + 1 for pembrolizumab in the combination.
Pembrolizumab KN-869 Database Cutoff Date: 13Mar2023, KN-A39 Database Cutoff Date: 08Aug2023

Table SIII.68: Clinical Trial Exposure by Age and Gender HNSCC Subjects KN012 B+B2 Cohorts, KN040, KN055 and KN048 Monotherapy (ASaT Population)

Age Category (Years)	Subjects			Mean Duration (days)			Subject Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	493	93	586	149.8	174.4	153.7	202.2	44.4	246.6
≥65	259	64	323	179.8	185.4	180.9	127.5	32.5	160.0
Total	752	157	909	160.1	178.9	163.3	329.6	76.9	406.5

Duration of Exposure is calculated as last dose date - first dose date +1.
MK-3475 Database Cutoff Date for HNSCC (KN012-HNSCC: 26APR2016, KN040: 15MAY2017, KN048: 13JUN2018, KN055: 22APR2016)

Data on File

Table SIII.69: Clinical Trial Exposure to Pembrolizumab by Age and Gender HNSCC Subjects in KN048 Combination Therapy (ASaT Population)

Age Category (Years)	Subjects			Mean Duration (days)			Subject Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	136	40	176	209.2	198.4	206.7	77.9	21.7	99.6
≥65	84	16	100	266.0	178.8	252.1	61.2	7.8	69.0
Total	220	56	276	230.9	192.8	223.2	139.1	29.6	168.6

Duration of Exposure is calculated as last dose date - first dose date +1.
MK-3475 Database Cutoff Date for KN048: 13JUN2018.

Data on File

Table SIII.70: Clinical Trial Exposure to Pembrolizumab by Age and Gender HNSCC Subjects in KN689 Monotherapy or Combination Therapy (APaT Population)

Age Category (Years)	Participants			Mean Duration (days)			Participant Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	200	47	247	248.7	246.1	248.2	136.2	31.7	167.9
≥65	84	30	114	213.8	135.7	193.3	49.2	11.1	60.3
Total	284	77	361	238.4	203.1	230.9	185.4	42.8	228.2

Duration of Exposure (days) is calculated as last dose date - first dose date +1 for pembrolizumab.
Database cutoff date for KN689: 25JUL2024

Table SIII.71: Clinical Trial Exposure to Pembrolizumab by Age and Gender RCC Subjects KN426 Combination Therapy (ASaT Population)

Age Category (Years)	Subjects			Mean Duration (days)			Subject Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	189	68	257	298.9	262.7	289.3	154.7	48.9	203.6
≥65	117	55	172	273.2	305.9	283.7	87.5	46.1	133.6
Total	306	123	429	289.0	282.0	287.0	242.2	95.0	337.1

Duration of Exposure is calculated as last dose date - first dose date + 1 for pembrolizumab in the combo.
MK-3475 Database Cutoff Date for KN426: 24AUG2018.

Data on File

Table SIII.72: Clinical Trial Exposure to Pembrolizumab by Age and Gender RCC Participants KN581 Combination Therapy (APaT Population)

Age Category (years)	Participants			Mean Duration (days)			Participant Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	142	51	193	479.8	456.0	473.5	186.6	63.7	250.2
≥65	110	49	159	426.1	337.6	398.9	128.3	45.3	173.6
Total	252	100	352	456.4	398.0	439.8	314.9	109.0	423.9

Duration of Exposure is calculated as last dose date – first dose date + 1 for pembrolizumab in the combo.MK-3475 Database Cutoff Date for KN581: 28AUG2020.

Data on File

Table SIII.73: Clinical Trial Exposure by Age and Gender RCC Participants KN564 Monotherapy (APaT Population)

Age Category (Years)	Participants			Mean Duration (days)			Person Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	239	94	333	287.0	272.5	282.9	187.8	70.1	258.0
≥65	101	54	155	241.4	267.4	250.5	66.8	39.5	106.3
Total	340	148	488	273.5	270.7	272.6	254.6	109.7	364.3

Duration of Exposure is calculated as last dose date - first dose date + 1.
Database cutoff date for RCC (KN564: 14DEC2020)

Table SIII.74: Clinical Trial Exposure by Age and Gender CRC Subjects in KN177 Monotherapy (AsaT Population)

Age Category (Years)	Subjects			Mean Duration (days)			Subject Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	46	34	80	490.9	405.4	454.6	61.8	37.7	99.6
≥65	25	48	73	399.0	320.4	347.3	27.3	42.1	69.4
Total	71	82	153	458.5	355.7	403.4	89.1	79.9	169.0

Duration of Exposure is calculated as last dose date - first dose date + 1.
Database cutoff date for CRC (KN177: 19FEB2020)

Data on File

Table SIII.75: Clinical Trial Exposure to Pembrolizumab by Age and Gender Esophageal Cancer Subjects KN590 Combination Therapy (ASaT Population)

Age Category (Years)	Subjects			Mean Duration (days)			Subject Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	165	36	201	233.2	225.5	231.9	105.4	22.2	127.6
≥65	140	29	169	238.8	235.9	238.3	91.5	18.7	110.3
Total	305	65	370	235.8	230.1	234.8	196.9	41.0	237.9

Duration of Exposure is calculated as last dose date - first dose date + 1.
Database Cutoff date for Esophageal (KN590: 02JUL2020)

Data on File

Table SIII.76: Clinical Trial Exposure to Pembrolizumab by Age and Gender TNBC Subjects KN355 Combination Therapy (ASaT Population)

Age Category (Years)	Subjects		Mean Duration (days)		Subject Time (years)	
	Female	Total	Female	Total	Female	Total
<65	459	459	236.5	236.5	297.3	297.3
≥65	137	137	288.2	288.2	108.1	108.1
Total	596	596	248.4	248.4	405.3	405.3

Duration of Exposure is calculated as last dose date - first dose date + 1 for pembrolizumab in the combo.
Database Cutoff Date for KN355: 11DEC2019.

Data on File

Table SIII.77: Clinical Trial Exposure to Pembrolizumab by Age and Gender TNBC Subjects KN522 Combination Therapy (APaT Population)

Age Category (Years)	Participants			Mean Duration (days)			Person Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	0	695	695	0.0	338.9	338.9	0.0	644.9	644.9
≥65	1	82	83	388.0	284.0	285.2	1.1	63.8	64.8
Total	1	777	778	388.0	333.1	333.2	1.1	708.6	709.7

Duration of exposure is the time from the first dose date to the last dose date for pembrolizumab.
Database cutoff date for TNBC: (KN522: 23MAR2021)

Data on File

Table SIII.78: Clinical Trial Exposure to Pembrolizumab by Age Endometrial Carcinoma Participants KN146 and KN775 Combination Therapy (APaT Population)

Age Category (year)	Participants	Mean Duration (days)	Participant Time (years)
<65	252	278.9	192.5
≥65	278	246.1	187.3
Total	530	261.7	379.8

Duration of exposure (day) is calculated as last dose date - first dose date + 1.
Database cutoff date for endometrial carcinoma (KN146: 18AUG2020, KN775: 26OCT2020)

Data on File

Table SIII.79: Clinical Trial Exposure to Pembrolizumab by Age Endometrial Carcinoma Participants KN868 Combination Therapy (APaT Population)

Age Category (years)	Participants	Mean Duration (days)	Participants Time (years)
<65	172	215.5	101.5
≥65	210	209.6	120.5
Total	382	212.3	222.0

Duration of Exposure is calculated as last dose date - first dose date +1 for pembrolizumab.
Database cutoff date (KN868: 16DEC2022 for dMMR participants and 06DEC2022 for pMMR participants)

Data on File

Table SIII.80: Clinical Trial Exposure by Age and Gender MSI-H Subjects KN158 Cohort K and KN164 Cohorts A and B Monotherapy (ASaT Population)

Age Category (Years)	Subjects			Mean Duration (days)			Subject Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	144	161	305	306.2	263.0	283.4	120.7	115.9	236.7
≥65	69	101	170	281.8	341.1	317.0	53.2	94.3	147.6
Total	213	262	475	298.3	293.1	295.4	174.0	210.3	384.2

Duration of Exposure is calculated as last dose date - first dose date + 1.
 Database cutoff date for MSI-H (KN158-cohort K: 05OCT2020)
 Database cutoff date for Colorectal (KN164-cohorts A and B: 09SEP2019)

Data on File

Table SIII.81: Clinical Trial Exposure to Pembrolizumab by Age Cervical Cancer Participants in KNA18 and KN826 Combination Therapy (APaT Population)

Age Category (Years)	Participants	Mean Duration (days)	Participant Time (years)
<65	731	359.1	718.7
≥65	104	363.7	103.5
Total	835	359.7	822.3

Duration of exposure is calculated as last dose date - first dose date +1 for pembrolizumab in the combo.
 Database cutoff date for Cervical Cancer (KNA18: 09JAN2023, KN826: 03MAY2021).

Data on file

Table SIII.82: Clinical Trial Exposure to Pembrolizumab by Age and Gender Gastric Cancer Participants KN811 Combination Therapy (APaT Population)

Age Category (Years)	Patients			Mean Duration (days)			Patient Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	169	36	205	324.8	395.9	337.3	150.3	39.0	189.3
≥65	115	30	145	304.4	362.9	316.5	95.8	29.8	125.6
Total	284	66	350	316.5	380.9	328.7	246.1	68.8	315.0

Duration of Exposure is calculated as last dose date – first dose date + 1.
 Database Cutoff Date: 25MAY2022

Data on file

Table SIII.83: Clinical Trial Exposure to Pembrolizumab by Age and Gender Gastric Cancer Participants KN859 Combination Therapy (APaT Population)

Age Category (years)	Participants			Mean Duration (days)			Participant Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	321	162	483	275.1	224.6	258.2	241.8	99.6	341.4
≥65	201	101	302	263.9	249.6	259.1	145.2	69.0	214.2
Total	522	263	785	270.8	234.2	258.5	387.0	168.7	555.6

Each participant is counted once on each applicable duration category row.
Duration of exposure is calculated as last dose date - first dose date + 1.
Database cutoff date for Gastric (KN859: 03OCT2022)

Data on File

Table SIII.84: Clinical Trial Exposure to Pembrolizumab by Age and Gender BTC Participants in KN966 Combination Therapy (APaT Population)

Age Category (Years)	Participants			Mean Duration (days)			Participants Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	142	124	266	243.9	209.3	227.8	94.8	71.0	165.9
≥65	137	126	263	234.6	246.0	240.0	88.0	84.9	172.9
Total	279	250	529	239.3	227.8	233.9	182.8	155.9	338.7

Duration of Exposure is calculated as last dose date - first dose date + 1 for pembrolizumab in the combo.
(pembrolizumab KN966 Database Cutoff Date: 15DEC2022)

Data on File

Table SIII.85: Clinical Trial Exposure to Pembrolizumab by Age and Gender Mesothelioma Participants in KN483 and KNA17 Combination Therapy (APaT Population)

Age Category (Years)	Participants			Mean Duration (days)			Participant Time (years)		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
<65	38	17	55	242.9	291.5	258.0	25.3	13.6	38.8
≥65	158	47	205	284.2	294.6	286.6	122.9	37.9	160.8
Total	196	64	260	276.2	293.8	280.5	148.2	51.5	199.7

Duration of Exposure is calculated as last dose date - first dose date +1.
Database Cutoff Date for Thoracic Malignancy (KN483: 16SEP2022, KNA17: 21SEP2022).

Data on File

Clinical Trial Exposure by Racial / Ethnic Origin

Clinical trial exposure to pembrolizumab monotherapy by race for all melanoma patients in the pooled KN001, KN002 and KN006 population is summarized in Table SIII.86. The majority of patients treated have been White (1530) followed by Asian (19) and Black or African American (7).

Clinical trial exposure to pembrolizumab monotherapy by race for all melanoma patients in KN054 and KN716 is summarized in Table SIII.87. The majority of patients treated have been White (432) followed by Asian (4), Black or African American (3), American Indian or Alaska

Native (1) and Multiracial (1). Race was not reported for 551 patients as the majority of patients were from KN054 where information on race was not collected due to the European Organisation for Research and Treatment of Cancer (EORTC) standards.

Clinical trial exposure to pembrolizumab monotherapy by race for all NSCLC patients in KN001, KN010, KN024 and KN042 is summarized in Table SIII.88. The majority of patients treated have been White (1,466) followed by Asian (428) and Black or African American (53).

Clinical trial exposure to pembrolizumab by race for all NSCLC patients in KN189, KN021 (Cohorts A, C and G) and KN407 who received pembrolizumab in combination with chemotherapy is summarized in Table SIII.89. The majority of patients treated have been White (681) followed by Asian (73), Black or African American (30), American Indian or Alaska Native (1) and Native Hawaiian or Other Pacific Islander (1). Five (5) patients did not report race.

Clinical trial exposure to pembrolizumab monotherapy by race for all NSCLC patients in KN091 is summarized in Table SIII.90. The majority of patients treated have been White (442) followed by Asian (107), Other (5), Multiracial (4), American Indian or Alaska Native (1). Twenty-one (21) patients did not report race.

Clinical trial exposure to pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy by race for all NSCLC patients in KN671 is summarized in Table SIII.91. The majority of patients treated have been White (249), followed by Asian (124), Black or African American (6), Multiracial (3), and American Indian or Alaska Native (1). Thirteen (13) patients did not report race.

Clinical trial exposure to pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy by race for all NSCLC patients in MK-3475A-D77 is summarized in Table SIII.92. The majority of patients treated have been White (158), followed by Asian (74), Multiracial (12), Black or African American (5), and American Indian or Alaska Native (2).

Clinical trial exposure to pembrolizumab monotherapy by race for all HL patients in KN013, KN087, and KN204 is summarized in Table SIII.93. The majority of patients treated have been White (330) followed by Asian (25), Black or African American (11), Multiracial (7), American Indian or Alaska Native (2), and Native Hawaiian or Other Pacific Islander (1). Thirteen (13) patients did not report race.

Clinical trial exposure to pembrolizumab monotherapy by race for all UC patients in KN045 and KN052 is summarized in Table SIII.94. The majority of patients treated have been White (512) followed by Asian (90) and Black or African American (13).

Clinical trial exposure to pembrolizumab by race for all UC patients in KNA39 and KN869 who received pembrolizumab + EV combination therapy is summarized in Table SIII.95. The majority of patients treated have been White (410), followed by Asian (107), Black or African American (9), Other (5) and American Indian or Alaska Native (2). Thirty-one (31) patients did not report race.

Clinical trial exposure to pembrolizumab monotherapy by race for all HNSCC patients in KN012 B+B2 Cohorts, KN040, KN055 and KN048 is summarized in Table SIII.96. The majority of patient treated have been White (723) followed by Asians (109) and Black or African American (28).

Clinical trial exposure to pembrolizumab by race for HNSCC patients in KN048 who received pembrolizumab combination therapy is summarized in Table SIII.97. The majority of patients treated have been White (200) followed by Asian (60) and Black or African American (10).

Clinical trial exposure to pembrolizumab by race for HNSCC patients in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy is summarized in Table SIII.98. The majority of patients treated have been White (284) followed by Asian (50), Multiracial (9), Black or African American (8), and Native Hawaiian or Other Pacific Islander (1). Race was not reported for nine (9) patients.

Clinical trial exposure to pembrolizumab by race for all RCC patients in KN426 who received pembrolizumab combination therapy is summarized in Table SIII.99. The majority of patients treated have been White (342) followed by Asians (66); Black or African American (9); American Indian or Alaska Native (2); Multiracial (1); Native Hawaiian or Other Pacific Islander (1). Race was not reported for 8 patients.

Clinical trial exposure to pembrolizumab by race for all RCC patients in KN581 who received pembrolizumab combination therapy is summarized in Table SIII.100. The majority of the patients treated have been White (260) followed by Asian (81); Other race (4); and Black or African American (2). Five (5) patients did not report race.

Clinical trial exposure to pembrolizumab monotherapy by race for all RCC patients in KN564 is summarized in Table SIII.101. The majority of patients treated have been White (365) followed by Asian (63); American Indian or Alaska Native (10); Multiracial (8); and Black or African American (6). Thirty-six (36) patients did not report race.

Clinical trial exposure to pembrolizumab monotherapy by race for all CRC patients in KN177 is summarized in Table SIII.102. The majority of patients treated have been White (113), followed by Asian (24), and Black or African American (9). Seven (7) patients did not report race.

Clinical trial exposure to pembrolizumab by race for all esophageal cancer patients in KN590 who received pembrolizumab combination therapy is summarized in Table SIII.103. The majority of patients treated have been Asian (201), followed by White (136) American Indian or Alaska Native (9), Black or African American (5), and Multiracial (5). Fourteen (14) patients did not report race.

Clinical trial exposure to pembrolizumab by race for all TNBC patients in KN355 who received pembrolizumab combination therapy is summarized in Table SIII.104. The majority of patients treated have been White (399), followed by Asian (138), Black or African American (21), Multiracial (11), and American Indian or Alaska Native (10). Seventeen (17) patients did not report race.

Clinical trial exposure to pembrolizumab by race for all TNBC subjects in KN522 who received pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy is summarized in Table SIII.105. The majority of patients treated have been White (500), followed by Asian (148), Black or African American (38), American Indian or Alaska Native (14), Multiracial (13), Native Hawaiian or Other Pacific Islander (1). Sixty-four (64) patients did not report race.

Clinical trial exposure to pembrolizumab by race for all endometrial carcinoma patients in KN146 and KN775 who received pembrolizumab combination therapy is summarized in Table SIII.106. The majority of patients treated have been White (364), followed by Asian (90), Black or African American (24); Multiracial (7), American Indian or Alaska Native (5), Native Hawaiian or Other Pacific Islander (2), and Other race (2). Thirty-six (36) patients did not report race.

Clinical trial exposure to pembrolizumab by race for all endometrial carcinoma patients in KN868 who received pembrolizumab combination therapy is summarized in Table SIII.107. The majority of patients treated have been White (288), followed by Black or African American (53), Asian (17), Multiracial (1), American Indian Or Alaska Native (2), and Native Hawaiian Or Other Pacific Islander (1). Twenty (20) patients did not report race.

Clinical trial exposure to pembrolizumab monotherapy by race for all MSI-H patients in KN158 Cohort K and KN164 Cohorts A and B is summarized in Table SIII.108. The majority of patients treated have been White (368), followed by Asian (64), American Indian or Alaska Native (18), Black or African American (16) and Multiracial (7). Two (2) patients did not report race.

Clinical trial exposure to pembrolizumab by race for all cervical cancer patients in KNA18 and KN826 who received pembrolizumab combination therapy is summarized in Table SIII.109. The majority of patients treated have been White (423), followed by Asian (219), Multiracial (110), American Indian or Alaska Native (42), Black or African American (18), Native Hawaiian or Other Pacific Islander (2). Twenty-one (21) patients did not report race.

Clinical trial exposure to pembrolizumab by race for gastric cancer patients in KN811 who received pembrolizumab combination therapy is summarized in Table SIII.110. The majority of patients treated have been White (217), followed by Asian (119), Multiple (6), American Indian or Alaska Native (5), and Black or African American (2). Race was not reported for one (1) patient.

Clinical trial exposure to pembrolizumab by race for gastric cancer patients in KN859 who received pembrolizumab combination therapy is summarized in Table SIII.111. The majority of patients treated have been White (422), followed by Asian (269), Multiracial (43), American Indian or Alaska Native (31), Black or African American (12) and Native Hawaiian or Other Pacific Islander (1). Race was not reported for seven (7) patients.

Clinical trial exposure to pembrolizumab by race for all BTC patients in KN966 who received pembrolizumab combination therapy is summarized in Table SIII.112. The majority of patients treated have been White (254), followed by Asian (243), Black or African American (11),

Multiracial (5), American Indian or Alaska Native (2) and Native Hawaiian or Other Pacific Islander (1). Thirteen (13) patients did not report race.

Clinical trial exposure to pembrolizumab by race for all mesothelioma patients in KN483 and KNA17 who received pembrolizumab combination therapy is summarized in Table SIII.113. The majority of patients treated have been White (193), followed by Asian (20). Forty-seven (47) patients did not report race.

Table SIII.86: Clinical Trial Exposure by Race Melanoma Subjects KN001, KN002 and KN006 Monotherapy (APaT Population)

Race	Subjects	Mean Duration (days)	Subject Time (years)
American Indian Or Alaska Native	1	267.0	0.7
Asian	19	133.8	7.0
Black Or African American	7	171.0	3.3
Multiple	4	160.5	1.8
Multiracial	2	330.0	1.8
Native Hawaiian Or Other Pacific Islander	1	274.0	0.8
White	1530	229.3	960.4
Null	3	182.3	1.5

Duration of Exposure is calculated as last dose date - first dose date +1.
 (pembrolizumab KN001 Database Cutoff Date: 18APR2014).
 (pembrolizumab KN002 Database Cutoff Date: 28FEB2015).
 (pembrolizumab KN006 Database Cutoff Date: 03MAR2015).

Data on File

Table SIII.87: Clinical Trial Exposure by Race Melanoma Participants KN054 and KN716 Monotherapy (APaT Population)

Race	Participants	Mean Duration (days)	Participant Time (years)
American Indian Or Alaska Native	1	112.0	0.3
Asian	4	132.2	1.4
Black Or African American	3	230.7	1.9
Multiracial	1	255.0	0.7
White	432	251.2	297.1
NULL	551	281.0	423.9
Total	992	267.1	725.4

Duration of Exposure is calculated as last dose date - first dose date + 1.
 Race data were not collected for KN054.
 Database cutoff date for Melanoma (KN054: 02OCT2017, KN716: 04DEC2020)

Data on File

Table SIII.88: Clinical Trial Exposure by Race NSCLC Subjects KN001, KN010, KN024 and KN042 Monotherapy (APaT Population)

Race	Subjects	Mean Duration (days)	Subject Time (years)
American Indian Or Alaska Native	16	206.6	9.1
Asian	428	197.8	231.8
Black Or African American	53	187.4	27.2
Multiracial	35	292.0	28.0
Native Hawaiian Or Other Pacific Islander	3	57.3	0.5
White	1466	204.1	819.2
NULL	21	117.4	6.8
Total	2022	202.8	1122.5

Duration of Exposure is calculated as last dose date - first dose date +1.
 (pembrolizumab KN001 Database Cutoff Date for Lung: 23JAN2015).
 (pembrolizumab KN010 Database Cutoff Date: 30SEP2015).
 (pembrolizumab KN024 Database Cutoff Date: 10JUL2017).
 (pembrolizumab KN042 Database Cutoff Date: 26FEB2018).

Data on File

Table SIII.89: Clinical Trial Exposure to Pembrolizumab by Race NSCLC Subjects KN021 (Cohorts A, C and G), KN189 and KN407 Combination Therapy (APaT Population)

Race	Subjects	Mean Duration (days)	Subject Time (years)
American Indian Or Alaska Native	1	643.0	1.8
Asian	73	159.4	31.9
Black Or African American	30	224.6	18.4
Native Hawaiian Or Other Pacific Islander	1	512.0	1.4
White	681	235.5	439.1
NULL	5	163.0	2.2
Total	791	228.5	494.8

Duration of Exposure is calculated as last dose date - first dose date +1.
 MK-3475 Database Cutoff Date for Lung Combination Therapy (KN021 Cohort A: 07NOV2016, Cohort G/C: 31MAY2017, KN189: 8NOV2017; KN407: 03APR2018)

Data on File

Table SIII.90: Clinical Trial Exposure to Pembrolizumab by Race NSCLC Participants KN091 Monotherapy Therapy (APaT Population)

Race	Patients	Mean Duration (days)	Patient Time (years)
American Indian Or Alaska Native	1	365.0	1.0
Asian	107	292.6	85.7
Multiracial	4	335.7	3.7
Other	5	313.8	4.3
White	442	255.3	309.0
NULL	21	261.9	15.1
Total	580	263.7	418.7

Duration of Exposure is calculated as last dose date - first dose date + 1.
 KN091 Database Cutoff Date: 20SEP2021

Data on File

Table SIII.91: Clinical Trial Exposure to Pembrolizumab by Race NSCLC Participants in KN671 Combination Therapy (APaT Population)

Race	Patients	Mean Duration (days)	Patient Time (years)
American Indian Or Alaska Native	1	413.0	1.1
Asian	124	290.5	98.6
Black Or African American	6	268.5	4.4
Multiracial	3	340.0	2.8
White	249	279.4	190.5
NULL	13	281.0	10.0
Total	396	283.5	307.4

Duration of Exposure is calculated as last dose date - first dose date + 1.
Database Cutoff Date: 29JUL2022

Data on File

Table SIII.92: Clinical Trial Exposure to Pembrolizumab SC by Race NSCLC Participants in MK-3475A-D77 Combination Therapy (APaT Population)

Race	Participants	Mean Duration (days)	Participant Time (years)
American Indian Or Alaska Native	2	189.0	1.0
Asian	74	204.6	41.5
Black Or African American	5	155.0	2.1
Multiracial	12	217.4	7.1
White	158	188.7	81.6
Total	251	194.1	133.4

Duration of Exposure (days) is calculated as last dose date - first dose date +1 for pembrolizumab.
Database Cutoff Date for Lung (MK-3475A-D77: 12JUL2024).

Data on File

Table SIII.93: Clinical Trial Exposure by Race HL Subjects KN013, KN087, and KN204 Monotherapy (APaT Population)

Race	Subjects	Mean Duration (days)	Subject Time (years)
American Indian Or Alaska Native	2	596.0	3.3
Asian	25	416.9	28.5
Black Or African American	11	469.0	14.1
Multiracial	7	328.7	6.3
Native Hawaiian Or Other Pacific Islander	1	715.0	2.0
White	330	377.8	341.4
NULL	13	289.5	10.3
Total	389	381.1	405.8

Duration of Exposure is calculated as last dose date - first dose date + 1.
Database Cutoff Date (KN013-Cohort 3: 28SEP2018, KN087: 21MAR2019, KN204: 16JAN2020)

Data on File

Table SIII.94: Clinical Trial Exposure by Race UC Subjects KN045 and KN052 Monotherapy (APaT Population)

Race	Subjects	Mean Duration (days)	Subject Time (years)
American Indian Or Alaska Native	2	65.0	0.4
Asian	90	135.8	33.5
Black Or African American	13	73.2	2.6
Multiracial	3	160.0	1.3
White	512	137.1	192.2
NULL	16	142.8	6.3
Total	636	135.6	236.2

Duration of Exposure is calculated as last dose date - first dose date +1.
Includes all subjects with urothelial carcinoma who received at least one dose of Pembrolizumab in KN045 and KN052.
Pembrolizumab KN045 Database Cutoff Date: 07SEP2016.
Pembrolizumab KN052 Database Cutoff Date: 01SEP2016.

Data on File

Table SIII.95: Clinical Trial Exposure by Race UC Participants KNA39 and KN869 Pembrolizumab + EV Combination Therapy (APaT Population)

Race	Participants	Mean Duration (months)	Participant Time (years)
American Indian or Alaska Native	2	11.15	1.86
Asian	107	10.27	91.58
Black Or African American	9	9.71	7.28
White	410	8.70	297.42
Other	5	6.26	2.61
Missing/Unknown	31	8.82	22.78
Total	564	9.01	423.53

Duration of exposure is calculated as last dose date - first dose date + 1 for pembrolizumab in the combination.
Pembrolizumab KN-869 Database Cutoff Date: 13Mar2023, KN-A39 Database Cutoff Date: 08Aug2023

Data on File

Table SIII.96: Clinical Trial Exposure by Race HNSCC Subjects KN012 B+B2 Cohorts, KN040, KN055 and KN048 Monotherapy (ASaT Population)

Race	Subjects	Mean Duration (days)	Subject Time (years)
American Indian Or Alaska Native	9	185.2	4.6
Asian	109	165.5	49.4
Black Or African American	28	178.0	13.6
Multi-Racial	12	179.3	5.9
Multiracial	6	154.8	2.5
White	723	162.6	321.8
NULL	22	144.4	8.7
Total	909	163.3	406.5

Duration of Exposure is calculated as last dose date - first dose date +1.
MK-3475 Database Cutoff Date for HNSCC (KN012-HNSCC: 26APR2016, KN040: 15MAY2017, KN048: 13JUN2018, KN055: 22APR2016)

Data on File

Table SIII.97: Clinical Trial Exposure to Pembrolizumab by Race HNSCC Subjects in KN048 Combination Therapy (ASaT Population)

Race	Subjects	Mean Duration (days)	Subject Time (years)
American Indian Or Alaska Native	3	244.7	2.0
Asian	60	161.6	26.5
Black Or African American	10	227.5	6.2
Multiracial	3	249.3	2.0
White	200	240.7	131.8
Total	276	223.2	168.6

Duration of Exposure is calculated as last dose date - first dose date +1.
MK-3475 Database Cutoff Date for KN048: 13JUN2018.

Data on File

Table SIII.98: Clinical Trial Exposure to Pembrolizumab by Race HNSCC Participants in KN689 Monotherapy or Combination Therapy (APaT Population)

Race	Participants	Mean Duration (days)	Participant Time (years)
Asian	50	211.1	28.9
Black Or African American	8	198.1	4.3
Multiracial	9	242.9	6.0
Native Hawaiian Or Other Pacific Islander	1	104.0	0.3
White	284	236.7	184.1
NULL	9	187.9	4.6
Total	361	230.9	228.2

Duration of Exposure (days) is calculated as last dose date - first dose date +1 for pembrolizumab.
Database cutoff date for KN689: 25JUL2024

Table SIII.99: Clinical Trial Exposure to Pembrolizumab by Race RCC Subjects KN426 Combination Therapy (ASaT Population)

Race	Subjects	Mean Duration (days)	Subject Time (years)
American Indian Or Alaska Native	2	372.5	2.0
Asian	66	273.6	49.4
Black Or African American	9	222.4	5.5
Missing	8	370.0	8.1
Multiracial	1	358.0	1.0
Native Hawaiian Or Other Pacific Islander	1	347.0	1.0
White	342	288.5	270.1
Total	429	287.0	337.1

Duration of Exposure is calculated as last dose date - first dose date +1.for pembrolizumab in the combo.
MK-3475 Database Cutoff Date for KN426: 24AUG2018.

Data on File

Table SIII.100: Clinical Trial Exposure to Pembrolizumab by Race RCC Participants KN581 Combination Therapy (APaT)

Race	Participants	Mean Duration (days)	Participant Time (years)
White	260	445.5	317.1
Black Or African American	2	713.5	3.9
Asian	81	426.2	94.5
American Indian Or Alaska Native	0	---	---
Native Hawaiian Or Other Pacific Islander	0	---	---
Other	4	214.0	2.3
Missing	5	435.4	6.0
Total	352	439.8	423.9

Duration of Exposure is calculated as last dose date - first dose date + 1 for pembrolizumab in the combo.MK-3475 Database Cutoff Date for KN581: 28AUG2020.

Data on File

Table SIII.101: Clinical Trial Exposure by Race RCC Participants KN564 Monotherapy (APaT Population)

Race	Participants	Mean Duration (days)	Person Time (years)
American Indian Or Alaska Native	10	268.6	7.4
Asian	63	283.4	48.9
Black Or African American	6	279.8	4.6
Multiracial	8	315.3	6.9
White	365	268.7	268.5
Missing	36	283.9	28.0
Total	488	272.6	364.3

Duration of Exposure is calculated as last dose date - first dose date + 1.
Database cutoff date for RCC (KN564: 14DEC2020)

Data on File

Table SIII.102: Clinical Trial Exposure by Race CRC Subjects KN177 Monotherapy (ASaT Population)

Race	Subjects	Mean Duration (days)	Subject Time (years)
Asian	24	427.5	28.1
Black Or African American	9	303.6	7.5
White	113	406.6	125.8
Missing	7	398.0	7.6
Total	153	403.4	169.0

Duration of Exposure is calculated as last dose date - first dose date +1.
Database cutoff date for CRC (KN177: 19FEB2020)

Data on File

Table SIII.103: Clinical Trial Exposure to Pembrolizumab by Race Esophageal Cancer Subjects KN590 Combination Therapy (ASaT Population)

Race	Subjects	Mean Duration (days)	Subject Time (years)
American Indian Or Alaska Native	9	146.4	3.6
Asian	201	223.3	122.9
Black Or African American	5	258.4	3.5
Multiracial	5	217.0	3.0
White	136	246.0	91.6
Missing	14	346.6	13.3
Total	370	234.8	237.9

Duration of Exposure is calculated as last dose date - first dose date +1.
Database Cutoff date for Esophageal (KN590: 02JUL2020)

Data on File

Table SIII.104: Clinical Trial Exposure to Pembrolizumab by Race TNBC Subjects in KN355 Combination Therapy (ASaT Population)

Race	Subjects	Mean Duration (days)	Subject Time (years)
American Indian Or Alaska Native	10	203.6	5.6
Asian	138	283.2	107.0
Black Or African American	21	237.2	13.6
Multiracial	11	244.3	7.4
White	399	241.1	263.4
Missing	17	180.3	8.4
Total	596	248.4	405.3

Duration of Exposure is calculated as last dose date - first dose date +1 for pembrolizumab in the combo.
Database Cutoff Date for KN355: 11DEC2019.

Data on File

Table SIII.105: Clinical Trial Exposure to Pembrolizumab by Race TNBC Subjects in KN522 Combination Therapy (APaT Population)

Race	Participants	Mean Duration (days)	Person Time (years)
American Indian Or Alaska Native	14	328.8	12.6
Asian	148	358.4	145.2
Black Or African American	38	325.3	33.8
Multiracial	13	343.2	12.2
Native Hawaiian Or Other Pacific Islander	1	397.0	1.1
White	500	325.3	445.3
Missing	64	339.2	59.4
Total	778	333.2	709.7

Duration of exposure is the time from the first dose date to the last dose date for pembrolizumab.
Database cutoff date for TNBC: (KN522: 23MAR2021)

Data on File

Table SIII.106: Clinical Trial Exposure to Pembrolizumab by Race Endometrial Carcinoma Participants KN146 and KN775 Combination Therapy (APaT Population)

Race	Participants	Mean Duration (days)	Participants Time (years)
American Indian Or Alaska Native	5	329.0	4.5
Asian	90	256.1	63.1
Black Or African American	24	236.0	15.5
Multiracial	7	196.7	3.8
Native Hawaiian Or Other Pacific Islander	2	11.0	0.1
Other	2	69.5	0.4
White	364	265.8	264.9
Missing	36	279.8	27.6
Total	530	261.7	379.8

Duration of exposure (day) is calculated as last dose date - first dose date + 1.
 Database cutoff date for endometrial carcinoma (KN146: 18AUG2020, KN775: 26OCT2020)

Data on File

Table SIII.107: Clinical Trial Exposure to Pembrolizumab by Race Endometrial Carcinoma Participants in KN868 Combination Therapy (APaT)

Race	Participants	Mean Duration (days)	Participants Time (years)
American Indian Or Alaska Native	2	415.5	2.3
Asian	17	304.4	14.2
Black Or African American	53	196.4	28.5
Multiracial	1	1.0	0.0
Native Hawaiian Or Other Pacific Islander	1	380.0	1.0
White	288	211.1	166.4
Missing	20	175.6	9.6
Total	382	212.3	222.0

Duration of Exposure is calculated as last dose date - first dose date + 1 for pembrolizumab.
 Database cutoff date (KN868: 16DEC2022 for dMMR participants and 06DEC2022 for pMMR participants)

Data on File

Table SIII.108: Clinical Trial Exposure by Race MSI-H Subjects in KN158 Cohort K and KN164 Cohorts A and B Monotherapy (ASaT Population)

Race	Subjects	Mean Duration (days)	Subject Time (years)
American Indian Or Alaska Native	18	286.8	14.1
Asian	64	315.0	55.2
Black Or African American	16	432.1	18.9
Multiracial	7	152.0	2.9
White	368	288.3	290.5
NULL	2	465.5	2.5
Total	475	295.4	384.2

Duration of Exposure is calculated as last dose date - first dose date + 1.
 Database cutoff date for MSI-H (KN158-cohort K: 05OCT2020)
 Database cutoff date for Colorectal (KN164-cohorts A and B: 09SEP2019)

Data on File

Table SIII.109: Clinical Trial Exposure to Pembrolizumab by Race Cervical Cancer Participants in KNA18 + KN826 Combination Therapy (APaT Population)

Race	Participants	Mean Duration (days)	Participant Time (years)
American Indian Or Alaska Native	42	348.2	40.0
Asian	219	431.4	258.7
Black or African American	18	300.8	14.8
Multiracial	110	367.7	110.7
Native Hawaiian Or Other Pacific Islander	2	283.5	1.6
White	423	327.9	379.8
NULL	21	290.9	16.7
Total	835	359.7	822.3

Duration of exposure is calculated as last dose date - first dose date +1 for pembrolizumab in the combo
 Database cutoff date for Cervical Cancer (KNA18: 09JAN2023, KN826: 03MAY2021).

Data on File

Table SIII.110: Clinical Trial Exposure to Pembrolizumab by Race in Gastric Cancer Participants KN811 Combination Therapy (APaT Population)

Race	Patients	Mean Duration (days)	Patient Time (years)
American Indian Or Alaska Native	5	282.6	3.9
Asian	119	336.8	109.7
Black Or African American	2	478.0	2.6
Multiple	6	342.3	5.6
White	217	323.7	192.3
NULL	1	304.0	0.8
Total	350	328.7	315.0

Duration of Exposure is calculated as last dose date - first dose date + 1.
 Database Cutoff Date: 25MAY2022

Data on File

Table SIII.111: Clinical Trial Exposure to Pembrolizumab by Race in Gastric Cancer Participants KN859 Combination Therapy (APaT Population)

Race	Participants	Mean Duration (days)	Participant Time (years)
American Indian Or Alaska Native	31	220.2	18.7
Asian	269	272.3	200.6
Black Or African American	12	235.9	7.8
Multiracial	43	314.0	37.0
Native Hawaiian Or Other Pacific Islander	1	28.0	0.1
White	422	249.1	287.8
NULL	7	197.0	3.8
Total	785	258.5	555.6

Each participant is counted once on each applicable duration category row.
Duration of exposure is calculated as last dose date - first dose date + 1.
Database cutoff date for Gastric (KN859: 03OCT2022)

Data on File

Table SIII.112: Clinical Trial Exposure to Pembrolizumab by Race BTC Participants in KN966 Combination Therapy (APaT Population)

Race	Participants	Mean Duration (days)	Participant Time (years)
American Indian Or Alaska Native	2	417.0	2.3
Asian	243	229.3	152.5
Black Or African American	11	175.9	5.3
Multiracial	5	122.6	1.7
Native Hawaiian Or Other Pacific Islander	1	279.0	0.8
White	254	238.2	165.7
Missing	13	294.5	10.5
Total	529	233.9	338.7

Duration of exposure is calculated as last dose date - first dose date + 1 for pembrolizumab in the combo.
(pembrolizumab KN966 Database Cutoff Date: 15DEC2022)

Data on File

Table SIII.113: Clinical Trial Exposure to Pembrolizumab by Race Mesothelioma Participants in KN483 and KNA17 Combination Therapy (APaT Population)

Race	Subjects	Mean Duration (days)	Participant Time (years)
Asian	20	323.8	17.7
White	193	272.3	143.9
NULL	47	295.7	38.1
Total	260	280.5	199.7

Duration of Exposure is calculated as last dose date - first dose date +1.
Database Cutoff Date for Thoracic Malignancy (KN483: 16SEP2022, KNA17: 21SEP2022).;

Data on File

PART II: MODULE SIV - POPULATIONS NOT STUDIED IN CLINICAL TRIALS

SIV.1 Exclusion Criteria in Pivotal Clinical Studies Within the Development Program

Table SIV.1.1: Exclusion Criteria in Pivotal Clinical Studies Within the Development Program

Exclusion Criterion	Reason for Exclusion	Is it Considered to be Missing Information?	Rationale (if not Included as Missing Information)
Hypersensitivity to the active substance or to any of the excipients	Avoid potential harm to the patient	No	Pembrolizumab should not be administered to persons with known hypersensitivity to the active substance or to any of the excipients.
Severe renal or moderate to severe hepatic impairment	No clinical trials have been conducted in subjects with severe renal or moderate or severe hepatic comorbidities	No	Adequately reflected in the SmPC (sections 4.2, 4.4 and 5.2)
Diagnosis of immunosuppression or is receiving systemic steroid therapy or any other form of immunosuppressive therapy within 7 days	May confound understanding of drug efficacy or safety profile. Use of immunosuppressive agents may negatively affect efficacy	No	Adequately reflected in the SmPC (sections 4.4, 4.5)
Active infection requiring therapy	Avoid factors that may confound understanding of drug safety profile needed for subsequent drug development and use.	No	Patients that have an active infection should have the infection treated prior to starting.
Autoimmune disease requiring systemic treatment in past 2 years. Replacement therapy allowed.	Avoid factors that may confound understanding of drug efficacy or safety profile needed for subsequent drug development and use.	No	Adequately reflected in the SmPC (section 4.4)
Known history of Human Immunodeficiency Virus (HIV) (HIV 1/2 antibodies). Patient is positive for active Hepatitis B (HBsAg reactive) or Hepatitis C (HCV RNA) [qualitative] is detected). Patients with negative Hepatitis C antibody testing may not require RNA testing if deemed appropriate by treating physician	The impact of this agent on HIV or Hepatitis B or C remains unknown. These subjects were excluded given the theoretical safety concern. Hepatitis is an identified adverse event of anti-PD-1 antibody treatment	No	Adequately reflected in the SmPC (section 4.4)

Table SIV.1.1: Exclusion Criteria in Pivotal Clinical Studies Within the Development Program

Exclusion Criterion	Reason for Exclusion	Is it Considered to be Missing Information?	Rationale (if not Included as Missing Information)
History of non-infectious pneumonitis that required steroids	Avoid increasing risk of pneumonitis from pembrolizumab	No	Pneumonitis is a known risk and is included as an important identified risk for pembrolizumab. Patients with a history of pneumonitis may experience pneumonitis during treatment. However, given the life-threatening nature of the indication and the ability to reverse pneumonitis should it occur during treatment with pembrolizumab, these patients should still have the option of treatment.
Known psychiatric or substance abuse disorders that would interfere with cooperation with the requirements of the trial.	Avoid factors that may confound understanding of drug efficacy or safety profile needed for subsequent drug development and use.	No	Exclusion is specific to requirements of clinical trial participation rather than real world use.
Prior treatment with IPI or other anti-CTLA-4 agent, any anti-PD-1, anti-PD-L1, or anti-PD-L2 agent.	Avoid factors that may confound understanding drug efficacy and safety	No	Exclusion is specific to understanding of efficacy and safety in clinical trial environment rather than real world use.
Being pregnant or breastfeeding, or expecting to conceive or father children within the projected duration of the study, starting with the Pre-screening or Screening visit through 120 days after the last dose of study medication.	Avoid potential harm to the unborn fetus or breastfeeding newborn.	No	Adequately reflected in the SmPC (section 4.6)
Clinically active CNS involvement	High risk patients where active CNS involvement portends poor prognosis and symptoms of the disease. To ensure patients are receiving optimal treatment, per their physician's discretion for their co-morbid conditions which may not always be possible within the confines of a clinical trial.	No	Patients with treated brain metastases were shown to have similar safety and activity of pembrolizumab. Given the fatal nature of their cancer, physicians should be allowed to use discretion on the management of brain metastases with systemic and local treatment options.

Table SIV.1.1: Exclusion Criteria in Pivotal Clinical Studies Within the Development Program

Exclusion Criterion	Reason for Exclusion	Is it Considered to be Missing Information?	Rationale (if not Included as Missing Information)
Having had chemotherapy, radioactive, or biological cancer therapy within 2 weeks prior to the first dose of study therapy	Avoid factors that may confound understanding of drug safety profile and efficacy needed for subsequent drug development and use. To ensure patients are receiving optimal treatment, per their physician’s discretion for their co-morbid conditions which may not always be possible within the confines of a clinical trial.	No	Experience to date demonstrates that pembrolizumab is well tolerated for oncology patients. Substantial documented benefits of treatment in life threatening, or otherwise lethal disease may outweigh risks.
Currently participating or participated in a study of an investigational agent or using an investigational device within 30 days of administration of pembrolizumab	Avoid factors that may confound understanding of the efficacy and safety of pembrolizumab in a clinical trial	No	This will not be applicable in the post-marketing environment as the appropriate population will be defined in the SmPC.
Known additional malignancy that is progressing or requires active treatment	To ensure patients are receiving optimal treatment, per their physician’s discretion for their co-morbid conditions which may not always be possible within the confines of a clinical trial.	No	Physicians should be allowed to determine which tumor possess the greatest risk and is the highest priority for treatment.
Received a live vaccine within 30 days prior to first dose	Avoid factors that may confound understanding of the efficacy of pembrolizumab in a clinical trial	No	Exclusion is specific to understanding of efficacy and safety in clinical trial environment.
History or current evidence of any condition, therapy, or laboratory abnormality that might confound the results of the study, interfere with the patient’s participation for the full duration of the study, or is not in the best interest of the patient to participate, in the opinion of the treating investigator	Avoid factors that may confound understanding of drug efficacy or safety profile needed for subsequent drug development and use.	No	Information on the conditions are known risks and may be included in other sections of the SmPC.

SIV.2 Limitations to Detect Adverse Reactions in Clinical Trial Development Program

The clinical development programme is unlikely to detect certain types of adverse reactions such as rare adverse reactions, adverse reactions with a long latency, or those caused by prolonged or cumulative exposure.

SIV.3 Limitations in Respect to Populations Typically Under-represented in Clinical Trial Development Program

Those populations detailed in Section SIV will require additional investigation during the development of pembrolizumab. Areas of missing information remain and will be assessed through the clinical development program and post marketing safety surveillance. The European Medicines Agency (EMA) Decision for the first paediatric investigation plan (PIP) covering the condition ‘Treatment of all conditions included in the category of malignant neoplasms (except nervous system, haematopoietic and lymphoid tissue)’ has been adopted on March 7, 2014. The first PIP was subsequently modified and the EMA decision for the modifications was adopted on February 16, 2018. The EMA Decision for the second PIP covering the condition ‘Treatment of Hodgkin Lymphoma’ has been adopted on August 1, 2016. The second PIP was subsequently modified and the EMA decision for the modification was adopted on January 30, 2018, and December 6, 2021, respectively. The marketing authorisation holder (MAH) will follow the PIP plans that have been adopted.

The first clinical study in paediatric patients, KEYNOTE-051, began enrollment in March 2015 and a clinical study report was prepared and submitted to the Agency in February 2019, resulting in adoption of an opinion for a PI update on June 27, 2019.

With RMP version 30.0, an updated clinical study report for KEYNOTE-051, with an additional 16 months of follow-up, was included to support the indication in relapsed or refractory classical HL in the paediatric population.

Table SIV.3.1: Exposure of Special Populations Included or not in Clinical Trial Development Programs

Type of Special Population	Exposure
Pregnant and breastfeeding women	<p>Pregnant and breastfeeding women were not included in the pre-authorisation clinical development program.</p> <p>There have been no trials evaluating pembrolizumab in pregnant or lactating women. In clinical trials, women of childbearing potential have been advised to avoid pregnancy and to use effective contraception during the dosing period and for a period of at least 120 days thereafter. A pregnancy test was conducted on each pre-menopausal female of childbearing potential within 72 hours prior to administration of pembrolizumab. If pregnancy was confirmed during the course of the trial, study drug administration was discontinued. Any subject who became pregnant while in a study was followed to the completion/termination of the pregnancy. If the pregnancy continued to term, the outcome (health of infant) was to be reported to the MAH.</p>
<p>Patients with relevant comorbidities:</p> <ul style="list-style-type: none"> • Patients with hepatic impairment • Patients with renal impairment • Patients with cardiovascular impairment • Immunocompromised patients • Patients with a disease severity different from inclusion criteria in clinical trials 	<p>Patients with the following conditions were excluded from clinical trials: active CNS metastases; ECOG PS ≥ 2 (except for urothelial carcinoma and RCC); HIV, hepatitis B or hepatitis C infection; active systemic autoimmune disease; interstitial lung disease; prior pneumonitis requiring systemic corticosteroid therapy; a history of severe hypersensitivity to another monoclonal antibody; receiving immunosuppressive therapy and a history of severe immune-related adverse reactions from treatment with ipilimumab, defined as any Grade 4 toxicity or Grade 3 toxicity requiring corticosteroid treatment (> 10 mg/day prednisone or equivalent) for greater than 12 weeks. Patients with active infections were excluded from clinical trials and were required to have their infection treated prior to receiving pembrolizumab. Patients with active infections occurring during treatment with pembrolizumab were managed with appropriate medical therapy. Patients with clinically significant renal (creatinine > 1.5 x Upper Limit of Normal (ULN)) or hepatic (bilirubin > 1.5 x ULN, alanine aminotransferase (ALT), aspartate aminotransferase (AST) > 2.5 x ULN in the absence of liver metastases) abnormalities at baseline were excluded from clinical trials, therefore information is limited in patients with severe renal and moderate to severe hepatic impairment.</p> <p>These exclusion criteria were applied to ensure that the population under study had adequate organ function in order to have a homogeneous patient population and to avoid potential confounding variables in the interpretation of safety data from the studies.</p>
Population with relevant different ethnic origin	Clinical trials (included in this RMP) exposure by race is presented in PART II: MODULE SIII.
Subpopulations carrying relevant genetic polymorphisms	Subpopulations carrying relevant genetic polymorphisms were not included in the pre-authorisation clinical development program.

PART II: MODULE SV - POST-AUTHORISATION EXPERIENCE

SV.1 Post-Authorisation Exposure

SV.1.1 Method Used to Calculate Exposure

A summary of the worldwide distribution of pembrolizumab for the cumulative period from market introduction to 03-SEP-2024 is presented below based on the available data.

Patient exposure estimates were calculated from our Company's internal distribution data from the Worldwide Financial Reporting System (WFRS), and the Financial Sharing Area databases. This data provides a more complete and consistent methodology for the estimate of patient exposure worldwide for current Company products. Patient exposure estimates were calculated from expanded distribution categories to provide a more accurate estimate of patient exposure worldwide. The effects of this update (starting 01-AUG-2018) may be apparent when comparing current estimates of patient exposure to those of prior reporting periods.

Pembrolizumab IV is available in 20 mg, 50 mg, and 100 mg vials. The dosing recommendation in adults is either 200 mg every 3 weeks or 400 mg every 6 weeks. Since patient exposure for the doses of 200 mg every 3 weeks or 400 mg every 6 weeks are essentially equivalent, standard dosing of 200 mg every 3 weeks was used for patient exposure estimation.

- The total number of vials for 20 mg vial, 50 mg vial, and 100 mg vial was divided by the average number of vials required to formulate a standard dose of 200mg. Specifically, the number of units utilized when dosing with 20 mg vials was on average 10 units; 50 mg vials was on average 4 units; and 100 mg vials was on average 2 units. Therefore, the total number of vials for the 20 mg strength was divided by 10, the total number of vials for the 50 mg strength was divided by 4, and the total number of vials for the 100 mg strength was divided by 2, to determine the number of doses (cycles).
- The estimated number of doses (cycles) per vial was then divided by the average number of doses estimated per patient per year of treatment with pembrolizumab. The average number of doses estimated per year of treatment with approximately 23 days between doses (recommended 21 days with 2-day allowance for weekends), or an average of 15.9 doses per patient regardless of strength (365 days/23 days).

SV.1.2 Exposure

The estimated number of doses of pembrolizumab distributed worldwide from product launch through 03-SEP-2024 is 49,260,870. This corresponds to 1,518,204 patient years of treatment.

PART II: MODULE SVI - ADDITIONAL EU REQUIREMENTS FOR THE SAFETY SPECIFICATION

Potential for Misuse for Illegal Purposes

Pembrolizumab is a potent and highly selective humanized mAb of the IgG4/kappa isotype designed to directly block the interaction between PD-1 and its ligands, PD-L1 and PD-L2. Neither pembrolizumab nor its components are known to possess addictive properties.

PART II: MODULE SVII - IDENTIFIED AND POTENTIAL RISKS

SVII.1 Identification of Safety Concerns in the Initial RMP Submission

Not applicable.

SVII.1.1 Risks Not Considered Important for Inclusion in the List of Safety Concerns in the RMP

Not applicable.

SVII.1.2 Risks Considered Important for Inclusion in the List of Safety Concerns in the RMP

Not applicable.

SVII.2 New Safety Concerns and Reclassification With a Submission of an Updated RMP

Not applicable.

SVII.3 Details of Important Identified Risks, Important Potential Risks, and Missing Information

SVII.3.1 Presentation of Important Identified Risks and Important Potential Risks

The characterization of the risk profile of pembrolizumab is presented based on four data sets (n=2799), the melanoma and NSCLC patient populations in KN001, the melanoma patient population in KN002, the melanoma patient population in KN006, and the previously treated NSCLC patient population in KN010, and is considered as the reference safety dataset for pembrolizumab. The reference safety dataset serves as the locked dataset for the characterization of each risk associated with pembrolizumab and is presented in tabular format.

The Important Identified and Potential Risks apply to both intravenous and subcutaneous routes of administration. The studies listed in the RMP consist of trials which utilized pembrolizumab administered intravenously and subcutaneously.

Safety data related to the previously untreated NSCLC population in KN024 (database cutoff date 10-JUL-2017) and KN042 (database cutoff date 26-FEB-2018) (total n=790) was evaluated and is presented in text for each risk. Overall, the safety profile from the NSCLC population in KN024 and KN042 was generally similar to the observed safety profile from the reference safety dataset.

Safety data related to the NSCLC population in KN189 (database cutoff date 08-NOV-2017) and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (database cutoff date 31-MAY-2017) (total n=488) were evaluated and are presented in text for each risk. Overall, the safety profile from the NSCLC population in KN189 and KN021 (Cohorts C and G)

pembrolizumab plus chemotherapy was generally similar to the observed safety profile from the reference safety dataset.

Safety data related to the Hodgkin Lymphoma (HL) populations in KN013 (database cutoff date 28-SEP-2018), KN087 (database cutoff date 21-MAR-2019), and KN204 (database cutoff date 16-JAN-2020) (total n=389) were evaluated and are presented in text for each risk. Overall, the safety profile from the HL population was generally similar to the observed safety profile in the reference safety dataset, with the exception of a higher incidence of pneumonitis (8% versus 3.4%) and hypothyroidism (17% versus 8.5%). The higher incidence of pneumonitis was likely reflective of high rates of prior exposure to radiation and/or chemotherapy; agents with a known association of pulmonary toxicity. The majority of pneumonitis events were Grade 3 and below and resolved with systemic corticosteroids. There were no fatal events of pneumonitis. All reported events of hypothyroidism were Grade 1 and 2 and resolved with standard therapeutic measures. The observed incidence of hypothyroidism was considered attributable to a prior history of regional radiation and to a higher rate of pre-existing hypothyroidism among participants in this group at baseline.

Safety data related to the urothelial carcinoma (UC) indication in KN052 (database cutoff date 01-SEP-2016) (n=370) and KN045 (database cutoff date 07-SEP-2016) (n=266) were evaluated and are presented in text for each risk. Overall, the safety profile from the UC population in KN052 and in KN045 was generally similar to the observed safety profile in the reference safety dataset.

Safety data related to the UC population in KN869 and KNA39 who received pembrolizumab plus enfortumab vedotin (database cutoff date 13-Mar-2023 and 08-Aug-2023, respectively) (total n=564) were evaluated and are presented in text for each risk. Overall, the safety findings observed for pembrolizumab in combination with enfortumab vedotin in the UC population in KNA39 and KN869 was generally similar to the known safety profiles of pembrolizumab and enfortumab vedotin monotherapies. The incidence of rash maculo-papular with the combination was higher than with pembrolizumab monotherapy (36.0 % in KNA39 and KN869 vs 3.9% in the pembrolizumab monotherapy reference safety dataset (RSD), respectively), reflecting the contribution of enfortumab vedotin and longer duration of exposure.

Safety data related to the HNSCC population in KN040 (database cutoff date 15-MAY-2017), KN012 (database cutoff date 26-APR-2016), and KN055 (database cutoff date 22-APR-2016) (total n=609) were evaluated and are presented in text for each risk. Overall, the safety profile from the HNSCC population was generally similar to the reference safety dataset.

Safety data related to the HNSCC population in KN048 (database cutoff date 13-JUN-2018) (pembrolizumab monotherapy n=300; pembrolizumab plus chemotherapy n=276) were evaluated and are presented in text for each risk. Overall, the safety profile of pembrolizumab monotherapy from the HNSCC population in KN048 was generally similar to the reference safety dataset. The AE profile observed in pembrolizumab plus chemotherapy was generally similar to the known safety profiles of either pembrolizumab monotherapy or platinum/5-FU chemotherapy.

Safety data related to the melanoma indication in KN054 (database cutoff date 02-OCT-2017) (n= 509) was evaluated and is presented in text for each risk. Overall, the safety profile from the melanoma population in KN054 was generally similar to the observed safety profile from the reference safety dataset. Higher frequencies of some immune-mediated AEs, mostly mild to moderate in severity and manageable with treatment interruption or discontinuation and/or corticosteroid therapy were observed in KN054 in comparison to the reference safety dataset. This could be a reflection of the >2-fold longer exposure to pembrolizumab in KN054 subjects compared to subjects in the reference safety dataset.

Safety data related to the NSCLC population in KN021 Cohort A (database cutoff date 07-NOV-2016) and KN407 (database cutoff 03-APR-2018) (total n=303) were evaluated and are presented in text for each risk. Overall, the risk profile from the NSCLC population observed in the pooled KN407 and KN021 Cohort A combo was generally similar to the risk profile for pembrolizumab as characterized in the reference safety dataset, and the AE summary profile observed was generally similar to the known safety profiles of either pembrolizumab monotherapy or carboplatin/paclitaxel (or nab-paclitaxel) chemotherapy.

Safety data related to the RCC population in KN426 who received pembrolizumab plus axitinib (database cutoff date 24-AUG-2018), were evaluated and are presented in text for each risk. The AE profile observed in pembrolizumab plus axitinib was generally similar to the known safety profiles of the combined safety profiles of pembrolizumab monotherapy (both in the reference safety dataset and in 1L RCC) and axitinib monotherapy in 1L RCC, with the exception of a higher than expected incidence of Grade 3-4 transaminase elevations. In KN426, Grades 3 and 4 ALT increased (20%) and AST increased (13%) were observed in the pembrolizumab plus axitinib group. The median time to onset of ALT ≥ 3 times ULN was 2.3 months (range: 7 days to 19.8 months). In patients with ALT ≥ 3 times ULN (Grade 2-4, n=116), ALT resolved to Grade 0-1 in 94%. Sixty-one percent of the patients with increased ALT received systemic corticosteroids. Of the patients who recovered, 92 (84%) were rechallenged with either pembrolizumab (3%) or axitinib (31%) monotherapy or with both (50%). Of these patients, 55% had no recurrence of ALT >3 times ULN, and of those patients with recurrence of ALT >3 times ULN, all recovered. There were no Grade 5 hepatic events.

Safety data related to the CRC population in KN177 (database cutoff 19-FEB-2020) (n=153) were evaluated and are presented in text for each risk. Overall, the safety profile of pembrolizumab monotherapy in the CRC population in KN177 was generally similar to the observed safety profile in the reference safety dataset.

Safety data related to the esophageal cancer population in KN590 (database cutoff 02-JUL-2020) who received pembrolizumab plus chemotherapy (total n=370) were evaluated and are presented in text for each risk. The AE profile observed in pembrolizumab plus chemotherapy group was generally similar to the known safety profiles of either pembrolizumab monotherapy or platinum/5-FU chemotherapy.

Safety data related to the TNBC population in KN355 (database cutoff 11-DEC-2019) who received pembrolizumab plus chemotherapy (total n=596) were evaluated and are presented in text for each risk. Overall, the observed safety profile of pembrolizumab in combination with

chemotherapy in TNBC was generally similar to the known safety profiles of pembrolizumab monotherapy and the chemotherapy administered.

Safety data related to the RCC population in KN581 who received pembrolizumab plus lenvatinib (database cutoff 28-AUG-2020) (n=352) were evaluated and are presented in text for each risk. Overall, the safety profile observed in pembrolizumab in combination with lenvatinib in the RCC population in KN581 was generally similar to the known safety profiles of pembrolizumab monotherapy (reference safety dataset) and lenvatinib monotherapy, except for a higher incidence of immune-mediated AEs primarily due to hypothyroidism (47.2%), mostly grade 1 and 2 events. Smaller increases in some immune-mediated AEs were also observed. The nature (severity, management, and outcome) of the immune-mediated AEs observed with the pembrolizumab and lenvatinib combination was similar to the nature of immune-mediated AEs previously seen with pembrolizumab monotherapy.

Safety data related to the endometrial carcinoma population in KN146 (database cutoff 18-AUG-2020) and KN775 (database cutoff 26-OCT-2020) (total n=530) who received pembrolizumab plus lenvatinib, were evaluated and are presented in text for each risk. Overall, the safety profile observed in pembrolizumab in combination with lenvatinib in endometrial carcinoma was generally similar to the known safety profiles of pembrolizumab monotherapy (reference safety dataset) and lenvatinib monotherapy, except for a higher incidence of hypothyroidism (55.8%), mostly due to grade 1 and 2 events. Smaller increases in some immune-mediated AEs were observed, including colitis (4.9%) and hyperthyroidism (10.2%). The nature (severity, management, and outcome) of the immune-mediated AEs observed with the pembrolizumab and lenvatinib combination was similar to the nature of immune-mediated AEs previously seen with pembrolizumab monotherapy.

Safety data related to the RCC population in KN564 (database cutoff 14-DEC-2020) (n=488) were evaluated and are presented in text for each risk. Overall, the safety profile of pembrolizumab monotherapy in the RCC population in KN564 was generally similar to the observed safety profile in the reference safety dataset, except for a higher incidence of immune-mediated events primarily due to hypothyroidism (21.1%) and hyperthyroidism (11.9%), due to grade 1 and 2 events. The nature (severity, management, and outcome) of the immune-mediated AEs was similar to the nature of immune-mediated AEs previously seen with pembrolizumab monotherapy.

Safety data related to the MSI-H population in KN158 Cohort K (database cutoff 05-OCT-2020) and KN164 Cohorts A and B (database cutoff 09-SEP-2019) (n=475) were evaluated and are presented in text for each risk. Overall, the safety profile of pembrolizumab monotherapy in the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B was generally similar to the observed safety profile in the reference safety dataset.

Safety data related to the cervical cancer population in KN826 (database cutoff 03-MAY-2021) who received pembrolizumab plus chemotherapy ± bevacizumab (total n=307) were evaluated and are presented in text for each risk. Overall, the observed safety profile of pembrolizumab in combination with chemotherapy ± bevacizumab in cervical cancer was generally similar to the known safety profiles of pembrolizumab monotherapy and bevacizumab or the chemotherapy administered. The overall exposure-adjusted incidence of AEOSIs was higher

in the pembrolizumab + chemotherapy ± bevacizumab group compared with the pembrolizumab monotherapy reference safety dataset. Most AEOSIs reported in the pembrolizumab + chemotherapy ± bevacizumab group were mild to moderate in severity (Grade 1 or 2). The higher incidence of exposure-adjusted AEOSIs in the pembrolizumab + chemotherapy ± bevacizumab group was driven primarily by infusion reactions (13.4%), reflecting the contribution of pembrolizumab, chemotherapy and bevacizumab.

Safety data related to the TNBC population in KN522 (database cutoff 23-MAR-2021) (n=783) were evaluated and are presented in text for each risk. Overall, the safety profile of pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy in TNBC in KN522 was generally similar to the known safety profiles of pembrolizumab monotherapy and the chemotherapy administered. A higher frequency of immune-mediated events was observed in the pembrolizumab in combination with neoadjuvant chemotherapy group followed by continued adjuvant pembrolizumab monotherapy treatment group in comparison to the reference safety dataset. This was primarily driven by hypothyroidism (15.1%), infusion reactions (18%), and severe skin reactions (5.7%), reflecting the contribution of both pembrolizumab and the neoadjuvant chemotherapy administered. Most AEOSIs were grade 1 and 2. The nature (type, severity, management, and outcome) of the immune-mediated AEs observed with pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy in KN522 was similar to the nature of immune-mediated AEs previously seen with pembrolizumab monotherapy.

Safety data related to the melanoma population in KN716 (database cutoff 04-DEC-2020) (total n=483) were evaluated and are presented in text for each risk. Overall, the safety profile of pembrolizumab monotherapy in the melanoma population in KN716 was generally similar to the observed safety profile in the reference safety dataset. Increases in some immune-mediated AEs were observed. The nature (type, severity, management, and outcome) of the immune-mediated AEs observed with pembrolizumab in KN716 was similar to the nature of immune-mediated AEs previously seen with pembrolizumab monotherapy. The increases in some immune-mediated AEs observed in KN716 could be a reflection of the >2-fold longer duration of exposure to pembrolizumab in KN716 subjects compared to subjects in the reference safety dataset.

Safety data related to the gastric cancer population in KN811 (database cutoff 25-MAY-2022) who received pembrolizumab plus chemotherapy and trastuzumab (n=350) were evaluated and are presented in text for each risk. Overall, the safety profile of pembrolizumab combination therapy in the gastric cancer population in KN811 was generally similar to the observed safety profile in the reference safety dataset, except for a higher incidence of immune-mediated events primarily due to infusion-related reactions (16.0%) and pneumonitis (6.0%). The higher incidence of these events may be driven by the addition of trastuzumab and chemotherapy, since these are known risks for the SOC regimen. The longer duration of exposure compared to the reference safety dataset may also have contributed to the observed higher incidence of these AEOSIs. The nature (severity, management, and outcome) of the immune-mediated AEs was similar to the nature of immune-mediated AEs previously seen with pembrolizumab monotherapy.

Safety data related to the NSCLC population in KN091 (database cutoff date 20-Sep-2021) (total n=580 in pembrolizumab group) were evaluated and are presented in text for each risk. Overall, the safety and tolerability of pembrolizumab administered in KN091 is generally consistent with the known safety profile of pembrolizumab in the advanced cancer setting, except for a higher incidence of immune-mediated events primarily due to a higher cumulative frequency of hypothyroidism (20.7%) and hyperthyroidism (10.7%), due to grade 1 and 2 events. However, the incidences of hyperthyroidism and hypothyroidism in the KN091 pembrolizumab group are consistent with those reported in other adjuvant studies with pembrolizumab monotherapy in melanoma (KN716) and RCC (KN564) populations. The incidence of pneumonitis (6.9%) in pembrolizumab group was higher than in the pembrolizumab reference safety dataset but, consistent with the incidence of pneumonitis (3.8 to 8.3%) reported in individual studies of patients with NSCLC treated with pembrolizumab monotherapy. The increases in some immune-mediated AEs in KN091 could be a reflection of the >2-fold longer duration of exposure to pembrolizumab compared to subjects in the reference safety dataset. The nature (severity, management, and outcome) of the immune-mediated AEs was similar to the nature of immune-mediated AEs previously seen with pembrolizumab monotherapy.

Safety data related to the gastric cancer population in KN859 (database cutoff 03-OCT-2022) who received pembrolizumab plus chemotherapy (n=785) were evaluated and are presented in text for each risk. Overall, the safety profile of pembrolizumab combination therapy in the gastric cancer population in KN859 was generally similar to the observed safety profiles of pembrolizumab monotherapy and the chemotherapy administered, except for a higher incidence of immune-mediated event of hypothyroidism (15.3%) as compared to the RSD. After adjustment for exposure the event rates for hypothyroidism were similar to the RSD. The events of hypothyroidism in KN859, were mostly Grade 1 and 2 events and are generally consistent in nature with the events of hypothyroidism previously reported for pembrolizumab.

Safety data related to the BTC population in KN966 (database cutoff date: 15-Dec-2022) (total n=529) were evaluated and presented in text for each risk. Overall, safety and tolerability of pembrolizumab in combination with chemotherapy in the BTC population in KN966 was generally similar to the known safety profile of pembrolizumab monotherapy and the safety profile of chemotherapy. The AEOSI severity in most participants were Grade 1 or 2. The nature (type, severity, management, and outcome) of the immune-mediated AEs observed with pembrolizumab in combination with chemotherapy was similar to the nature of immune-mediated AEs previously observed with pembrolizumab monotherapy in the reference safety dataset.

Safety data related to the NSCLC population in KN671 (database cutoff date 29-Jul-2022) (total n=396) were evaluated and are presented in text for each risk. Overall, the safety and tolerability of pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy administered in KN671 is generally consistent with the known safety profile of pembrolizumab monotherapy and the known safety profile of the individual chemotherapy components.

Safety data related to the endometrial carcinoma population in KN868 (database cutoff 16-Dec-2022 for dMMR participants and 06-Dec-2022 for pMMR participants) (total n=382) who

received pembrolizumab plus chemotherapy were evaluated and presented in text for each risk. Overall, the safety profile observed with pembrolizumab in combination with chemotherapy in the endometrial carcinoma population in KN868 (dMMR and pMMR) was generally similar to the known safety profiles of pembrolizumab monotherapy or chemotherapy administered as monotherapy. No new indication-specific AEOSI were identified (ie, immune-mediated events causally associated with pembrolizumab) when pembrolizumab was administered concurrently with chemotherapy. The types and incidences of the most frequently reported AEOSI in the pembrolizumab plus chemotherapy group were generally consistent with the pembrolizumab monotherapy RSD, with the exception of infusion reactions. Infusion reactions were balanced between the pembrolizumab plus chemotherapy group and placebo plus chemotherapy group in KN868 and were primarily low grade. AEOSI in the pembrolizumab plus chemotherapy group were manageable with concomitant corticosteroids and/or treatment interruption/discontinuation.

Safety data related to the cervical cancer population in KNA18 (database cutoff 09-Jan-2023) who received pembrolizumab in combination with chemoradiotherapy (total n=528) were evaluated and presented in text for each risk. Overall, the observed safety profile of pembrolizumab in combination with chemoradiotherapy administered in KNA18 was generally consistent with the established safety profiles of pembrolizumab monotherapy and the cisplatin-based chemoradiotherapy regimen. The overall incidence of AEOSI was higher in the pembrolizumab plus chemoradiotherapy group compared with pembrolizumab monotherapy, primarily driven by low-grade events of hypothyroidism and hyperthyroidism. The nature (severity, outcome) of AEOSI observed in the pembrolizumab plus chemoradiotherapy group were generally consistent with pembrolizumab monotherapy. The overall increased incidence of some AEOSI observed in KNA18 may reflect a longer duration of exposure in the combination arm in KNA18 participants, as well as a background incidence of these AEs in the underlying population.

Safety data related to the mesothelioma population in KN483 and KNA17 (database cutoff 16-SEP-2022 and 21-SEP-2022, respectively) were evaluated and are presented for each risk. Overall, pembrolizumab in combination with pemetrexed/platinum chemotherapy in participants with previously untreated advanced and/or metastatic MPM has a manageable safety profile that is generally consistent with the established profile of pembrolizumab monotherapy and known safety profile of the individual chemotherapy components. No new safety concerns were identified. AEs were effectively managed by standard clinical practice, as applicable for treatment with pembrolizumab or pemetrexed/platinum chemotherapy.

Safety data related to the HNSCC population in KN689 (database cutoff date 25-JUL-2024) (pembrolizumab group n=361) were evaluated and are presented in text for each risk. Overall, the safety profile of pembrolizumab as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy is generally consistent with the known safety profile of pembrolizumab monotherapy and the known safety profile of radiation and chemotherapy except for a higher incidence of hypothyroidism as compared to the reference safety dataset. This higher incidence is anticipated in this study population due to the addition of RT in addition to a longer duration of exposure.

Safety data related to the NSCLC population in MK-3475A-D77 (database cutoff date 12-Jul-2024) dosed with pembrolizumab administered as MK-3475A subcutaneously (total n=251) were evaluated and are presented for each risk. The results from MK-3475-D77 show that the safety profile of pembrolizumab SC in combination with chemotherapy is consistent with the safety data of pembrolizumab monotherapy and chemotherapy, including data from other studies with participants with NSCLC receiving pembrolizumab IV and chemotherapy.

All identified and potential risks for pembrolizumab are reviewed routinely for a change in characterization (including severity, frequency and risk factors) in accordance with the EMA Guideline on good pharmacovigilance practices (GVP) Module V Risk management systems and Module IX Signal Management. Per this process, data regarding reports of immune-mediated AEOSIs with a fatal outcome were reviewed. Review of the data revealed that a number of AEOSIs had at least 1 event with a reported fatal outcome.

A series of database queries was done to identify all CT and postmarketing AEOSIs with a fatal outcome.

Clinical Trial (CT) Data:

The Aggregate Safety Evaluation (ASE) dataset, which is a locked aggregate CT safety database, was queried. The current 3Q2017 dataset contains all safety data for subjects who received pembrolizumab monotherapy in an open-label or unmasked, completed randomized trial for which there has been a database lock (DBL) on or before 30-Sep-2017 (n=7730). The ASE was queried to identify any CT AEOSI with a reported Common Terminology Criteria for Adverse Events (CTCAE) Grade 5 (fatal outcome) severity.

Global Safety Database and Postmarketing (PM) Data:

The global safety database contains clinical trial (including MAH sponsored clinical trials, Investigator Initiated Study Programs, MAH Sponsored Individual Patient Use, Business Partner Cases, and Academic Collaboration Studies) information for all serious events as well as other reportable events such as cancer, overdose, pregnancy, lactation, and protocol-specified events of clinical interest. The global safety database also contains all cases from postmarketing sources including health care providers, consumers and scientific literature as well as competent authorities worldwide. The safety database was queried as of 31-Dec-2017 to identify all Grade 5 CT AEOSIs from interventional, pembrolizumab monotherapy trials.

The global safety database was also queried as of 31-Dec-2017, to identify all PM and non-interventional reports containing a reported immune-mediated AEOSI with a fatal outcome. Two separate queries were done: one to identify reports with a fatal AEOSI only, excluding those with other fatal events due to disease progression/related terms, and another query including cases with other reported causes of death including those due to disease progression.

The results indicated that even with the broader global safety database analysis that included cases in which fatal AEOSIs were reported in the setting of disease progression, the number of cases with fatal immune-mediated AEOSIs remains low.

While the frequency of fatal immune-mediated AEOSIs is low, and while there remain some immune-mediated AEOSIs for which there are no reports of fatal outcome, based on the totality of data and medical importance of the information, the MAH updated the Warning and Precautions section of the prescribing information to add language to further qualify and communicate that immune-mediated AEOSIs may be fatal.

The immune-mediated adverse reactions are presumed to have a common etiology. Of these immune-mediated adverse reactions, information on severity and frequency data for pneumonitis, colitis, hepatitis, nephritis and endocrinopathies is presented, in the risk tables below as part of the important identified risk.

Table SVII.3.1.1: Details of Important Identified Risk: Immune-Mediated Adverse Reactions-Pneumonitis

Important Identified Risk:	Immune-mediated Pneumonitis
<u>Potential Mechanisms:</u>	Pembrolizumab is a potent and highly-selective humanized monoclonal antibody (mAb) of the Immunoglobulin G4 (IgG4)/kappa isotype directed to the programmed cell death-1 (PD-1) receptor and designed to directly block the interaction between the receptor and its ligands, PD-L1 and PD-L2. The PD-1 pathway represents a major immune control switch which may be engaged by tumor cells to overcome active T-cell immune surveillance. T-cell activation in proximity to normal tissue may lead to inflammation and injury to normal lung tissue.
<u>Evidence Source(s) and Strength of Evidence:</u>	Review of pembrolizumab clinical trial data, post-marketing experience and literature regarding pneumonitis represent sufficient evidence of a causal association with pembrolizumab exposure. CTD 2.7.4 pembrolizumab KN001 Database Cutoff Date: 18APR2014 pembrolizumab KN001 Database Cutoff Date for Lung: 23JAN2015 pembrolizumab KN002 Database Cutoff Date: 28FEB2015 pembrolizumab KN006 Database Cutoff Date: 03MAR2015 pembrolizumab KN010 Database Cutoff Date: 30SEP2015 pembrolizumab KN013 Database Cutoff Date for Hodgkin Lymphoma: 28SEP2018 pembrolizumab KN024 Database Cutoff Date: 10JUL2017 pembrolizumab KN087 Database Cutoff Date: 21MAR2019 pembrolizumab KN045 Database Cutoff Date: 07SEP2016 pembrolizumab KN052 Database Cutoff Date: 01SEP2016 pembrolizumab KN021 Database Cutoff Date Cohort A: 07NOV2016, Cohort G/C: 31MAY2017 pembrolizumab KN189 Database Cutoff Date: 08NOV2017 pembrolizumab KN040 Database Cutoff Date: 15MAY2017 pembrolizumab KN012 Database Cutoff Date: 26APR2016 pembrolizumab KN055 Database Cutoff Date: 22APR2016 pembrolizumab KN054 Database Cutoff Date: 02OCT2017 pembrolizumab KN407 Database Cutoff Date: 03APR2018 pembrolizumab KN426 Database Cutoff Date: 24AUG2018 pembrolizumab KN048 Database Cutoff Date: 13JUN2018 pembrolizumab KN042 Database Cutoff Date: 26FEB2018 pembrolizumab KN177 Database Cutoff Date: 19FEB2020 pembrolizumab KN204 Database Cutoff Date: 16JAN2020 pembrolizumab KN590 Database Cutoff Date: 02JUL2020 pembrolizumab KN355 Database Cutoff Date: 11DEC2019

Table SVII.3.1.1: Details of Important Identified Risk: Immune-Mediated Adverse Reactions-Pneumonitis

	<p>pembrolizumab KN581 Database Cutoff Date: 28AUG2020 pembrolizumab KN146 Database Cutoff Date: 18AUG2020 pembrolizumab KN775 Database Cutoff Date: 26OCT2020 pembrolizumab KN564 Database Cutoff Date: 14DEC2020 pembrolizumab KN158 Database Cutoff Date Cohort K: 05OCT2020 pembrolizumab KN164 Database Cutoff Date Cohorts A and B: 09SEP2019 pembrolizumab KN826 Database Cutoff Date: 03MAY2021 pembrolizumab KN522 Database Cutoff Date: 23MAR2021 pembrolizumab KN716 Database Cutoff Date: 04DEC2020 pembrolizumab KN811 Database Cutoff Date: 25MAY2022 pembrolizumab KN091 Database Cutoff Date: 20SEP2021 pembrolizumab KN859 Database Cutoff Date: 03OCT2022 pembrolizumab KN966 Database Cutoff Date: 15DEC2022 pembrolizumab KN671 Database Cutoff Date: 29JUL2022 pembrolizumab KN869 Database Cutoff Date: 13MAR2023 pembrolizumab KNA39 Database Cutoff Date: 08AUG2023 pembrolizumab KN868 Database Cutoff Date: 16DEC2022 for dMMR participants and 06DEC2022 for pMMR participants pembrolizumab KNA18 Database Cutoff Date: 09JAN2023 pembrolizumab KN483 Database Cutoff Date: 16SEP2022 pembrolizumab KNA17 Database Cutoff Date: 21SEP2022 pembrolizumab KN689 Database Cutoff Date: 25JUL2024 pembrolizumab MK-3475A-D77 Database Cutoff Date: 12JUL2024</p>								
<p>Characterisation of the Risk: Immune-mediated Pneumonitis</p>									
<p>Frequency with 95%CI</p>	<p align="center">95% Confidence Interval for the Overall Incidence (%) of Pneumonitis KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab</p> <table border="1" data-bbox="630 1255 1279 1430"> <thead> <tr> <th rowspan="2"></th> <th colspan="2">Cumulative MEL 001/002/006 + NSCLC 001/010</th> </tr> <tr> <th>n (%)</th> <th>95% CI</th> </tr> </thead> <tbody> <tr> <td>Pneumonitis</td> <td>94 (3.4)</td> <td>(2.7,4.1)</td> </tr> </tbody> </table> <p><u>Pembrolizumab IV</u> The overall number and proportion of patients with of pneumonitis in the NSCLC population in KN024 and KN042 (n=790) was 65 (8.2%) (95% CI- 6.4, 10.4). In individual studies of patients with NSCLC (total n=2022), the incidence of pneumonitis (all Grades) ranged from 3.8% to 8.3%. The proportion of patients with events of pneumonitis in KN024 and KN042, although somewhat more common than in the reference safety dataset, were generally consistent in nature with the proportion of events of pneumonitis previously reported for pembrolizumab. The overall number and proportion of patients with pneumonitis in the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488) was 22 (4.5%) (95% CI- 2.8, 6.7), which is generally similar to the proportion of patients with the event of pneumonitis in the reference safety dataset for melanoma and NSCLC. The overall number and proportion of patients with pneumonitis in the HL population in KN013, KN087, and KN204 (n=389) was 31 (8.0%) (95% CI-5.5, 11.1). These proportion of patients with events of pneumonitis, although somewhat more common than the reference safety dataset, were</p>		Cumulative MEL 001/002/006 + NSCLC 001/010		n (%)	95% CI	Pneumonitis	94 (3.4)	(2.7,4.1)
	Cumulative MEL 001/002/006 + NSCLC 001/010								
	n (%)	95% CI							
Pneumonitis	94 (3.4)	(2.7,4.1)							

Table SVII.3.1.1: Details of Important Identified Risk: Immune-Mediated Adverse Reactions-Pneumonitis

	<p>generally consistent in nature with the proportion of events of pneumonitis previously reported for pembrolizumab.</p> <p>The overall number and proportion of patients with pneumonitis in the UC population in KN052 (n=370) was 7 (1.9%) (95%CI-0.8,3.9) and in KN045 (n=266) was 11 (4.1%) (95%CI-2.1,7.3), which is generally similar to the proportion of patients with the event of pneumonitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with pneumonitis in the HNSCC population in KN040, KN012 and KN055 (n=609) was 23 (3.8%) (95% CI- 2.4,5.6), which is generally similar to the proportion of patients with the event of pneumonitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with pneumonitis in the HNSCC population in KN048 pembrolizumab monotherapy (n=300) and pembrolizumab plus chemotherapy (n=276) groups were 18 (6.0%) (95% CI- 3.6, 9.3) and 15 (5.4%) (95% CI-3.1,8.8) respectively, which is generally similar to the overall proportion of patients with the event of pneumonitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with pneumonitis in the melanoma population in KN054 (n=509) was 17 (3.3%) (95% CI-2.0, 5.3), which is generally similar to the proportion of patients with the events of pneumonitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with pneumonitis in the NSCLC population in KN407 and KN021 (Cohort A) (n=303) was 19 (6.3%) (95% CI-3.8, 9.6). The proportion of patients with events of pneumonitis in KN407 and KN021 (Cohort A), although somewhat more common than in the reference safety dataset, were generally consistent in nature with the proportion of events of pneumonitis previously reported for pembrolizumab.</p> <p>The overall number and proportion of patients with pneumonitis in the RCC population in KN426 pembrolizumab plus axitinib (n=429) was 12 (2.8%) (95% CI-1.5,4.8), which is similar to the proportion of patients with the event of pneumonitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with pneumonitis in the CRC population in KN177 (n=153) was 6 (3.9%) (95% CI-1.5, 8.3) which is generally similar to the proportion of patients with the event of pneumonitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with pneumonitis in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370) was 23 (6.2%) (95% CI-4,9.2), which is similar to the proportion of patients with the event of pneumonitis reported for the reference safety dataset. The proportion of patients with events of pneumonitis in KN590 were generally consistent in nature with the proportion of events of pneumonitis previously reported for pembrolizumab.</p> <p>The overall number and proportions of patients with pneumonitis in the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596) was 15 (2.5%) (95% CI-1.4,4.1), which is generally similar to the proportion of patients with the event of pneumonitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with pneumonitis in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352) was 19 (5.4%) (95% CI-3.3,8.3), which is generally similar to the proportion of patients with the event of pneumonitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with pneumonitis in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530) was 9 (1.7%) (95% CI- 0.8, 3.2), which is generally similar to the proportion of patients with the event of pneumonitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with pneumonitis in the RCC population in KN564 pembrolizumab monotherapy (n=488) was 11 (2.3%) (95% CI-1.1,4), which is generally similar to the proportion of patients with the event of pneumonitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with pneumonitis in the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475) was 15 (3.2%) (95% CI-1.8, 5.2) which is generally similar to the proportion of patients with the event of pneumonitis in the reference safety dataset.</p>
--	---

Table SVII.3.1.1: Details of Important Identified Risk: Immune-Mediated Adverse Reactions-Pneumonitis

	<p>The overall number and proportion of patients with pneumonitis in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307) was 6 (2.0%) (95% CI-0.7,4.2), which is generally similar to the proportion of patients with the event of pneumonitis in the reference safety dataset.</p> <p>The overall number and proportion of pneumonitis in the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) was 17 (2.2%) (95% CI-1.3,3.5), which is generally similar to the proportion of patients with the event of pneumonitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with pneumonitis in the melanoma population in KN716 pembrolizumab monotherapy (n=483) was 9 (1.9%) (95% CI-0.9, 3.5), which is generally similar to the proportion of patients with the event of pneumonitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with of pneumonitis in the gastric cancer population in KN811 pembrolizumab combination therapy (n=350) was 21 (6.0%) (95% CI-3.8, 9.0), which is generally similar to the proportion of patients with events of pneumonitis in the reference safety database.</p> <p>The overall number and proportion of patients with pneumonitis in the NSCLC population in KN091 pembrolizumab monotherapy (n=580) was 40 (6.9%) (95% CI-5.0,9.3), The proportion of patients with the event of pneumonitis in KN091, although somewhat more common than the reference safety dataset, was consistent with the proportion of events of pneumonitis previously reported for pembrolizumab monotherapy NSCLC studies.</p> <p>The overall number and proportion of patients with pneumonitis in the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785) was 25 (3.2%) (95% CI-2.1,4.7), which is generally similar to the proportion of patients with the events of pneumonitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with pneumonitis in the BTC population in KN966 pembrolizumab plus chemotherapy (n=529) was 26 (4.9%) (95% CI-3.2, 7.1), which is generally similar to the proportion of patients with the events of pneumonitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with pneumonitis in the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396) was 22 (5.6%) (95% CI-3.5, 8.3), which is more common than in the reference safety dataset. However, after adjusting for exposure, the rates for pneumonitis were similar to those reported in the reference dataset.</p> <p>The overall number and proportion of patients with pneumonitis in the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564) was 54 (9.6%) (95% CI-7.3, 12.3), which was higher than that in the reference safety dataset, reflecting the contribution of enfortumab vedotin and longer duration of exposure.</p> <p>The overall number and proportion of patients with pneumonitis in the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382) was 4 (1.0%) (95% CI-0.3, 2.7), which is generally similar to the proportion of patients with the events of pneumonitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with pneumonitis in the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528) was 7 (1.3%) (95% CI-0.5, 2.7), which is generally similar to the proportion of patients with the event of pneumonitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with pneumonitis in the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260) was 17 (6.5%) (95% CI-3.9, 10.3), which is generally similar to the proportion of patients with the events of pneumonitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with pneumonitis in the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361) were 20 (5.5%) (95% CI 3.4, 8.4) respectively, which is generally similar to the proportion of patients with the events of pneumonitis in the reference safety dataset.</p>
--	--

Table SVII.3.1.1: Details of Important Identified Risk: Immune-Mediated Adverse Reactions-Pneumonitis

	<p><u>Pembrolizumab SC</u></p> <p>The overall number and proportion of patients with pneumonitis in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251) was 13 (5.2%) (95% CI-2.8, 8.7), which is generally similar to the proportion of patients with the events of pneumonitis in the reference safety dataset.</p>																																		
<p>Outcomes</p>	<p>Percent overall, fatal, not resolved, resolved, and unknown for identified risk of pneumonitis – pembrolizumab (AEs for 30 days after discontinuation were included)</p> <p style="text-align: center;">Pneumonitis (AEOSI)</p> <p style="text-align: center;">KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment-Emergent)</p> <table border="1" data-bbox="488 611 1414 940"> <thead> <tr> <th colspan="2"></th> <th colspan="2">Cumulative NSCLC 001/010 + MEL 001/002/006</th> </tr> <tr> <th colspan="2"></th> <th>n</th> <th>(%)</th> </tr> </thead> <tbody> <tr> <td>Subjects in population</td> <td>Outcome</td> <td>2,799</td> <td></td> </tr> <tr> <td rowspan="7">With one or more adverse events</td> <td>Overall</td> <td>94</td> <td>(3.4)</td> </tr> <tr> <td>Fatal</td> <td>4</td> <td>(0.1)</td> </tr> <tr> <td>Not Resolved</td> <td>30</td> <td>(1.1)</td> </tr> <tr> <td>Resolved</td> <td>54</td> <td>(2.0)</td> </tr> <tr> <td>Resolving</td> <td>5</td> <td>(0.2)</td> </tr> <tr> <td>Sequelae</td> <td>1</td> <td>(0.0)</td> </tr> <tr> <td>Unknown</td> <td>0</td> <td>(0.0)</td> </tr> </tbody> </table> <p>Every subject is counted once on each applicable row. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the percent incidence specified in the report title, after rounding. Outcome: Resolved = RECOVERED/RESOLVED, Not resolved = NOT RECOVERED/NOT RESOLVED.</p> <p><u>Pembrolizumab IV</u></p> <p>In the NSCLC population in KN024 and KN042 (n=790), the outcomes for the 65 patients with events of pneumonitis were as follows: 33 (4.2%) resolved, 11 (1.4%) resolving, 18 (2.3%) not resolved, and 1 (0.1%) unknown. There were 2 (0.3%) patients with events of pneumonitis with a fatal outcome.</p> <p>In the NSCLC population in KN189 and KN021(Cohorts C and G) pembrolizumab plus chemotherapy (n=488), the outcomes for the 22 (4.5%) patients with events of pneumonitis were as follows: 13 (2.7%) resolved, 1 (0.2%) resolving, 1 (0.2%) resolved with sequelae, 4 (0.8%) not resolved. There were 3 (0.6%) patients with events of pneumonitis with a fatal outcome reported in KN189.</p> <p>In the HL population in KN013, KN087, and KN204 (n=389), the outcomes for the 31 (8.0%) patients with events of pneumonitis were as follows: 24 (6.2%) resolved, 1 (0.3%) resolving, and 6 (1.5%) not resolved. There were no patients with events of pneumonitis with a fatal outcome.</p> <p>In the UC population in KN052 (n=370), the outcomes for the 7 (1.9%) patients with events of pneumonitis were as follows: 3 (0.8%) resolved and 4 (1.1%) not resolved and in KN045 (n=266), the outcomes for the 11 (4.1%) patients with events of pneumonitis were as follows: 5 (1.9%) resolved, 1 (0.4%) resolving, 4 (1.5%) not resolved. There was 1 (0.4%) patient with an event of pneumonitis with a fatal outcome reported in KN045.</p> <p>In the HNSCC population in KN040, KN012 and KN055 (n=609), the outcomes for the 23 patients with events of pneumonitis were as follows: In the HNSCC population in KN040, KN012 and KN055 (n=609), the outcomes for the 23 patients with events of pneumonitis were as follows: 11 (1.8%) resolved, 3 (0.5%) resolving, 1 (0.2%) resolving with sequelae, 7 (1.1%) not resolved. There was 1 (0.2%) patient with an event of pneumonitis with a fatal outcome reported in KN055.</p> <p>In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), the outcomes for the 18 patients with events of pneumonitis were as follows: 11 (3.7%) resolved, 3 (1.0%) resolving, and 3 (1.0%) not resolved. There was 1 (0.3%) patient with an event of pneumonitis with a fatal outcome. In the HNSCC population in KN048 pembrolizumab plus chemotherapy group</p>			Cumulative NSCLC 001/010 + MEL 001/002/006				n	(%)	Subjects in population	Outcome	2,799		With one or more adverse events	Overall	94	(3.4)	Fatal	4	(0.1)	Not Resolved	30	(1.1)	Resolved	54	(2.0)	Resolving	5	(0.2)	Sequelae	1	(0.0)	Unknown	0	(0.0)
		Cumulative NSCLC 001/010 + MEL 001/002/006																																	
		n	(%)																																
Subjects in population	Outcome	2,799																																	
With one or more adverse events	Overall	94	(3.4)																																
	Fatal	4	(0.1)																																
	Not Resolved	30	(1.1)																																
	Resolved	54	(2.0)																																
	Resolving	5	(0.2)																																
	Sequelae	1	(0.0)																																
	Unknown	0	(0.0)																																

Table SVII.3.1.1: Details of Important Identified Risk: Immune-Mediated Adverse Reactions-Pneumonitis

<p>(n=276), the outcomes for the 15 patients with events of pneumonitis were as follows: 10 (3.6%) resolved, 2 (0.7%) resolving, 2 (0.7%) resolved with sequelae. There was 1 (0.4%) patient with an event of pneumonitis with a fatal outcome.</p> <p>In the melanoma population in KN054 (n=509), the outcome of the 17 patients with events of pneumonitis were as follows: 8 (1.6%) resolved and 9 (1.8%) not resolved. There were no patients with events of pneumonitis with a fatal outcome.</p> <p>In the NSCLC population in KN407 and KN021 (Cohort A) (n=303), the outcomes for the 19 patients with events of pneumonitis were as follows: 11 (3.6%) resolved, 2 (0.6%) resolving, 1 (0.3%) resolved with sequelae, 4 (1.3%) not resolved. There was 1 (0.3%) patient with an event of pneumonitis with a fatal outcome.</p> <p>In the RCC population in KN426 pembrolizumab plus axitinib (n=429), the outcomes for the 12 patients with events of pneumonitis were as follows: 1 (0.2%) fatal, 7 (1.6%) resolved, 2 (0.5%) not resolved and 2 (0.5%) resolving.</p> <p>In the CRC population in KN177 (n=153), the outcomes for the 6 patients with events of pneumonitis were as follows: 2 (1.3%) resolved, 1 (0.7%) resolving, 1 (0.7%) resolved with sequelae and 2 (1.3%) not resolved. There were no patients with events of pneumonitis with a fatal outcome.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), the outcomes for the 23 patients with events of pneumonitis were as follows: 11 (3%) resolved, 5 (1.4%) resolving, and 5 (1.4%) not resolved. There were 2 (0.5%) patients with events of pneumonitis with a fatal outcome.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), the outcomes for the 15 patients with events of pneumonitis were as follows: 11 (1.8%) resolved, 1 (0.2%) resolving, and 3 (0.5%) not resolved. There were no patients with events of pneumonitis with a fatal outcome.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), the outcomes for the 19 patients with events of pneumonitis were as follows: 7 (2%) resolved, 2 (0.6%) resolved with sequelae, 2 (0.6%) resolving, and 7 (2%) not resolved. There was 1 (0.3%) patient with an event of pneumonitis with a fatal outcome.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), the outcome for the 9 patients with events of pneumonitis was as follows: 5 (0.9%) resolved, 3 (0.6%) not resolved, and 1 (0.2%) resolving. There were no patients with events of pneumonitis with a fatal outcome.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488), the outcomes for the 11 patients with events of pneumonitis were as follows: 10 (2%) resolved and 1 (0.2%) not resolved. There were no patients with events of pneumonitis with a fatal outcome.</p> <p>In the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), the outcomes for the 15 patients with events of pneumonitis were as follows: 7 (1.5%) resolved, 3 (0.6%) resolving, 2 (0.4%) resolved with sequelae and 3 (0.6%) not resolved. There were no patients with events of pneumonitis with a fatal outcome.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), the outcomes for the 6 patients with events of pneumonitis were as follows: 2 (0.7%) not resolved, 2 (0.7%) resolving, and 2 (0.7%) resolved. There were no patients with events of pneumonitis with a fatal outcome.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), the outcomes for the 17 patients with events of pneumonitis were as follows: 13 (1.7%) resolved, 2 (0.3%) not resolved, and 1 (0.1%) resolving. There was 1 (0.1%) patient with an event of pneumonitis with a fatal outcome.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483). The outcomes for the 9 patients with events of pneumonitis were as follows: 5 (1%) resolved, 2 (0.4%) not resolved, and 2 (0.4%) resolving. There were no patients with events of pneumonitis with a fatal outcome.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), the outcomes for the 21 patients with events of pneumonitis were as follows: 13 (3.7%) resolved, 5</p>
--

Table SVII.3.1.1: Details of Important Identified Risk: Immune-Mediated Adverse Reactions-Pneumonitis

<p>(1.4%) not resolved, and 1 (0.3%) resolving. There were 2 (0.6%) patients with events of pneumonitis with a fatal outcome.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580), the outcomes for the 40 patients with events of pneumonitis were as follows: 27 (4.7%) resolved, 11 (1.9%) not resolved, and 2 (0.3%) resolved with sequelae. There were no patients with events of pneumonitis with a fatal outcome.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), the outcomes for the 25 patients with events of pneumonitis were as follows: 13 (1.7%) not resolved, 8 (1.0%) resolved, 2 (0.3%) resolving, and 1 (0.1%) resolved with sequelae. There was 1 (0.1%) patient with a fatal event of pneumonitis.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), the worst reported outcomes for the 26 patients with events of pneumonitis were as follows: 9 (1.7%) resolved, 9 (1.7%) not resolved, 6 (1.1%) resolving and 1 (0.2%) resolved with sequelae. There was 1 (0.2%) patient with an event of pneumonitis with a fatal outcome.</p> <p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy group (n=396), the outcomes for the 22 patients with events of pneumonitis were as follows: 10 (2.5%) resolved, 6 (1.5%) not resolved, 4 (1.0%) resolving, and 1 (0.3%) resolved with sequelae. There was 1 (0.3%) patient with an event of pneumonitis with a fatal outcome.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), the outcomes for the 54 patients with events of pneumonitis were as follows: 26 (4.6%) resolved, 3 (0.5%) resolved with sequelae, 16 (2.8%) not resolved, and 7 (1.2%) resolving. There were 2 (3.7%) patients with an event of pneumonitis with a fatal outcome.</p> <p>In the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382), the outcomes for patients with events of pneumonitis were not recorded per study protocol (managed by external collaboration).</p> <p>In the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528), the outcomes for the 7 patients with events of pneumonitis were as follows: 4 (0.8%) resolved, 2 resolving (0.4%), and 1 (0.2%) not resolved. There were no patients with an event of pneumonitis with a fatal outcome.</p> <p>In the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), the reported outcomes for the 17 patients with events of pneumonitis were as follows: 9 (3.5%) resolved, 8 (3.1%) not resolved. There were no patients with events of pneumonitis with a fatal outcome.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), the outcomes for the 20 patients with events of pneumonitis were as follows: 15 (4.2%) resolved, and 4 (1.1%) not resolved. There was 1 (1.1%) patient with an event of pneumonitis with a fatal outcome.</p> <p><u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251), the outcomes for the 13 patients with events of pneumonitis were as follows: 7 (2.8%) not resolved, 3 (1.2%) resolved, 2 (0.8%) resolving. There was 1 (0.4%) patient with an event of pneumonitis with a fatal outcome.</p> <p><u>Aggregate Review</u></p> <p>Review of the ASE dataset (n=7730) yielded 12 fatal cases of pneumonitis. Review of the global safety database yielded the following: 31 fatal cases from ongoing interventional monotherapy clinical trials (including the 12 cases from ASE) and 100 fatal cases from the PM or non-interventional environment. Of the 100, there were 84 cases in which fatal pneumonitis was the only grade 5 event and 16 cases that reported a grade 5 pneumonitis, as well as a grade 5 malignant neoplasm progression (MNP) or other related fatal event(s).</p>

Table SVII.3.1.1: Details of Important Identified Risk: Immune-Mediated Adverse Reactions-Pneumonitis

Seriousness	Percent serious for identified risk of pneumonitis - pembrolizumab (Serious AEs for 90 days after discontinuation were included.)	
	Pneumonitis - Serious (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment Emergent)	
		Cumulative NSCLC 001/010 + MEL 001/002/006
		n (%)
	Subjects in population	2,799
	with one or more adverse events	49 (1.8)
	Interstitial lung disease	3 (0.1)
	Pneumonitis	46 (1.6)
	Every subject is counted a single time for each applicable row and column.	
	A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns meets the incidence criterion in the report title, after rounding.	
Include all treated subjects in KN001 Part B1, B2, B3, D, C, F1, F2, F3 and all subjects in KN002 treated with Pembrolizumab in the original phase and all subjects in KN006 and KN010 treated with Pembrolizumab.		
<u>Pembrolizumab IV</u>		
Of the 65 patients with events of pneumonitis in the NSCLC population in KN024 and KN042 (n=790), 37 (4.7%) experienced serious events of pneumonitis.		
Of the 22 patients with events of pneumonitis in the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), 13 (2.7%) experienced serious events of pneumonitis.		
Of the 31 patients with events of pneumonitis in the HL population in KN013, KN087, and KN204 (n=389), 15 (3.9%) experienced serious events of pneumonitis.		
Of the 7 patients with events of pneumonitis in the UC population in KN052 (n=370), 3 (0.8%) experienced serious events of pneumonitis and of the 11 patients with events of pneumonitis in the UC population in KN045 (n=266), 7 (2.6%) experienced serious events of pneumonitis.		
Of the 23 patients with events of pneumonitis in the HNSCC population in KN040, KN012 and KN055 (n=609), 12 (2%) experienced serious events of pneumonitis.		
Of the 18 patients with events of pneumonitis in the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), 4 (1.3%) were serious. Of the 15 events of pneumonitis in the HNSCC pembrolizumab plus chemotherapy group (n=276), 5 (1.8%) experienced serious events of pneumonitis.		
Of the 17 patients with events of pneumonitis in the melanoma population in KN054 (n=509), 8 (1.6%) experienced serious events of pneumonitis.		
Of the 19 patients with events of pneumonitis in the NSCLC population in KN407 and KN021 (Cohort A) (n=303), 9 (3.0%) experienced serious events of pneumonitis.		
Of the 12 patients with events of pneumonitis in the RCC population in KN426 pembrolizumab plus axitinib (n=429), 6 (1.4%) experienced serious events of pneumonitis.		
Of the 6 patients with events of pneumonitis in the CRC population in KN177 (n=153), 1 (0.7%) experienced a serious event of pneumonitis.		
Of the 23 patients with events of pneumonitis in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370) 14 (3.8%) experienced serious events of pneumonitis.		
Of the 15 patients with events of pneumonitis in the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), 7 (1.2%) experienced serious events of pneumonitis.		
Of the 19 patients with events of pneumonitis in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), 9 (2.6%) experienced serious events of pneumonitis.		
There were 4 (0.8%) patients with serious events of pneumonitis reported in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530).		

Table SVII.3.1.1: Details of Important Identified Risk: Immune-Mediated Adverse Reactions-Pneumonitis

	<p>Of the 11 patients with events of pneumonitis in the RCC population in KN564 pembrolizumab monotherapy (n=488), 5 (1%) experienced serious events of pneumonitis.</p> <p>Of the 15 patients with events of pneumonitis in the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475) 6 (1.3%) experienced serious events of pneumonitis.</p> <p>Of the 6 patients with events of pneumonitis in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), 1 (0.3%) experienced a serious event of pneumonitis.</p> <p>Of the 17 patients with events of pneumonitis in the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), 9 (1.1%) experienced serious events of pneumonitis.</p> <p>Of the patients with 9 events of pneumonitis in the melanoma population in KN716 pembrolizumab monotherapy (n=483), 3 (0.6%) experienced serious events of pneumonitis.</p> <p>Of the 21 patients with events of pneumonitis in the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), 8 (2.3%) experienced serious events of pneumonitis.</p> <p>Of the 40 patients with events of pneumonitis in the NSCLC population in KN091 pembrolizumab monotherapy (n=580), 16 (2.8%) experienced serious events of pneumonitis.</p> <p>Of the 25 patients with events of pneumonitis in the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), 12 (1.5%) experienced a serious event of pneumonitis.</p> <p>Of the 26 patients with events of pneumonitis in the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), 8 (1.5%) experienced serious events of pneumonitis.</p> <p>Of the 22 patients with events of pneumonitis in the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396), 11 (2.8%) experienced serious events of pneumonitis.</p> <p>Of the 54 patients with events of pneumonitis in the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), 22 (3.9%) experienced serious events of pneumonitis.</p> <p>Of the 4 patients with events of pneumonitis in the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382), 1 (0.3%) experienced a serious event of pneumonitis.</p> <p>Of the 7 patients with events of pneumonitis in the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528), 2 (0.4%) experienced serious events of pneumonitis.</p> <p>Of the 17 patients with events of pneumonitis in the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), 10 (3.8%) experienced serious events of pneumonitis.</p> <p>Of the 20 patients with events of pneumonitis in the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), 6 (1.7%) experienced serious events of pneumonitis.</p> <p><u>Pembrolizumab SC</u></p> <p>Of the 13 patients with events of pneumonitis in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251), 5 (2.0%) experienced serious events of pneumonitis.</p>
--	---

Table SVII.3.1.1: Details of Important Identified Risk: Immune-Mediated Adverse Reactions-Pneumonitis

Severity and Nature of the Risk	Identified treatment emergent adverse events (AE) of pneumonitis (all events) in participants summarized by grade– pembrolizumab (Table includes events for 30 days after discontinuation)		
	Pneumonitis (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment Emergent)		
		Cumulative NSCLC 001/010 + MEL 001/002/006	
		n	(%)
	Subjects in population	2,799	
	with one or more adverse events	94	(3.4)
	Interstitial lung disease	7	(0.3)
	Grade 1	3	(0.1)
	Grade 2	2	(0.1)
	Grade 3	1	(0.0)
	Grade 5	1	(0.0)
	Pneumonitis	87	(3.1)
	Grade 1	19	(0.7)
	Grade 2	34	(1.2)
	Grade 3	24	(0.9)
	Grade 4	7	(0.3)
	Grade 5	3	(0.1)
	Every subject is counted a single time for each applicable specific adverse event. A subject with multiple adverse events within a system organ class is counted a single time for that system organ class. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the incidence specified in the report title, after rounding. Only the highest reported grade of a given adverse event is counted for the individual subject. Grades are based on National Cancer Institute (NCI) Common Terminology Criteria for Adverse Events (CTCAE) version 4.0.		
	<u>Pembrolizumab IV</u>		
	In the NSCLC population in KN024 and KN042 (n=790), there were 19 (2.4%) patients with Grade 3, 6 (0.8%) patients with Grade 4 and 2 (0.3%) patients with Grade 5 events of pneumonitis.		
	In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), there were 8 (1.6%) patients with Grade 3, 1 (0.2%) patient with Grade 4, and 3 (0.6%) patients with Grade 5 events of pneumonitis.		
	In the HL population in KN013, KN087, and KN204 (n=389), there were 6 (1.5%) patients with Grade 3, 3 (0.8%) patients with Grade 4 and no patients with Grade 5 events of pneumonitis.		
	In the UC population in KN052 (n=370), there were 3 (0.8%) patients with Grade 3 events and no patients with Grade 4 or 5 events of pneumonitis; in the UC population in KN045 (n=266), there were 5 (1.9%) patients with Grade 3 events, no patients with Grade 4 events, and 1 (0.4%) patient with a Grade 5 event of pneumonitis.		
In the HNSCC population in KN040, KN012 and KN055 (n=609), there were 6 (1.0%) patients with Grade 3 events and 1 (0.2%) patient with a Grade 5 event of pneumonitis.			
In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), there were 4 (1.3%) patients with Grade 3 events and 1 (0.3%) patient with a Grade 5 event of pneumonitis; in the pembrolizumab plus chemotherapy group (n=276), there were 3 (1.1%) patients with Grade 3 events and 1 (0.4%) patient with a Grade 5 events of pneumonitis.			
In the melanoma population in KN054 (n=509), there were 4 (0.8%) patients with Grade 3 events of pneumonitis reported. There were no patients with Grade 4 or 5 events of pneumonitis.			
In the NSCLC population in KN407 and KN021 (Cohort A) (n=303), there were 6 (2.0%) patients with Grade 3, no patients with Grade 4, and 1 (0.3%) patient with a Grade 5 event of pneumonitis.			
In the RCC population in KN426 pembrolizumab plus axitinib (n=429), there was 1 (0.2%) patient with a Grade 3 event and 1 (0.2%) patient with a Grade 5 event of pneumonitis reported.			

Table SVII.3.1.1: Details of Important Identified Risk: Immune-Mediated Adverse Reactions-Pneumonitis

	<p>In the CRC population in KN177 (n=153) there were no Grade 3-5 events of pneumonitis reported.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370) there were 7 (1.9%) patients with Grade 3 events, no patients with Grade 4 events, and 2 (0.5%) patients with Grade 5 events of pneumonitis reported.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), there were 5 (0.8%) patients with Grade 3 events, and 1 (0.2%) patient with a Grade 4 event. There were no patients with Grade 5 events of pneumonitis reported.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352) there were 5 (1.4%) patients with Grade 3 events, 1 (0.3%) patient with a Grade 4 event, and 1 (0.3%) patient with a Grade 5 event of pneumonitis reported.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530) there were 4 (0.8%) patients with Grade 3 and no patients with Grade 4 or 5 events of pneumonitis reported.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488) there were 2 (0.4%) patients with Grade 3, 2 (0.4%) patients with Grade 4, and no patients with Grade 5 events of pneumonitis reported.</p> <p>In the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), there were 5 (1.1%) patients with Grade 3 events of pneumonitis. There were no patients with Grade 4 or 5 events of pneumonitis reported.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), there was 1 (0.3%) patient with a Grade 3 and no patients with Grade 4 or 5 events of pneumonitis reported.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) there were 6 (0.8%) patients with Grade 3 events, no patients with Grade 4 events, and 1 (0.1%) patient with a Grade 5 event of pneumonitis reported.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483) there was 1 (0.2%) patient with a Grade 3 and no patients with Grade 4 or 5 events of pneumonitis reported.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350) there were 3 (0.9%) patients with a Grade 3, 1 (0.3%) patient with a Grade 4, and 2 (0.6%) patients with Grade 5 events of pneumonitis reported.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580), there were 6 (1.0%) patients with Grade 3, 2 (0.3%) patients with Grade 4, and no patients with Grade 5 events of pneumonitis reported.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), there were 8 (1.0%) patients with a Grade 3, 1 (0.1%) patient with a Grade 4, and 1 (0.1%) patient with a Grade 5 event of pneumonitis reported.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529, there were 5 (0.9%) patients with Grade 3 events, no patients with Grade 4 events, and 1 (0.2%) patient with a Grade 5 event of pneumonitis reported.</p> <p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396) there were 6 (1.5%) patients with Grade 3, 1 (0.3%) patient with a Grade 4, and 1 (0.3%) patient with a Grade 5 event of pneumonitis reported.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), there were 17 (3.0%) patients with Grade 3 events, 2 (0.4%) patients with Grade 4 events, and 2 (0.4%) patients with a Grade 5 event of pneumonitis reported.</p> <p>In the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382) there were 2 (0.5%) patients with Grade 3 events of pneumonitis. There were no patients with Grade 4 or 5 events of pneumonitis reported.</p> <p>In the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528) there was 1 (0.2%) patient with a Grade 3 event, and 1 (0.2%) patient with a Grade 4 event of pneumonitis reported. There were no Grade 5 events of pneumonitis reported.</p>
--	--

Table SVII.3.1.1: Details of Important Identified Risk: Immune-Mediated Adverse Reactions-Pneumonitis

	<p>In the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), there were 2 (0.8%) patients with Grade 3 events. There were no patients with Grade 4 or 5 events of pneumonitis reported.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), there were 3 (0.8%) patients with Grade 3 events, 2 patients (0.6%) with Grade 4 events, and 1 (0.3%) patient with a Grade 5 event of pneumonitis.</p> <p><u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251), there were 5 (2.0%) patients with Grade 3, and 1 (0.4%) patient with a Grade 5 event of pneumonitis reported. There were no patients with Grade 4 events of pneumonitis reported.</p> <p>In the global safety database through 31-DEC-2017, (pembrolizumab monotherapy trials only), there were 31 Grade 5 clinical trial reports of pneumonitis.</p> <p>Review of pembrolizumab clinical trial data from ongoing studies and postmarketing data regarding immune-mediated pneumonitis, including fatal cases, is consistent with the risk as characterized in the RMP.</p>
<u>Risk Factors and Risk Groups:</u>	<p>Patients with a history of non-infectious pneumonitis that required steroids or current pneumonitis were excluded from the clinical trials. These patients are considered to be a risk group for the development of pneumonitis; in the interim analysis of the KN001 NSCLC cohort, possible risk factors identified that might predispose subjects to pneumonitis were a documented history of prior thoracic radiation to the chest (≥ 30Gy). According to the literature, risk factors for interstitial lung disease may include occupational exposure to toxins, chest irradiation, some chemotherapies, smoking and advanced age.</p>
<u>Preventability:</u>	<p>Although the development of pneumonitis cannot be completely prevented; patients should be monitored for signs and symptoms of pneumonitis. Suspected pneumonitis should be confirmed with radiographic imaging and other causes excluded. Withholding/discontinuation of pembrolizumab and appropriate medical intervention including corticosteroids following the onset of pneumonitis may result in recovery.</p>
<u>Impact on the Risk-Benefit Balance of the Product:</u>	<p>Severe pneumonitis may be life-threatening or fatal in individual patients. However, given the fatal outcome of untreated cancer, the risk of pneumonitis which generally can be managed, is outweighed by the potential benefit.</p>
<u>Public Health Impact:</u>	<p>This risk has minimal public health impact outside its effect on individual patients.</p>

Table SVII.3.1.2: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Colitis

<u>Important Identified Risk:</u>	<p>Immune-mediated Colitis</p>
<u>Potential Mechanisms:</u>	<p>Pembrolizumab is a potent and highly-selective humanized monoclonal antibody (mAb) of the Immunoglobulin G4 (IgG4)/kappa isotype directed to the programmed cell death-1 (PD-1) receptor and designed to directly block the interaction between the receptor and its ligands, PD-L1 and PD-L2. The PD-1 pathway represents a major immune control switch which may be engaged by tumor cells to overcome active T-cell immune surveillance. A potential mechanism of action is T-cell activation in proximity to normal tissue resulting in inflammation and injury to normal GI tissue.</p>
<u>Evidence Source(s) and Strength of Evidence:</u>	<p>Review of pembrolizumab clinical trial data, post-marketing experience and literature regarding colitis represent sufficient evidence of a causal association with pembrolizumab exposure.</p> <p>CTD 2.7.4</p>

Table SVII.3.1.2: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Colitis

	Please reference evidence source cited in Table SVII.3.1.1; Details of Important Identified Risk: Immune-Mediated Pneumonitis										
Characterisation of the Risk: Immune-mediated Colitis											
Frequency with 95%CI	<p style="text-align: center;">95% Confidence Interval for the Overall Incidence (%) of Colitis KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab</p> <table border="1" data-bbox="487 520 1421 646"> <thead> <tr> <th data-bbox="487 520 841 583"></th> <th colspan="2" data-bbox="841 520 1421 583">Cumulative MEL 001/002/006 + NSCLC 001/010</th> </tr> <tr> <th data-bbox="487 583 841 615"></th> <th data-bbox="841 583 1096 615">n (%)</th> <th data-bbox="1096 583 1421 615">95% CI</th> </tr> </thead> <tbody> <tr> <td data-bbox="487 615 841 646">Colitis</td> <td data-bbox="841 615 1096 646">48 (1.7)</td> <td data-bbox="1096 615 1421 646">(1.3,2.3)</td> </tr> </tbody> </table> <p><u>Pembrolizumab IV</u></p> <p>The overall number and proportion of patients with colitis in the NSCLC population in KN024 and KN042 (n=790) was 13 (1.6%) (95% CI-0.9, 2.8), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with colitis in the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488) was 14 (2.9%) (95% CI- 1.6, 4.8), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with colitis in the HL population in KN013, KN087, and KN204 (n=389) was 6 (1.5%) (95%CI-0.6, 3.3), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with colitis in the UC population in KN052 (n=370) was 9 (2.4%) (95%CI-1.1,4.6) and in KN045 (n=266) was 6 (2.3%) (95%CI-0.8,4.8), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with colitis in the HNSCC population in KN040, KN012 and KN055 (n=609) was 4 (0.7%) (95% CI- 0.2,1.7), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with colitis in the HNSCC population in KN048 pembrolizumab monotherapy (n=300) and pembrolizumab plus chemotherapy (n=276) groups were 3 (1.0%) (95% CI- 0.2, 2.9) and 7 (2.5) (95% CI- 1.5,2) respectively, which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with colitis in the melanoma population in KN054 (n=509) was 19 (3.7%) (95% CI-2.3, 5.8). The proportion of patients with events of colitis in KN054 although somewhat more common than in the reference safety dataset were generally consistent in nature with the proportion of patients with events of colitis previously reported for pembrolizumab.</p> <p>The overall number and proportion of patients with colitis in the NSCLC population in KN407 and KN021 (Cohort A) (n=303) was 9 (3.0%) (95% CI-1.4, 5.6), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with colitis in the RCC population in KN426 pembrolizumab plus axitinib (n=429) was 11 (2.6%) (95% CI-1.3,4.5), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with colitis in the CRC population in KN177 (n=153) was 10 (6.5%) (95% CI-3.2,11.7). Colitis was reported more frequently in the KN177 pembrolizumab safety dataset compared with the RSD (6.5% vs 1.7%, respectively); however, when adjusted for exposure, the incidences were similar (6.1 vs 4.1 events per 100 person-years)</p>			Cumulative MEL 001/002/006 + NSCLC 001/010			n (%)	95% CI	Colitis	48 (1.7)	(1.3,2.3)
	Cumulative MEL 001/002/006 + NSCLC 001/010										
	n (%)	95% CI									
Colitis	48 (1.7)	(1.3,2.3)									

Table SVII.3.1.2: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Colitis

	<p>and the proportion of events were generally consistent in nature with the proportion of events of colitis previously reported for pembrolizumab.</p> <p>The overall number and proportion of patients with colitis in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370) was 8 (2.2%) (95%CI-0.9,4.2), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with colitis in the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596) was 11 (1.8) (95% CI- 0.9,3.3), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with colitis in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352) was 9 (2.6%) (95% CI-1.2, 4.8), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with colitis in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530) was 26 (4.9%) (95% CI-3.2, 7.1). The proportion of patients with events of colitis in KN146 and KN775, although higher in frequency than in the reference safety dataset, were generally consistent in nature with the proportion of events of colitis previously reported for pembrolizumab.</p> <p>The overall number and proportion of patients with colitis in the RCC population in KN564 pembrolizumab monotherapy (n=488) was 8 (1.6%) (95% CI-0.7,3.2), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with colitis in the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475) was 10 (2.1%) (95% CI-1.0, 3.8) which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with colitis in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307) was 16 (5.2%) (95% CI-3.0, 8.3). Although more common than in the reference safety dataset, the proportion of events of colitis were generally consistent in nature with the proportion of events of colitis previously reported for pembrolizumab.</p> <p>The overall number and proportion of patients with colitis in the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) was 13 (1.7%) (95% CI-0.9,2.8), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with colitis in the melanoma population in KN716 pembrolizumab monotherapy (n=483) was 17 (3.5%) (95% CI-2.1, 5.6), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with colitis in the gastric cancer population in KN811 pembrolizumab combination therapy (n=350) was 17 (4.9%) (95% CI 2.9, 7.7). Colitis was reported more frequently in the KN811 pembrolizumab safety dataset compared with the RSD, however when adjusted for exposure, the incidences were similar (0.5 vs 0.3 events per 100 person-months) and the proportion of patients with events of colitis were generally consistent in nature with the proportion of patients with events of colitis previously reported for pembrolizumab.</p> <p>The overall number and proportion of patients with colitis in the NSCLC population in KN091 pembrolizumab monotherapy (n=580) was 14 (2.4%) (95% CI-1.3,4.0), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with colitis in the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785) was 26 (3.3%) (95% CI-2.2, 4.8) which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with colitis in the BTC population in KN966 pembrolizumab plus chemotherapy (n=529) was 9 (1.7%) (95% CI-0.8,3.2), which is the same as the proportion of patients with events of colitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with colitis in the NSCLC population in KN671 pembrolizumab in combination with chemotherapy neoadjuvant treatment followed by continued</p>
--	--

Table SVII.3.1.2: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Colitis

	<p>adjuvant pembrolizumab monotherapy (n=396) was 5 (1.3%) (95% CI-0.4,2.9), which is generally similar to the overall proportion of patients with events of colitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with colitis in the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564) was 18 (3.2%) (95% CI-1.9%, 5.0%), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with colitis in the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382) was 7 (1.8%) (95% CI-0.7, 3.7), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset</p> <p>The overall number and proportion of patients with colitis in the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528) was 14 (2.7%) (95% CI-1.5, 4.4), which is generally similar to the overall frequency in the reference safety dataset.</p> <p>The overall number and proportion of patients with colitis in the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260) was 6 (2.3%) (95% CI-0.9,5), which is generally similar to the overall frequency in the reference safety dataset.</p> <p>The overall number and proportion of patients with colitis in the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361) group were 9 (2.5%) (95% CI- 1.1, 4.7), which is generally similar to the proportion of patients with the event of colitis in the reference safety dataset.</p> <p><u>Pembrolizumab SC</u></p> <p>The overall number and proportion of patients with colitis in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251) was 3 (1.2%) (95% CI-0.2, 3.5), which is generally similar to the overall proportion of patients with events of colitis in the reference safety dataset.</p>																															
<p>Outcomes</p>	<p>Percent overall, fatal, not resolved, resolved, and unknown for identified risk of colitis - pembrolizumab (Table includes events for 30 days after discontinuation.)</p> <p style="text-align: center;">Colitis (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment Emergent)</p> <table border="1" data-bbox="487 1192 1419 1507"> <thead> <tr> <th colspan="2"></th> <th colspan="2">Cumulative NSCLC 001/010 + MEL 001/002/006</th> </tr> <tr> <th></th> <th>Outcome</th> <th>n</th> <th>(%)</th> </tr> </thead> <tbody> <tr> <td>Subjects in population</td> <td></td> <td>2,799</td> <td></td> </tr> <tr> <td rowspan="6">With one or more adverse events</td> <td>Overall</td> <td>48</td> <td>(1.7)</td> </tr> <tr> <td>Fatal</td> <td>0</td> <td>(0.0)</td> </tr> <tr> <td>Not Resolved</td> <td>6</td> <td>(0.2)</td> </tr> <tr> <td>Resolved</td> <td>41</td> <td>(1.5)</td> </tr> <tr> <td>Resolving</td> <td>1</td> <td>(0.0)</td> </tr> <tr> <td>Unknown</td> <td>0</td> <td>(0.0)</td> </tr> </tbody> </table> <p>Every subject is counted once on each applicable row. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the percent incidence specified in the report title, after rounding. Outcome: Resolved = RECOVERED/RESOLVED, Not resolved = NOT RECOVERED/NOT RESOLVED.</p> <p><u>Pembrolizumab IV</u></p> <p>In the NSCLC population in KN024 and KN042 (n=790), the outcomes of the 13 patients with events of colitis were as follows: 11 (1.4%) resolved, 2 (0.3%) not resolved. There were no patients with events of colitis with a fatal outcome.</p> <p>In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), the outcomes of the 14 patients with events of colitis were as follows: 10</p>			Cumulative NSCLC 001/010 + MEL 001/002/006			Outcome	n	(%)	Subjects in population		2,799		With one or more adverse events	Overall	48	(1.7)	Fatal	0	(0.0)	Not Resolved	6	(0.2)	Resolved	41	(1.5)	Resolving	1	(0.0)	Unknown	0	(0.0)
		Cumulative NSCLC 001/010 + MEL 001/002/006																														
	Outcome	n	(%)																													
Subjects in population		2,799																														
With one or more adverse events	Overall	48	(1.7)																													
	Fatal	0	(0.0)																													
	Not Resolved	6	(0.2)																													
	Resolved	41	(1.5)																													
	Resolving	1	(0.0)																													
	Unknown	0	(0.0)																													

Table SVII.3.1.2: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Colitis

	<p>(2.1%) resolved, 4 (0.8%) not resolved. There were no patients with events of colitis with a fatal outcome.</p> <p>In the HL population in KN013, KN087, and KN204 (n=389), the outcomes of the 6 (1.5%) patients with events of colitis were as follows: 6 (1.5%) resolved. There were no patients with events of colitis with a fatal outcome.</p> <p>In the UC population in KN052 (n=370), the outcomes for the 9 (2.4%) patients with events of colitis were as follows: 5 (1.4%) resolved, 3 (0.8%) not resolved, 1 (0.3%) resolving and in KN045 (n=266), the outcomes for the 6 (2.3%) patients with events of colitis were as follows: 6 (2.3%) resolved. There were no patients with events of colitis with a fatal outcome.</p> <p>In the HNSCC population in KN040, KN012 and KN055 (n=609), the outcomes for the 4 patients with events were as follows: 3 (0.5%) resolved and 1 (0.2%) not resolved. There were no patients with events of colitis with a fatal outcome.</p> <p>In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), the outcomes for the 3 patients with events of colitis were as follows: 1 (0.3%) resolved, 1 (0.3%) resolved with sequelae and 1 (0.3%) resolving. There were no fatal outcomes. In the HNSCC population in KN048 pembrolizumab plus chemotherapy group (n=276), the outcomes for the 7 patients with events of colitis were as follows: 5 (1.8%) resolved and 2 (0.7%) not resolved. There were no patients with events of colitis with a fatal outcome.</p> <p>In the melanoma population in KN054 (n=509), the outcomes for the 19 patients with events of colitis were as follows: 10 (2.0%) not resolved, 8 (1.6%) resolved, and 1 (0.2%) resolved with sequelae. There were no patients with events of colitis with a fatal outcome.</p> <p>In the NSCLC population in KN407 and KN021 (Cohort A) (n=303), the outcomes of the 9 patients with events of colitis were as follows: 8 (2.6%) resolved, 1 (0.3%) resolving. There were no patients with events of colitis with a fatal outcome.</p> <p>In the RCC population in KN426 pembrolizumab plus axitinib (n=429), the outcomes for the 11 patients with events of colitis were as follows: 8 (1.9%) resolved, 1 (0.2%) not resolved, 1 (0.2%) resolving and 1 (0.2%) resolved with sequelae. There were no patients with events of colitis with a fatal outcome.</p> <p>In the CRC population in KN177 (n=153), the outcomes for the 10 patients with events of colitis were as follows: 9 (5.9%) resolved, and 1 (0.7%) not resolved. There were no patients with events of colitis with a fatal outcome.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), the outcomes of the 8 patients with events of colitis were as follows: 6 (1.6%) resolved and 2 (0.5%) resolving.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), the outcomes for the 11 patients with events of colitis were as follows: 8 (1.3%) resolved, 2 (0.3%) not resolved, and 1 (0.2%) resolved with sequelae. There were no patients with events of colitis with a fatal outcome.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), the outcomes for the 9 patients with events of colitis were as follows: 6 (1.7%) resolved, 2 (0.6%) not resolved, and 1 (0.3%) resolving. There were no patients with events of colitis with a fatal outcome.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), the outcomes for the 26 patients with events of colitis were as follows: 16 (3.0%) resolved, 7 (1.3%) not resolved, 1 (0.2%) resolving and 1 (0.2%) resolved with sequelae. There was 1 (0.2%) patient with an event of colitis with a fatal outcome.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488), the outcomes for the 8 patients with events of colitis were as follows: 6 (1.2%) resolved, 1 (0.2%) resolved with sequelae, and 1 (0.2%) not resolved. There were no patients with events of colitis with a fatal outcome.</p> <p>In the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), the outcome for the 10 (2.1%) patients with events of colitis was resolved. There were no patients with events of colitis with a fatal outcome.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), the outcomes for the 16 patients with events of colitis were as follows: 11 (3.6%) resolved, 2 (0.7%)</p>
--	--

Table SVII.3.1.2: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Colitis

	<p>resolving, and 3 (1.0%) not resolved. There were no patients with events of colitis with a fatal outcome.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), the outcomes for the 13 patients with events of colitis were as follows: 12 (1.5%) resolved, and 1 (0.1%) not resolved. There were no patients with events of colitis with a fatal outcome.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), the outcome for the 17 patients with events of colitis were as follows: 12 (2.5%) resolved, 2 (0.4%) not resolved, 2 (0.4%) resolving, and 1 (0.2%) resolved with sequelae. There were no patients with events of colitis with a fatal outcome.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), the outcomes for the 17 patients with events of colitis were as follows: 12 (3.4%) resolved, 3 (0.9%) not resolved, 1 (0.3%) resolving and 1 (0.3%) resolved with sequelae. There were no patients with events of colitis with a fatal outcome.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580), the outcomes for the 14 patients with events of colitis were as follows: 12 (2.1%) resolved, 1 (0.2%) resolved with sequelae, and 1 (0.2%) not resolved. There were no patients with events of colitis with a fatal outcome.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), the outcomes for the 26 patients with events of colitis were as follows: 22 (2.8%) resolved, 2 (0.3%) not resolved, and 2 (0.3%) resolving. There were no patients with an event of colitis with a fatal outcome.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), the worst reported outcomes for the 9 patients with events of colitis were as follows: 7 (1.3%) resolved, 1 (0.2%) not resolved and 1 (0.2%) resolving. There were no patients with events of colitis with a fatal outcome.</p> <p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy group (n=396), the outcomes for the 5 patients with events of colitis were as follows: 4 (1.0%) resolved, and 1 (0.3%) resolving. There were no patients with events of colitis with a fatal outcome.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), the outcomes for the 18 patients with events of colitis were as follows: 14 (2.5%) resolved, 3 (0.5%) not resolved, and 1 (0.2%) resolving. There were no patients with an event of colitis with a fatal outcome.</p> <p>In the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382), the outcomes for patients with events of colitis were not recorded per study protocol (managed by external collaboration).</p> <p>In the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528), the outcomes for the 14 patients with events of colitis were as follows: 9 (1.7%) resolved, 3 (0.6%) not resolved, and 2 (0.4%) resolving. There were no patients with events of colitis with a fatal outcome.</p> <p>In the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), the reported outcomes for the 6 patients with events of colitis were as follows: 5 (1.9%) resolved, 1 (0.4%) not resolved. There were no patients with events of colitis with a fatal outcome.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), the outcomes for the 9 patients with events of colitis were as follows: 9 (2.5%) resolved. There were no fatal outcomes.</p> <p><u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251), the outcomes for the 3 patients with events of colitis were as follows: 3 (1.2%) resolved. There were no patients with events of colitis with a fatal outcome.</p>
--	--

Table SVII.3.1.2: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Colitis

	<p>Aggregate Review</p> <p>Review of the ASE dataset (n=7730) yielded 2 fatal cases of colitis. Review of the global safety database yielded the following: 2 fatal cases from ongoing interventional monotherapy clinical trials (corresponding to the 2 cases from ASE) and 11 fatal cases from the PM or non-interventional environment. Of the 11, there were 10 cases in which fatal colitis was the only Grade 5 event and 1 case that reported a Grade 5 colitis, as well as a Grade 5 MNP or other related fatal event(s).</p>																				
<p>Seriousness</p>	<p>Percent serious for identified risk of colitis - Pembrolizumab (All Serious AEs for 90 days after discontinuation were included.)</p> <p style="text-align: center;">Colitis – Serious (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Patients Treated with Pembrolizumab (Treatment Emergent)</p> <table border="1" data-bbox="488 625 1419 852"> <thead> <tr> <th rowspan="2"></th> <th colspan="2">Cumulative NSCLC 001/010 + MEL 001/002/006</th> </tr> <tr> <th>n</th> <th>(%)</th> </tr> </thead> <tbody> <tr> <td>Subjects in population</td> <td>2,799</td> <td></td> </tr> <tr> <td>with one or more adverse events</td> <td>32</td> <td>(1.1)</td> </tr> <tr> <td>Colitis</td> <td>30</td> <td>(1.1)</td> </tr> <tr> <td>Colitis microscopic</td> <td>1</td> <td>(0.0)</td> </tr> <tr> <td>Enterocolitis</td> <td>1</td> <td>(0.0)</td> </tr> </tbody> </table> <p>Every subject is counted a single time for each applicable row and column. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns meets the incidence criterion in the report title, after rounding. Include all treated subjects in KN001 Part B1, B2, B3, D, C, F1, F2, F3 and all subjects in KN002 treated with Pembrolizumab in the original phase, and all subjects in KN006 and KN010 treated with Pembrolizumab.</p> <p><u>Pembrolizumab IV</u></p> <p>Of the 13 patients with events of colitis in the NSCLC population in KN024 and KN042 (n=790), 8 (1.0%) experienced serious events of colitis.</p> <p>Of the 14 patients with events of colitis in the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), 5 (1.0%) experienced serious events of colitis.</p> <p>Of the 6 patients with events of colitis in the HL population in KN013, KN087, and KN204 (n=389), 2 (0.5%) experienced serious events of colitis.</p> <p>Of the 9 patients with events of colitis in the UC population in KN052 (n=370), 2 (0.5%) experienced serious events of colitis and of the 6 patients with events of colitis in the UC population in KN045 (n=266), 5 (1.9%) experienced serious events of colitis.</p> <p>Of the 4 patients with events of colitis in the HNSCC population (n=609), all 4 (0.7%) events of colitis experienced serious events of colitis.</p> <p>Of the 3 patients with events of colitis in the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), 2 (0.7%) experienced serious events of colitis. Of the 7 patients with events of colitis in the HNSCC population in KN048 pembrolizumab plus chemotherapy group (n=276), 2 (0.7%) experienced serious events of colitis.</p> <p>Of the 19 patients with events of colitis in the melanoma population in KN054 (n=509), 10 (2.0%) experienced serious events of colitis.</p> <p>Of the 9 patients with events of colitis in the NSCLC population in KN407 and KN021 (Cohort A) (n=303), 7 (2.3%) experienced serious events of colitis.</p> <p>Of the 11 patients with events of colitis in the RCC population in KN426 pembrolizumab plus axitinib (n=429), 6 (1.4%) experienced serious events of colitis.</p> <p>Of the 10 patients with events of colitis in the CRC population in KN177 (n=153), 5 (3.3%) experienced serious events of colitis.</p> <p>Of the 8 patients with events of colitis in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), 6 (1.6%) experienced serious events of colitis.</p>		Cumulative NSCLC 001/010 + MEL 001/002/006		n	(%)	Subjects in population	2,799		with one or more adverse events	32	(1.1)	Colitis	30	(1.1)	Colitis microscopic	1	(0.0)	Enterocolitis	1	(0.0)
	Cumulative NSCLC 001/010 + MEL 001/002/006																				
	n	(%)																			
Subjects in population	2,799																				
with one or more adverse events	32	(1.1)																			
Colitis	30	(1.1)																			
Colitis microscopic	1	(0.0)																			
Enterocolitis	1	(0.0)																			

Table SVII.3.1.2: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Colitis

	<p>Of the 11 patients with events of colitis in the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), 1 (0.2%) experienced a serious event of colitis.</p> <p>Of the 9 patients with events of colitis in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), 4 (1.1%) experienced serious events of colitis.</p> <p>Of the 26 patients with events of colitis in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), 10 (1.9%) experienced serious events of colitis.</p> <p>Of the 8 patients with events of colitis in the RCC population in KN564 pembrolizumab monotherapy (n=488), 6 (1.2%) experienced serious events of colitis.</p> <p>Of the 10 patients with events of colitis in the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), 2 (0.4%) experienced serious events of colitis.</p> <p>Of the 16 patients with events of colitis in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), 5 (1.6%) experienced serious events of colitis.</p> <p>Of the 13 patients with events of colitis in the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), 7 (0.9%) experienced serious events of colitis.</p> <p>Of the 17 patients with events of colitis in the melanoma population in KN716 pembrolizumab monotherapy (n=483), 7 (1.4%) experienced serious events of colitis.</p> <p>Of the 17 patients with events of colitis in gastric cancer population in KN811 pembrolizumab combination therapy (n=350), 6 (1.7%) experienced serious events of colitis.</p> <p>Of the 14 patients with events of colitis in the NSCLC population in KN091 pembrolizumab monotherapy (n=580), 4 (0.7%) experienced serious events of colitis.</p> <p>Of the 26 patients with events of colitis in the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), 21 (2.7%) experienced a serious event of colitis.</p> <p>Of the 9 patients with events of colitis in the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), 6 (1.1%) experienced serious events of colitis.</p> <p>Of the 5 patients with events of colitis in the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396), 2 (0.5%) experienced serious events of colitis.</p> <p>Of the 18 patients with events of colitis in the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), 8 (1.4%) experienced serious events of colitis.</p> <p>Of the 7 patients with events of colitis in the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382), 2 (0.5%) experienced serious events of colitis.</p> <p>Of the 14 patients with events of colitis in the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528), 4 (0.8%) experienced serious events of colitis.</p> <p>Of the 6 patients with events of colitis in the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), 4 (1.5%) experienced serious events of colitis.</p> <p>Of the 9 patients with events of colitis in the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), 6 (1.7%) experienced serious events of colitis.</p> <p><u>Pembrolizumab SC</u></p> <p>Of the 3 patients with events of colitis in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251), 2 (0.8%) experienced serious events of colitis</p>
--	---

Table SVII.3.1.2: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Colitis

Severity and Nature of the Risk	Identified treatment emergent adverse events (AE) of colitis (all events) in participants summarized by grade- pembrolizumab. (AEs for 30 days after discontinuation were included)		
	Colitis (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Patients Treated with Pembrolizumab (Treatment Emergent)		
		Cumulative NSCLC 001/010 + MEL 001/002/006	
		n	(%)
	Subjects in population	2,799	
	with one or more adverse events	48	(1.7)
	Colitis	45	(1.6)
	Grade 1	5	(0.2)
	Grade 2	9	(0.3)
	Grade 3	29	(1.0)
	Grade 4	2	(0.1)
	Colitis microscopic	2	(0.1)
	Grade 1	1	(0.0)
	Grade 2	1	(0.0)
	Enterocolitis	1	(0.0)
	Grade 3	1	(0.0)
	Autoimmune colitis	1	(0.0)
	Grade 2	1	(0.0)
	Every subject is counted a single time for each applicable specific adverse event. A subject with multiple adverse events within a system organ class is counted a single time for that system organ class. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the incidence specified in the report title, after rounding. Only the highest reported grade of a given adverse event is counted for the individual subject. Grades are based on NCI CTCAE version 4.0.		
	Pembrolizumab IV		
In the NSCLC population in KN024 and KN042 (n=790), there were 7 (0.9%) patients with Grade 3 events of colitis. There were no patients with Grade 4 or 5 events of colitis.			
In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), there were 5 (1.0%) patients with Grade 3 events of colitis. There were no patients with Grade 4 or 5 events of colitis.			
In the HL population in KN013, KN087, and KN204 (n=389), there were 3 (0.8%) patients with Grade 3 events of colitis. There were no patients with Grade 4 or 5 events of colitis.			
In the UC population in KN052 (n=370), there were 3 (0.8%) patients with Grade 3 events, 1 (0.3%) patient with a Grade 4 event, and no patients with Grade 5 events of colitis; in the UC population in KN045 (n=266), there were 3 (1.1%) patients with Grade 3 events and no patients with Grade 4 or 5 events of colitis.			
In the HNSCC population in KN040, KN012 and KN055 (n=609), there was 1 (0.2%) patient with a Grade 3 event of colitis. There were no patients with Grade 4 or 5 events of colitis.			
In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), there were 2 (0.7%) patients with Grade 3 events of colitis. There were no patients with Grade 4 or 5 events of colitis. In the pembrolizumab plus chemotherapy group (n=276), there were 2 (0.7%) patients with Grade 3 events of colitis. There were no patients with Grade 4 or 5 events of colitis.			
In the melanoma population in KN054 (n=509), there were 10 (2.0%) patients with Grade 3 events of colitis. There were no patients with Grade 4 or 5 events of colitis.			
In the NSCLC population in KN407 and KN021 (Cohort A) (n=303), there were 4 (1.3%) patients with Grade 3 events and 2 (0.7%) patients with Grade 4 events. There were no patients with Grade 5 events of colitis.			
In the RCC population in KN426 pembrolizumab plus axitinib (n=429), there were 8 (1.9%) patients with Grade 3 events of colitis. There were no patients with Grade 4 or 5 events of colitis.			

Table SVII.3.1.2: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Colitis

	<p>In the CRC population in KN177 (n=153) there were 3 (2.0%) patients with Grade 3 events and 2 (1.3%) patients with Grade 4 events of colitis. There were no patients with Grade 5 events of colitis reported.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), there were 4 (1.1%) patients with Grade 3 events of colitis. There were no patients with Grade 4 or 5 events of colitis.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), there were 2 (0.3%) patients with Grade 3 events of colitis. There were no patients with Grade 4 or 5 events of colitis.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352) there were 4 (1.1%) patients with Grade 3 events of colitis. There were no patients with Grade 4 or 5 events of colitis.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530) there were 8 (1.5%) patients with Grade 3, 1 (0.2%) patient with a Grade 4, and 1 (0.2%) patient with a Grade 5 events of colitis.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488) there were 5 (1.0%) patients with Grade 3 events of colitis. There were no patients with Grade 4 or 5 events of colitis.</p> <p>In the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), there was 1 (0.2%) patient with a Grade 3 and 1 (0.2%) patient with a Grade 4 event of colitis. There were no patients with Grade 5 events of colitis reported.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), there were 5 (1.6%) patients with Grade 3 and no patients with Grade 4 or 5 events of colitis reported.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) there were 5 (0.6%) patients with Grade 3 events, and 1 (0.1%) patient with a Grade 4 event of colitis. There were no patients with Grade 5 events of colitis reported.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), there were 8 (1.7%) patients with Grade 3 events of colitis. There were no patients with Grade 4 or 5 events of colitis.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), there were 9 (2.6%) patients with Grade 3 events of colitis. There were no patients with Grade 4 and 5 events of colitis.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580) there were 4 (0.7%) patients with Grade 3 events of colitis. There were no patients with Grade 4 or 5 events of colitis.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), there were 18 (2.3%) patients with a Grade 3, and 1 (0.1%) patient with a Grade 4 event of colitis reported. There were no patients with a Grade 5 event of colitis reported.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), there were 4 (0.8%) patients with Grade 3 events and 1 (0.2%) patient with a Grade 4 event of colitis. There were no patients with Grade 5 events of colitis.</p> <p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396) there were 3 (0.8%) patients with Grade 3 events of colitis, there were no patients with Grade 4 or 5 events of colitis reported.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), there were 9 (1.6%) patients with Grade 3 events of colitis. There were no patients with Grade 4 or 5 events of colitis.</p> <p>In the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382) there was 1 (0.3%) patient with Grade 3 event of colitis. There were no patients with Grade 4 or 5 events of colitis.</p> <p>In the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528), there were 2 (0.4%) patients with Grade 3 events of colitis, and 2 (0.4%) patients with Grade 4 colitis. There were no Grade 5 events of colitis.</p> <p>In the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), there were no patients with Grade 3, 4 or 5 events of colitis.</p>
--	--

Table SVII.3.1.2: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Colitis

	<p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), there were 7 (1.9%) patients with Grade 3 and 1 (0.3%) patient with Grade 4 events of colitis. There were no patients with Grade 5 events of colitis.</p> <p><u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251) there were 2 (0.8%) patients with Grade 3 events of colitis. There were no patients with Grade 4 or 5 events of colitis reported.</p> <p>In the global safety database through 31-DEC-2017, (pembrolizumab monotherapy trials only), there were 2 Grade 5 clinical trial reports of colitis.</p> <p>Review of pembrolizumab clinical trial data from ongoing studies and postmarketing data regarding immune-mediated colitis, including fatal cases, is consistent with the risk as characterized in this RMP.</p>
<u>Risk Factors and Risk Groups:</u>	No specific risk factors for colitis and diarrhea associated with pembrolizumab were identified.
<u>Preventability:</u>	Although the development of colitis cannot be completely prevented; patients should be monitored for signs and symptoms of colitis and if they develop, other causes excluded. Withholding/discontinuation of pembrolizumab and appropriate medical intervention including corticosteroids following the onset of colitis may result in recovery.
<u>Impact on the Risk-Benefit Balance of the Product:</u>	Severe colitis may be life-threatening or fatal in individual patients. However, given the fatal outcome of untreated cancer, the risk of colitis which generally can be managed, is outweighed by the potential benefit of pembrolizumab treatment.
<u>Public Health Impact:</u>	This risk has minimal public health impact outside its effect on individual patients.

Table SVII.3.1.3: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Hepatitis

<u>Important Identified Risk:</u>	Immune-mediated Hepatitis
<u>Potential Mechanisms:</u>	Pembrolizumab is a potent and highly-selective humanized monoclonal antibody (mAb) of the Immunoglobulin G4 (IgG4)/kappa isotype directed to the programmed cell death-1 (PD-1) receptor and designed to directly block the interaction between the receptor and its ligands, PD-L1 and PD-L2. The PD-1 pathway represents a major immune control switch which may be engaged by tumor cells to overcome active T-cell immune surveillance. A potential mechanism of action is T-cell activation in proximity to normal tissue resulting inflammation and injury to normal liver tissue.
<u>Evidence Source(s) and Strength of Evidence:</u>	<p>Review of pembrolizumab clinical trial data, post-marketing experience and literature regarding hepatitis represent sufficient evidence of a causal association with pembrolizumab exposure.</p> <p>CTD 2.7.4</p> <p>Please reference evidence source cited in Table SVII.3.1.1; Details of Important Identified Risk: Immune-Mediated Pneumonitis</p>

Table SVII.3.1.3: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Hepatitis

Characterisation of the Risk: Immune-mediated Hepatitis							
Frequency with 95%CI	95% Confidence Interval for the Overall Incidence (%) of Hepatitis KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab						
		<table border="1"> <thead> <tr> <th colspan="2">Cumulative MEL 001/002/006 + NSCLC 001/010</th> </tr> <tr> <th>n (%)</th> <th>95% CI</th> </tr> </thead> <tbody> <tr> <td>Hepatitis</td> <td>18 (0.6) (0.4,1.0)</td> </tr> </tbody> </table>	Cumulative MEL 001/002/006 + NSCLC 001/010		n (%)	95% CI	Hepatitis
Cumulative MEL 001/002/006 + NSCLC 001/010							
n (%)	95% CI						
Hepatitis	18 (0.6) (0.4,1.0)						
<p><u>Pembrolizumab IV</u></p> <p>The overall number and proportion of patients with hepatitis reported in the NSCLC population in KN024 and KN042 (n=790) was 10 (1.3%) (95% CI-. (0.6, 2.3), which is generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hepatitis in the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488) was 5 (1%) (95% CI- 0.3, 2.4), which is generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hepatitis in the HL population in KN013, KN087, and KN204 (n=389) was 2 (0.5%) (95%CI-0.1, 1.8), which is generally similar to the overall proportion of patients with the event of hepatitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hepatitis in the UC population in KN052 (n=370) was 4 (1.1%) (95%CI-0.3,2.7), which is generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset for melanoma and NSCLC.</p> <p>There were no patients with events of hepatitis reported in the UC population in KN045.</p> <p>The overall number and proportion of patients with hepatitis in the HNSCC population in KN040, KN012 and KN055 (n=609) was 6 (1.0%) (95% CI- 0.4,2.1), which is generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hepatitis in the HNSCC population in KN048 pembrolizumab monotherapy (n=300) and pembrolizumab plus chemotherapy (n=276) groups were 2 (0.7%) (95% CI- 0.1,2.4) and 0 (0) (95% CI-0, 1.3) respectively, which is generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hepatitis in the melanoma population in KN054 (n=509) was 9 (1.8%) (95% CI-0.8, 3.3). The proportion of patients with events of hepatitis in KN054 although somewhat more common than in the reference safety dataset were generally consistent in nature with the proportion of events of hepatitis previously reported for pembrolizumab.</p> <p>The overall number and proportion of patients with hepatitis in the NSCLC population in KN407 and KN021 (Cohort A) (n=303), was 5 (1.7%) (95% CI-0.5, 3.8), which is generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hepatitis in the RCC population in KN426 pembrolizumab plus axitinib (n=429) was 12 (2.8%) (95% CI-1.5,4.8), which is higher than the proportion of patients with the event of hepatitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hepatitis in the CRC population in KN177 (n=153) was 4 (2.6%) (95% CI-(0.7,6.6)) which is generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hepatitis in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370) was 5 (1.4%) (95% CI-0.4,3.1), which is</p>							

Table SVII.3.1.3: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Hepatitis

<p>generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hepatitis in the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596) was 5 (0.8%) (95% CI-0.3,1.9), which is generally similar to the proportion of patients with the event of hepatitis in the reference data set for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hepatitis in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352) was 7 (2.0%) (95% CI-0.8,4.1), which is generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hepatitis in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530) was 8 (1.5%) (95% CI-0.7, 3.0) which is generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hepatitis in the RCC population in KN564 pembrolizumab monotherapy (n=488) was 5 (1%) (95% CI-0.3,2.4), which is generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hepatitis in the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475) was 4 (0.8%) (95% CI-0.2,2.1) which is generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hepatitis in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307) was 5 (1.6%) (95% CI-0.5, 3.8), which is generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hepatitis in the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) was 11 (1.4%) (95% CI-0.7,2.5), which is generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hepatitis in the melanoma population in KN716 pembrolizumab monotherapy (n=483) was 10 (2.1%) (95% CI-1.0, 3.8), which is generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hepatitis in the gastric cancer population in KN811 pembrolizumab combination therapy (n=350) was 2 (0.6%) (95% CI-0.1, 2.1), which is the same as the proportion of patients with the event of hepatitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hepatitis in the NSCLC population in KN091 pembrolizumab monotherapy (n=580) was 10 (1.7%) (95% CI-0.8,3.1), which is generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset</p> <p>The overall number and proportion of patients with hepatitis in the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785) was 9 (1.1%) (95% CI-0.5, 2.2), which is generally similar to the proportion of patients with the events of hepatitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hepatitis in the BTC population in KN966 pembrolizumab plus chemotherapy (n=529) was 9 (1.7%) (95% CI-0.8, 3.2), which is generally similar to the proportion of patients with the events of hepatitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hepatitis in the NSCLC population in KN671 pembrolizumab in combination with chemotherapy neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396) was 3 (0.8%) (95% CI-0.2, 2.2), which is generally similar to the overall proportion of patients with events of hepatitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hepatitis in the UC population in KN869 and KNA39 pembrolizumab +EV combination therapy (n=564) was 16 (2.8%) (95% CI-1.6%, 4.6%), which is generally similar to the proportion of patients with the events of hepatitis in the reference safety dataset.</p>
--

Table SVII.3.1.3: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Hepatitis

	<p>There were no patients with events of hepatitis reported in the endometrial carcinoma patient population in KN868 pembrolizumab plus chemotherapy (n=382).</p> <p>The overall number and proportion of patients with hepatitis in the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528) was 1 (0.2%) (95% CI- 0.0, 1.1), which is generally similar to the overall frequency in the reference safety dataset.</p> <p>There were no patients with events of hepatitis reported in the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260).</p> <p>The overall number and proportion of patients with hepatitis in the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361) were 8 (2.2%) (95% CI- 1.0,4.3), which is generally similar to the proportion of patients with the event of hepatitis in the reference safety dataset.</p> <p><u>Pembrolizumab SC</u></p> <p>The overall number and proportion of patients with hepatitis in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251) was 1 (0.4%) (95% CI-0.0, 2.2), which is generally similar to the overall proportion of patients with events of hepatitis in the reference safety dataset.</p>																															
<p>Outcomes</p>	<p>Percent overall, fatal, not resolved, resolved, and unknown for identified risk of hepatitis - pembrolizumab. (Table includes events for 30 days after discontinuation.)</p> <p style="text-align: center;">Hepatitis (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment Emergent)</p> <table border="1" data-bbox="493 982 1417 1268"> <thead> <tr> <th colspan="2"></th> <th colspan="2">Cumulative NSCLC 001/010 + MEL 001/002/006</th> </tr> <tr> <th></th> <th>Outcome</th> <th>n</th> <th>(%)</th> </tr> </thead> <tbody> <tr> <td>Subjects in population</td> <td></td> <td>2,799</td> <td></td> </tr> <tr> <td rowspan="6">With one or more adverse events</td> <td>Overall</td> <td>18</td> <td>(0.6)</td> </tr> <tr> <td>Fatal</td> <td>0</td> <td>(0.0)</td> </tr> <tr> <td>Not Resolved</td> <td>2</td> <td>(0.1)</td> </tr> <tr> <td>Resolved</td> <td>14</td> <td>(0.5)</td> </tr> <tr> <td>Resolving</td> <td>2</td> <td>(0.1)</td> </tr> <tr> <td>Unknown</td> <td>0</td> <td>(0.0)</td> </tr> </tbody> </table> <p>Every subject is counted once on each applicable row. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the percent incidence specified in the report title, after rounding. Outcome: Resolved = RECOVERED/RESOLVED, Not resolved = NOT RECOVERED/NOT RESOLVED.</p> <p><u>Pembrolizumab IV</u></p> <p>In the NSCLC population in KN024 and KN042, the outcomes of the 10 patients with events of hepatitis were as follows: 9 (1.1%) resolved, 1 (0.1%) resolving. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), the outcomes of the 5 patients with events were as follows: 3 (0.6%) resolved, 1 (0.2%) not resolved, 1 (0.2%) unknown. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the HL population in KN013, KN087, and KN204 (n=389), the outcome of the 2 patients with events of hepatitis were as follows: 1 resolved (0.3%) and 1 not resolved (0.3%). There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the UC population in KN052 (n=370), the outcomes for the 4 patients with events of hepatitis were as follows: 4 (1.1%) resolved. There were no patients with events of hepatitis with a fatal outcome.</p>			Cumulative NSCLC 001/010 + MEL 001/002/006			Outcome	n	(%)	Subjects in population		2,799		With one or more adverse events	Overall	18	(0.6)	Fatal	0	(0.0)	Not Resolved	2	(0.1)	Resolved	14	(0.5)	Resolving	2	(0.1)	Unknown	0	(0.0)
		Cumulative NSCLC 001/010 + MEL 001/002/006																														
	Outcome	n	(%)																													
Subjects in population		2,799																														
With one or more adverse events	Overall	18	(0.6)																													
	Fatal	0	(0.0)																													
	Not Resolved	2	(0.1)																													
	Resolved	14	(0.5)																													
	Resolving	2	(0.1)																													
	Unknown	0	(0.0)																													

Table SVII.3.1.3: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Hepatitis

<p>In the HNSCC population in KN040, KN012 and KN055 (n=609), the outcomes for the 6 patients with events of hepatitis were as follows: 4 (0.7%) resolved, 1 (0.2%) resolving and 1 (0.2%) not resolved. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), the outcomes for the 2 patients with events of hepatitis were as follows: 1 (0.3%) resolved and 1 (0.3%) resolving. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the HNSCC population in KN048 pembrolizumab plus chemotherapy group (n=276), there were no patients with events of hepatitis.</p> <p>In the melanoma population in KN054 (n=509), the outcomes for the 9 patients with events of hepatitis were as follows: 6 (1.2%) not resolved and 3 (0.6%) resolved. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the NSCLC population in KN407 and KN021 (Cohort A) (n=303), the outcomes of the 5 patients with events of hepatitis were as follows: 3 (1.0%) resolved, 1 (0.3%) not resolved and 1 (0.3%) resolving. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the RCC population in KN426 pembrolizumab plus axitinib (n=429), the outcomes for the 12 patients with events of hepatitis were as follows: 9 (2.1%) resolved, 2 (0.5%) resolving and 1 (0.2%) resolved with sequelae. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the CRC population in KN177 (n=153), the outcomes for the 4 patients with events of hepatitis were as follows: 4 (2.6%) resolved. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), the outcomes for the 5 patients with events of hepatitis were as follows: 2 (0.5%) resolved and 3 (0.8%) not resolved. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), the outcomes for the 5 patients with events of hepatitis were as follows: 2 (0.3%) resolved, 2 (0.3%) not resolved, and 1 (0.2%) resolving. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), the outcomes for the 7 patients with events of hepatitis were as follows: 3 resolved (0.9%), 2 (0.6%) resolving, 1 (0.3%) not resolved. There was 1 (0.3%) patient with an event of hepatitis that resulted in a fatal outcome.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), the outcomes for the 8 patients with events of hepatitis were as follows: 4 (0.8%) resolved, 3 (0.6%) not resolved, and 1 (0.2%) resolved with sequelae. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488), the outcomes for the 5 patients with events of hepatitis were as follows: 5 (1%) resolved. There were no events of hepatitis with a fatal outcome.</p> <p>In the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), the outcomes for the 4 (0.8%) patients with events of hepatitis were as follows: 3 (0.6%) resolved and 1 (0.2%) not resolved. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), the outcomes for the 5 patients with events of hepatitis were as follows: 1 (0.3%) resolving, and 4 (1.3%) resolved. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), the outcomes for the 11 patients with events of hepatitis were as follows: 8 (1.0%) resolved, and 3 (0.4%) not resolved. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), the outcomes for the 10 patients with events of hepatitis were as follows: 6 (1.2%) resolved, 2 (0.4%) resolving, 1 (0.2%) not resolved, and 1 (0.2%) resolved with sequelae. There were no patients with events of hepatitis with a fatal outcome.</p>

Table SVII.3.1.3: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Hepatitis

<p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), the outcome for the 2 events of hepatitis was as follows: 1 (0.3%) resolved. There was 1 (0.3%) patient with an event of hepatitis that resulted in a fatal outcome.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580), the outcomes for the 10 patients with events of hepatitis were as follows: 7 (1.2%) resolved, 2 (0.3%) not resolved, and 1 (0.2%) resolved with sequelae. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), the outcomes for the 9 patients with events of hepatitis were as follows: 6 (0.8%) resolved, 1 (0.1%) not resolved, 1 (0.1%) resolving, and 1 (0.1%) unknown. There were no patients with an event of hepatitis with a fatal outcome.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), the worst reported outcomes for the 9 patients with events of hepatitis were as follows: 8 (1.5%) resolved and 1 (0.2%) not resolved. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy group (n=396), the outcomes for the 3 patients with events of hepatitis were as follows: 2 (0.5%) resolved, and 1 (0.3%) resolved with sequelae. There were no patients with events of hepatitis with a fatal outcome.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), the outcomes for the 16 patients with events of hepatitis were as follows: 14 (2.5%) resolved, 1 (0.2%) not resolved, and 1 (0.2%) resolving. There were no patients with an event of hepatitis with a fatal outcome.</p> <p>In the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528), the outcome for the 1 (0.2%) event of hepatitis was resolved. There were no events of hepatitis with a fatal outcome.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), the outcomes for the 8 patients with events of hepatitis were as follows: 7 (1.9%) resolved and 1 (0.3%) resolved with sequelae. There were no patients with events of hepatitis with a fatal outcome.</p> <p><u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251), the outcome for the 1 (0.4) event of hepatitis was resolving. There were no patients with events of hepatitis with a fatal outcome.</p> <p><u>Aggregate Review</u></p> <p>Review of the ASE dataset (n=7730) yielded no fatal cases of hepatitis. Review of the global safety database yielded the following: 4 fatal cases from ongoing interventional monotherapy clinical trials and 9 fatal cases from the PM or non-interventional environment. Of the 9, all were cases in which fatal hepatitis was the only Grade 5 event.</p>
--

Table SVII.3.1.3: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Hepatitis

Seriousness	Percent serious for identified risk of hepatitis - Pembrolizumab. (Serious AEs for 90 days after discontinuation were included.)		
	Hepatitis - Serious (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment Emergent)		
	Cumulative NSCLC 001/010 + MEL 001/002/006		
		n	(%)
	Subjects in population	2,799	
	with one or more adverse events	11	(0.4)
	Autoimmune hepatitis	8	(0.3)
	Drug-induced liver injury	2	(0.1)
	Hepatitis	2	(0.1)
	<p>Every subject is counted a single time for each applicable row and column. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns meets the incidence criterion in the report title, after rounding. Include all treated subjects in KN001 Part B1, B2, B3, D, C, F1, F2, F3 and all subjects in KN002 treated with Pembrolizumab in the original phase, and all subjects in KN006 and KN010 treated with Pembrolizumab.</p>		

Pembrolizumab IV

Of the 10 patients with events of hepatitis in the NSCLC population in KN024 and KN042, 6 (0.8%) experienced serious events of hepatitis.

Of the 5 patients with events of hepatitis in the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), 3 (0.6%) experienced serious events of hepatitis.

Of the 2 patients with events of hepatitis in the HL population in KN013, KN087, and KN204 (n=389), 2 (0.5%) experienced serious events of hepatitis.

Of the 4 patients with events of hepatitis in the UC population in KN052 (n=370), 3 (0.8%) experienced serious events of hepatitis.

Of the 6 patients with events of hepatitis in the HNSCC population in KN040, KN012 and KN055 (n=609), 2 (0.3%) experienced serious events of hepatitis.

Of the 2 patients with events of hepatitis in the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), 2 (0.7%) experienced serious events of hepatitis. There were no patients with serious events of hepatitis reported in the pembrolizumab plus chemotherapy group (n=276).

Of the 9 patients with events of hepatitis in the melanoma population in KN054 (n=509), 4 (0.8%) experienced serious events of hepatitis.

Of the 5 patients with events of hepatitis in the NSCLC population in KN407 and KN021 (Cohort A) (n=303), 2 (0.7%) experienced serious. events of hepatitis

Of the 12 patients with events of hepatitis in the RCC population in KN426 pembrolizumab plus axitinib (n=429), 7 (1.6%) experienced serious events of hepatitis.

Of the 4 patients with events of hepatitis in the CRC population in KN177 (n=153), 4 (2.6%) experienced serious events of hepatitis.

Of the 5 patients with events of hepatitis in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), 3 (0.8%) experienced serious events of hepatitis.

Of the 5 patients with events of hepatitis in TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), 4 (0.7%) experienced serious events of hepatitis.

Of the 7 patients with events of hepatitis in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), 6 (1.7%) experienced serious events of hepatitis.

Of the 8 patients with events of hepatitis in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), 4 (0.8%) experienced serious events of hepatitis.

Table SVII.3.1.3: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Hepatitis

	<p>Of the 5 patients with events of hepatitis in the RCC population in KN564 pembrolizumab monotherapy (n=488), 3 (0.6%) experienced serious events of hepatitis.</p> <p>Of the 4 patients with events of hepatitis in the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), 1 (0.2%) experienced a serious event of hepatitis.</p> <p>Of the 5 patients with events of hepatitis in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), 3 (1.0%) experienced serious events of hepatitis.</p> <p>Of the 11 patients with events of hepatitis in the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), 7 (0.9%) experienced serious events of hepatitis.</p> <p>Of the 10 patients with events of hepatitis in the melanoma population in KN716 pembrolizumab monotherapy (n=483), 2 (0.4%) experienced serious events of hepatitis.</p> <p>Of the 2 patients with events of hepatitis in gastric cancer population in KN811 pembrolizumab combination therapy (n=350), 2 (0.6%) experienced serious events of hepatitis.</p> <p>Of the 10 patients with events of hepatitis in the NSCLC population in KN091 pembrolizumab monotherapy (n=580), 9 (1.6%) experienced serious events of hepatitis.</p> <p>Of the 9 patients with events of hepatitis in the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), 4 (0.5%) experienced serious events of hepatitis.</p> <p>Of the 9 patients with events of hepatitis in the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), 3 (0.6%) were serious.</p> <p>Of the 3 patients with events of hepatitis in the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396), 2 (0.5%) experienced serious events of hepatitis.</p> <p>Of the 16 patients with events of hepatitis in the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), 7 (1.2%) experienced serious events of hepatitis.</p> <p>There were no serious events of hepatitis reported in the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528).</p> <p>Of the 8 patients with events of hepatitis in the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), 7 (1.9%) experienced serious events of hepatitis.</p> <p><u>Pembrolizumab SC</u></p> <p>There were no serious events of hepatitis reported in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemoradiotherapy (n=251).</p>
--	--

Table SVII.3.1.3: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Hepatitis

Severity and Nature of the Risk	Identified treatment emergent adverse events (AE) of hepatitis (all events) in participants summarized by grade – Pembrolizumab. (AEs for 30 days after discontinuation were included)		
	Hepatitis (AEOSI)		
	KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment Emergent)		
		Cumulative NSCLC 001/010 + MEL 001/002/006	
		n	(%)
	Subjects in population	2,799	
	with one or more adverse events	18	(0.6)
	Autoimmune hepatitis	12	(0.4)
	Grade 2	4	(0.1)
	Grade 3	7	(0.3)
	Grade 4	1	(0.0)
	Drug-induced liver injury	2	(0.1)
	Grade 3	1	(0.0)
	Grade 4	1	(0.0)
	Hepatitis	5	(0.2)
Grade 1	1	(0.0)	
Grade 3	4	(0.1)	
Every subject is counted a single time for each applicable specific adverse event. A subject with multiple adverse events within a system organ class is counted a single time for that system organ class. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the incidence specified in the report title, after rounding. Only the highest reported grade of a given adverse event is counted for the individual subject. Grades are based on NCI CTCAE version 4.0.			
<u>Pembrolizumab IV</u>			
In the NSCLC population in KN024 and KN042, (n=790), there were 6 (0.8%) patients with Grade 3 events, 2 (0.3%) patients with Grade 4 events, and no patients with Grade 5 events of hepatitis.			
In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), there were 3 (0.6%) patients with Grade 3 events, 1 (0.2%) patient with a Grade 4 event, and no patients with Grade 5 events of hepatitis.			
In the HL population in KN013, KN087, and KN204 (n=389), there were 2 (0.5%) patients with Grade 3 events and no patients with Grade 4 or 5 events of hepatitis reported.			
In the UC population in KN052 (n=370), there were 4 (1.1%) patients with Grade 3 events and there were no patients with Grade 4 or 5 events of hepatitis.			
In the HNSCC population in KN040, KN012 and KN055 (n=609), there were 3 (0.5%) patients with Grade 3 events and 1 (0.2%) patient with a Grade 4 event of hepatitis.			
In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), there were 2 (0.7%) patients with Grade 4 events; there were no patients with Grade 5 events of hepatitis; in the pembrolizumab plus chemotherapy group (n=276), there were no patients with events of hepatitis.			
In the melanoma population in KN054 (n=509), there were 6 (1.2%) patients with Grade 3 events and 1 (0.2%) patient with a Grade 4 event of hepatitis. There were no patients with Grade 5 events of hepatitis.			
In the NSCLC population in KN407 and KN021 (Cohort A) (n=303), there were 4 (1.3%) patients with Grade 3 events, 1 (0.3%) patient with a Grade 4 event and no patients with Grade 5 events of hepatitis.			
In the RCC population in KN426 pembrolizumab plus axitinib (n=429), there were 7 (1.6%) patients with Grade 3 events and 3 (0.7%) patients with Grade 4 events of hepatitis. There were no patients with Grade 5 events of hepatitis.			
In the CRC population in KN177 (n=153) there were 4 (2.6%) patients with Grade 3 events. There were no patients with Grade 4 or 5 events of hepatitis.			

Table SVII.3.1.3: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Hepatitis

	<p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), there were 5 (1.4%) patients with Grade 3 events of hepatitis. There were no patients with Grade 4 or 5 events of hepatitis.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), there were 3 (0.5%) patients with Grade 3 and 1 (0.2%) patient with a Grade 4 event of hepatitis. There were no patients with Grade 5 events of hepatitis.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352) there were 3 (0.9%) patients with Grade 3, 1 (0.3%) patient with a Grade 4, and 1 (0.3%) patient with a Grade 5 events of hepatitis.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), there were 7 (1.3%) patients with Grade 3 and no patients with Grade 4 or 5 events of hepatitis.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488) there were 4 (0.8%) patients with Grade 3 events of hepatitis. There were no patients with Grade 4 or 5 events of hepatitis.</p> <p>In the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), there were 2 (0.4%) patients with Grade 3 events of hepatitis. There were no patients with Grade 4 or 5 events of hepatitis reported.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), there were 3 (1.0%) patients with Grade 3 and 1 (0.3%) patient with a Grade 4 events of hepatitis reported. There were no Grade 5 events of hepatitis.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) there were 6 (0.8%) patients with Grade 3 events, and 3 (0.4%) patients with Grade 4 events of hepatitis. There were no patients with Grade 5 events of hepatitis reported.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), there were 8 (1.7%) patients with Grade 3 events of hepatitis. There were no patients with Grade 4 or 5 events of hepatitis.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), there was 1 (0.3%) patient with a Grade 4 event and 1 (0.3%) patient with a Grade 5 event of hepatitis.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580) there were 5 (0.9%) patients with Grade 3 and 4 (0.7%) patients with Grade 4 events of hepatitis. There were no patients with Grade 5 events of hepatitis.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), there were 3 (0.4%) patients with a Grade 3 event of hepatitis reported. There were no patients with Grade 4 or 5 events of hepatitis reported.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), there were 5 (0.9%) patients with Grade 3 events of hepatitis reported. There were no patients with Grade 4 or 5 events of hepatitis.</p> <p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396) there was 1 (0.3%) patient with a Grade 4 event of hepatitis, there were no patients with Grade 3 or 5 events of hepatitis reported.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), there were 9 (1.6%) patients with Grade 3 events, and 1 (0.2%) with a Grade 4 event. There were no patients with Grade 5 events of hepatitis.</p> <p>In the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528), there were no Grade 3, 4 or 5 events of hepatitis.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), there were 5 (1.4%) patients with Grade 3 events and 2 (0.6%) with Grade 4 events; there were no patients with Grade 5 events of hepatitis.</p>
--	---

Table SVII.3.1.3: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Hepatitis

	<p><u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy, there were no patients with Grade 3, 4, or 5 events of hepatitis reported. In the global safety database through 31-DEC-2017, (pembrolizumab monotherapy trials only), there were 4 Grade 5 clinical trial reports of hepatitis.</p> <p>Review of pembrolizumab clinical trial data from ongoing studies and postmarketing data regarding immune-mediated hepatitis, including fatal cases, is consistent with the risk as characterized in this RMP.</p>
<u>Risk Factors and Risk Groups:</u>	Patients with moderate to severe liver dysfunction were excluded from clinical trials. No analysis of specific risk factors for immune-mediated hepatitis associated with pembrolizumab has been undertaken.
<u>Preventability:</u>	Although the development of hepatitis cannot be completely prevented; patients should be monitored for signs and symptoms of hepatitis and if they develop, other causes excluded. Withholding/discontinuation of pembrolizumab and appropriate medical intervention including corticosteroids following the onset of hepatitis may result in recovery.
<u>Impact on the Risk-Benefit Balance of the Product:</u>	Hepatic failure may be life-threatening or fatal in individual patients; however, the event can be managed with cessation of pembrolizumab and immunosuppression with corticosteroids. Given the fatal outcome of untreated cancer, the risk of immune-mediated hepatitis is outweighed by the potential benefit.
<u>Public Health Impact:</u>	This risk has minimal public health impact outside its effect on individual patients.

Table SVII.3.1.4: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Nephritis

<u>Important Identified Risk:</u>	Immune-mediated Nephritis								
<u>Potential Mechanisms:</u>	Pembrolizumab is a potent and highly-selective humanized monoclonal antibody (mAb) of the Immunoglobulin G4 (IgG4)/kappa isotype directed to the programmed cell death-1 (PD-1) receptor and designed to directly block the interaction between the receptor and its ligands, PD-L1 and PD-L2. The PD-1 pathway represents a major immune control switch which may be engaged by tumor cells to overcome active T-cell immune surveillance. T-cell activation in proximity to normal tissue may lead to inflammation and injury to normal tissue.								
<u>Evidence Source(s) and Strength of Evidence:</u>	Review of pembrolizumab clinical trial data, post-marketing experience and literature regarding nephritis represent sufficient evidence of a causal association with pembrolizumab exposure. CTD 2.7.4 Please reference evidence source cited in Table SVII.3.1.1; Details of Important Identified Risk: Immune-Mediated Pneumonitis								
<u>Characterisation of the Risk:</u> Immune-mediated Nephritis									
<u>Frequency with 95%CI</u>	<p>95% Confidence Interval for the Overall Incidence (%) of Nephritis KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab</p> <table border="1"> <thead> <tr> <th rowspan="2"></th> <th colspan="2">Cumulative MEL 001/002/006 + NSCLC 001/010</th> </tr> <tr> <th>n (%)</th> <th>95% CI</th> </tr> </thead> <tbody> <tr> <td>Nephritis</td> <td>9 (0.3)</td> <td>(0.1,0.6)</td> </tr> </tbody> </table> <p><u>Pembrolizumab IV</u></p> <p>The overall number and proportion of patients with nephritis in the NSCLC population in KN024 and KN042 (n=790) was 4 (0.5%) (95%CI- 0.1, 1.3), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset for melanoma and NSCLC.</p>		Cumulative MEL 001/002/006 + NSCLC 001/010		n (%)	95% CI	Nephritis	9 (0.3)	(0.1,0.6)
	Cumulative MEL 001/002/006 + NSCLC 001/010								
	n (%)	95% CI							
Nephritis	9 (0.3)	(0.1,0.6)							

Table SVII.3.1.4: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Nephritis

	<p>The overall number and proportion of patients with nephritis in the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488) was 7 (1.4%) (95% CI- 0.6, 2.9), which is marginally higher than the proportion of patients with the event of nephritis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with nephritis in the HL population in KN013, KN087, and KN204 (n=389) was 2 (0.5%) (95%CI-0.1, 1.8), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with nephritis in the UC population in KN052 (n=370) was 1 (0.3%) (95%CI-0,1.5) and in KN045 (n=266) was 2 (0.8%) (95%CI-0.1,2.7), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset for melanoma and NSCLC.</p> <p>There were no patients with events of nephritis reported in the HNSCC population in KN040, KN012 and KN055.</p> <p>The overall number and proportion of patients with nephritis in the HNSCC population in KN048 pembrolizumab monotherapy (n=300) and pembrolizumab plus chemotherapy (n=276) groups were 2 (0.7%) (95%CI-0.1, 2.4) and 2 (0.7%) (95%CI-0.1, 2.6) respectively, which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with nephritis in the melanoma population in KN054 (n=509) was 2 (0.4%) (95% CI-0,1.4), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with nephritis in the NSCLC population in KN407 and KN021 (Cohort A) (n=303), was 2 (0.7%) (95%CI-0.1, 2.4), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with nephritis in the RCC population in KN426 pembrolizumab plus axitinib (n=429) was 6 (1.4%) (95% CI-0.5,3), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with nephritis in the CRC population in KN177 (n=153) was 1 (0.7%) (95% CI-0,3.6) which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with nephritis in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370) was 1 (0.3%) (95% CI-0,1.5), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with nephritis in the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596) was 4 (0.7%) (95% CI-0.2,1.7), which is generally similar to the overall frequency in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with nephritis in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352) was 6 (1.7%) (95% CI-0.6, 3.7), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset.</p> <p>The overall number and proportion of patients with nephritis in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530) was 5 (0.9%) (95% CI-0.3, 2.2), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset.</p> <p>The overall number and proportion of patients with nephritis in the RCC population in KN564 pembrolizumab monotherapy (n=488) was 3 (0.6%) (95% CI-0.1,1.8), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with nephritis in the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475) was 2 (0.4%) (95% CI-0.1,1.5) which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset.</p>
--	---

Table SVII.3.1.4: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Nephritis

	<p>The overall number and proportion of patients with nephritis in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307) was 1 (0.3%) (95% CI-0.0, 1.8), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset.</p> <p>The overall number and proportion of patients with nephritis in the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) was 7 (0.9%) (95% CI-0.4,1.8), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset.</p> <p>The overall number and proportion of patients with nephritis in the melanoma population in KN716 pembrolizumab monotherapy (n=483) was 7 (1.4%) (95% CI-0.6, 3.0), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with nephritis in the gastric cancer population in KN811 pembrolizumab combination therapy (n=350) was 4 (1.1%) (95% CI-0.3, 2.9), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset.</p> <p>The overall number and proportion of patients with nephritis in the NSCLC population in KN091 pembrolizumab monotherapy (n=580) was 4 (0.7%) (95% CI-0.2, 1.8), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset.</p> <p>The overall number and proportion of patients with nephritis in the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785) was 4 (0.5%) (95% CI-0.1, 1.3), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset.</p> <p>The overall number and proportion of patients with nephritis in the BTC population in KN966 pembrolizumab plus chemotherapy (n=529) was 2 (0.4%) (95% CI-0.0, 1.4), which is generally similar to the proportion of patients with the events of nephritis in the reference safety dataset.</p> <p>There were no events of nephritis reported in the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396).</p> <p>The overall number and proportion of patients with nephritis in the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564) was 6 (1.1%) (95% CI-0.4 %, 2.3%), which is generally similar to the proportion of patients with the events of nephritis in the reference safety dataset.</p> <p>The overall number and proportion of patients with nephritis in the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382) was 2 (0.5%) (95% CI-0.1, 1.9), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset.</p> <p>The overall number and proportion of patients with nephritis in the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528) was 2 (0.4%) (95% CI-0.0, 1.4), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset.</p> <p>The overall number and proportion of patients with nephritis in the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260) was 2 (0.8%) (95% CI-0.1, 2.8), which is generally similar to the proportion of patients with the events of nephritis in the reference safety dataset.</p> <p>The overall number and proportion of patients with nephritis in the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361) were 3 (0.8%) (95%CI-0.2, 2.4), which is generally similar to the proportion of patients with the event of nephritis in the reference safety dataset.</p> <p><u>Pembrolizumab SC</u></p> <p>There were no events of nephritis reported in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251).</p>
--	---

Table SVII.3.1.4: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Nephritis

Outcomes	Percent overall, fatal, not resolved, resolved, and unknown for identified risk of nephritis as assessed by the Sponsor - pembrolizumab. (Table includes events for 30 days after discontinuation.)		
	Nephritis (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment Emergent)		
			Cumulative NSCLC 001/010 + MEL 001/002/006
		Outcome	n (%)
	Subjects in population		2,799
	With one or more adverse events	Overall	9 (0.3)
		Fatal	0 (0.0)
		Not Resolved	3 (0.1)
		Resolved	5 (0.2)
		Resolving	1 (0.0)
	Unknown	0 (0.0)	
Outcome: Resolved = RECOVERED/RESOLVED, Not resolved = NOT RECOVERED/NOT RESOLVED Every subject is counted once for the AE outcome, with the order: Fatal>Not Resolved>Resolving>Unknown>Sequelae>Resolved.			
<u>Pembrolizumab IV</u>			
In the NSCLC population in KN024 and KN042 (n=790), the outcomes of the 4 patients with events of nephritis were as follows: 2 (0.3%) resolved, 1 (0.1%) resolving, 1 (0.1%) resolved with sequelae.			
In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), the outcomes of the 7 patients with events of nephritis were as follows: 2 (0.4%) resolved, 3 (0.6%) resolving, 2 (0.4%) not resolved. There were no patients with events of nephritis with a fatal outcome.			
In the HL population in KN013, KN087, and KN204 (n=389), the outcome of the 2 (0.5%) patients with events of nephritis were as follows: 1 (0.3%) not resolved and 1 (0.3%) resolved. There were no fatal outcomes.			
In the UC population in KN052 (n=370), the outcome for the 1 patient with an event of nephritis was: 1 (0.3%) resolved and in the UC population in KN045 (n=266), the outcomes for the 2 patients with events of nephritis were as follows: 2 (0.8%) resolved. There were no patients with events of nephritis with a fatal outcome.			
In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), the outcomes for the 2 patients with events of nephritis were as follows: 1 (0.3%) not resolved and 1 (0.3%) resolved with sequelae. There were no fatal outcomes. In the HNSCC population in KN048 pembrolizumab plus chemotherapy group (n=276), the outcome for the 2 patients with events of nephritis were as follows: 1 (0.4%) resolved and 1 (0.4%) not resolved. There were no patients with events of nephritis with a fatal outcome.			
In the melanoma population in KN054 (n=509), the outcome of the 2 patients with events of nephritis were as follows: 1 (0.2%) not resolved and 1 (0.2%) resolved with sequelae. There were no patients with events of nephritis with a fatal outcome.			
In the NSCLC population in KN407 and KN021 (Cohort A) (n=303), the outcomes of the 2 patients with events of nephritis were as follows: 1 (0.3%) resolved, 1 (0.3%) resolving. There were no patients with events of nephritis with a fatal outcome.			
In the RCC population in KN426 pembrolizumab plus axitinib (n=429), the outcomes for the 6 patients with events of nephritis were as follows: 4 (0.9%) resolved and 2 (0.5%) not resolved. There were no patients with events of nephritis with a fatal outcome.			
In the CRC population in KN177 (n=153), the outcome of the 1 (0.7%) patient with an event of nephritis was not resolved. There were no patients with events of nephritis with a fatal outcome.			
In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), the outcome of the 1 patient with an event of nephritis was as follows: 1 (0.3%) not resolved. There were no patients with events of nephritis with a fatal outcome.			

Table SVII.3.1.4: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Nephritis

	<p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), the outcomes for the patients with events of nephritis were as follows: 2 (0.3%) resolved, 1 (0.2%) not resolved, and 1 (0.2%) resolved with sequelae. There were no patients with events of nephritis with a fatal outcome.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), the outcome of the 6 (1.7%) patients with events of nephritis were as follows: 2 (0.6%) resolved, 1 (0.3%) resolving, 1 (0.3%) resolved with sequelae, and 1 (0.3%) not resolved. There was 1 (0.3%) patient with an event of nephritis that resulted in a fatal outcome.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), the outcomes for the 5 patients with events of nephritis were as follows: 4 (0.8%) resolved and 1 (0.2%) not resolved. There were no patients with events of nephritis with a fatal outcome.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488), the outcomes for the 3 patients with events of nephritis were as follows: 1 (0.2%) resolved, 1 (0.2%) resolving, and 1 (0.2%) resolved with sequelae. There were no patients with events of nephritis with a fatal outcome.</p> <p>In the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), the outcomes for the 2 (0.4%) patients with events of nephritis were as follows: 1 (0.2%) resolved and 1 (0.2%) resolving. There were no patients with events of nephritis with a fatal outcome.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), the outcome of the 1 patient with an event of nephritis was as follows: 1 (0.3%) not resolved. There were no patients with events of nephritis with a fatal outcome.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), the outcomes for the 7 patients with events of nephritis were as follows: 5 (0.6%) resolved, 1 (0.1%) not resolved, and 1 (0.1%) unknown. There were no patients with events of nephritis with a fatal outcome.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), the outcomes of the 7 patients with events of nephritis were as follows: 6 (1.2%) resolved, and 1 (0.2%) not resolved. There were no patients with events of nephritis with a fatal outcome.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), the outcomes for the 4 patients with events of nephritis were as follows: 2 (0.6%) resolving and 2 (0.6%) not resolved. There were no patients with events of nephritis with a fatal outcome.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580), the outcomes for the 4 patients with events of nephritis were as follows: 3 (0.5%) resolved and 1 (0.2%) not resolved. There were no patients with events of nephritis with a fatal outcome.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), the outcomes for the 4 patients with events of nephritis were as follows: 3 (0.4%) not resolved, and 1 (0.1%) resolved. There were no patients with events of nephritis with a fatal outcome.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), the worst reported outcomes for the 2 patients with events of nephritis were as follows: 1 (0.2%) resolved and 1 (0.2%) not resolved. There were no patients with events of nephritis with a fatal outcome.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), the outcomes for the 6 patients with events of nephritis were as follows: 5 (0.9%) resolved, and 1 (0.2%) not resolved. There were no patients with an event of nephritis with a fatal outcome.</p> <p>In the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382), the outcomes for patients with events of nephritis were not recorded per study protocol (managed by external collaboration).</p> <p>In the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528), the outcomes for the 2 patients with events of nephritis were resolved (0.4%). There were no patients with events of nephritis with a fatal outcome.</p> <p>In the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), the reported outcome for the 2 patients with events of nephritis was 2 (0.8%) resolved. There were no patients with events of nephritis with a fatal outcome.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without g</p>
--	--

Table SVII.3.1.4: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Nephritis

	<p>chemotherapy and then as monotherapy (n=361), the outcome for the 3 patients with events of nephritis were as follows: 3 (0.8%) resolved. There were no fatal outcomes.</p> <p><u>Pembrolizumab SC</u></p> <p>There were no events of nephritis reported in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251).</p> <p>Aggregate Review</p> <p>Review of the ASE dataset (n=7730) yielded no fatal cases of nephritis. Review of the global safety database also yielded no fatal cases from ongoing interventional monotherapy clinical trials and no fatal cases from the PM or non-interventional environment.</p>																								
<p>Seriousness</p>	<p>Percent serious for identified risk of nephritis- pembrolizumab (Serious AEs for 90 days after discontinuation were included.)</p> <p style="text-align: center;">Nephritis - Serious (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment Emergent)</p> <table border="1" data-bbox="464 827 1383 1062"> <thead> <tr> <th></th> <th colspan="2">Cumulative NSCLC 001/010 + MEL 001/002/006</th> </tr> <tr> <th></th> <th>n</th> <th>(%)</th> </tr> </thead> <tbody> <tr> <td>Subjects in population</td> <td>2,799</td> <td></td> </tr> <tr> <td>with one or more adverse events</td> <td>8</td> <td>(0.3)</td> </tr> <tr> <td>Acute kidney injury</td> <td>2</td> <td>(0.1)</td> </tr> <tr> <td>Autoimmune nephritis</td> <td>1</td> <td>(0.0)</td> </tr> <tr> <td>Renal failure</td> <td>2</td> <td>(0.1)</td> </tr> <tr> <td>Tubulointerstitial nephritis</td> <td>3</td> <td>(0.1)</td> </tr> </tbody> </table> <p>Every subject is counted a single time for each applicable row and column. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns meets the incidence criterion in the report title, after rounding. Include all treated subjects in KN001 Part B1, B2, B3, D, C, F1, F2, F3 and all subjects in KN002 treated with Pembrolizumab in the original phase, and all subjects in KN006 and KN010 treated with Pembrolizumab.</p> <p><u>Pembrolizumab IV</u></p> <p>The 4 patients with events of nephritis reported in the NSCLC population in KN024 and KN042 (n=790) 3(0.4%) experienced serious events of nephritis.</p> <p>Of the 7 patients with events of nephritis in the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), 5 (1.0%) experienced serious events of nephritis.</p> <p>There was 1 (0.3%) patient with a serious event of nephritis reported in the HL population in KN013, KN087, and KN204 (n=389).</p> <p>The 1 (0.3%) patient with an event of nephritis in the UC population in KN052 (n=370) experienced a serious event of nephritis and the 2 (0.8%) patients with events of nephritis in the UC population in KN045 (n=266) experienced serious events of nephritis.</p> <p>Of the 2 patients with events of nephritis reported in the HNSCC pembrolizumab monotherapy population in KN048 (n=300), 1 (0.3%) experienced a serious event of nephritis. Of the 2 patients with events of nephritis reported in the HNSCC pembrolizumab plus chemotherapy group in KN048 (n=276), none experienced serious events of nephritis.</p> <p>The 2 patients with events of nephritis reported in the melanoma population in KN054 (n=509) experienced serious events of nephritis (0.4%).</p> <p>Of the 2 patients with events (0.7%) of nephritis in the NSCLC population in KN407 and KN021 (Cohort A) (n=303), both experienced serious events of nephritis.</p>		Cumulative NSCLC 001/010 + MEL 001/002/006			n	(%)	Subjects in population	2,799		with one or more adverse events	8	(0.3)	Acute kidney injury	2	(0.1)	Autoimmune nephritis	1	(0.0)	Renal failure	2	(0.1)	Tubulointerstitial nephritis	3	(0.1)
	Cumulative NSCLC 001/010 + MEL 001/002/006																								
	n	(%)																							
Subjects in population	2,799																								
with one or more adverse events	8	(0.3)																							
Acute kidney injury	2	(0.1)																							
Autoimmune nephritis	1	(0.0)																							
Renal failure	2	(0.1)																							
Tubulointerstitial nephritis	3	(0.1)																							

Table SVII.3.1.4: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Nephritis

	<p>Of the 6 patients with events of nephritis in the RCC population in KN426 pembrolizumab plus axitinib (n=429), 2 (0.5%) experienced serious events of nephritis.</p> <p>The one patient with an event of nephritis in the CRC population in KN177 (n=153) experienced a serious event of nephritis (0.7%).</p> <p>There were no patients with serious events of nephritis in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370).</p> <p>Of the 4 patients with events of nephritis in the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), 1 (0.2%) experienced a serious event of nephritis.</p> <p>Of the 6 (1.7%) patients with events of nephritis in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), 3 (0.9%) experienced serious events of nephritis.</p> <p>Of the 5 patients with events of nephritis in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), 3 (0.6%) experienced serious events of nephritis.</p> <p>Of the 3 patients with events of nephritis in the RCC population in KN564 pembrolizumab monotherapy (n=488), 1 (0.2%) experienced a serious event of nephritis.</p> <p>There were no patients with serious events of nephritis reported in the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475).</p> <p>There were no patients with serious events of nephritis in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307).</p> <p>Of the 7 patients with events of nephritis in the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), 6 (0.8%) experienced serious events of nephritis.</p> <p>Of the 7 patients with events of nephritis in the melanoma population in KN716 pembrolizumab monotherapy (n=483), 4 (0.8%) experienced serious events of nephritis.</p> <p>Of the 4 patients with events of nephritis in the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), 2 (0.6%) experienced serious events of nephritis.</p> <p>There were no patients with serious events of nephritis in the NSCLC population in KN091 pembrolizumab monotherapy (n=580).</p> <p>Of the 4 patients with events of nephritis in the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), 4 (0.5%) experienced serious events of nephritis.</p> <p>There were no patients with serious events of nephritis in the BTC population in KN966 pembrolizumab plus chemotherapy (n=529).</p> <p>Of the 6 patients with events of nephritis in the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), 4 (0.7%) experienced serious events of nephritis.</p> <p>There were no patients with serious events of nephritis in the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382).</p> <p>Of the 2 patients with events of nephritis in the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528), 2 (0.4%) experienced serious events of nephritis.</p> <p>Of the 2 patients with events (0.8%) of nephritis in the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), both experienced serious events of nephritis</p> <p>Of the 3 patients with events of nephritis reported in the HNSCC in KN689 population who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), 2 (0.6%) experienced a serious event of nephritis</p> <p><u>Pembrolizumab SC</u></p> <p>There were no events of nephritis reported in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251).</p>
--	---

Table SVII.3.1.4: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Nephritis

Severity and Nature of the Risk	Nephritis (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment Emergent)		
	Cumulative NSCLC 001/010 + MEL 001/002/006		
		n	(%)
	Subjects in population	2,799	
	with one or more adverse events	9	(0.3)
	Acute kidney injury	2	(0.1)
	Grade 3	1	(0.0)
	Grade 4	1	(0.0)
	Autoimmune nephritis	1	(0.0)
	Grade 2	1	(0.0)
	Renal failure	2	(0.1)
	Grade 3	2	(0.1)
	Tubulointerstitial nephritis	4	(0.1)
	Grade 1	1	(0.0)
	Grade 2	2	(0.1)
Grade 3	1	(0.0)	
<p>Every subject is counted a single time for each applicable specific adverse event. A subject with multiple adverse events within a system organ class is counted a single time for that system organ class. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the incidence specified in the report title, after rounding. Only the highest reported grade of a given adverse event is counted for the individual subject. Grades are based on NCI CTCAE version 4.0.</p>			
<u>Pembrolizumab IV</u>			
<p>In the NSCLC population in KN024 and KN042 (n=790), there were 2 (0.3%) patients with Grade 3 events of nephritis. There were no patients with Grade 4 or 5 events of nephritis.</p> <p>In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), there was 4 (0.8%) patients with Grade 3 events, 2 (0.4) patients with Grade 4 events and no patients with Grade 5 events of nephritis.</p> <p>In the HL population in KN013, KN087, and KN204 (n=389), there was 1 (0.3%) patient with a Grade 3 and 1 (0.3%) patient with a Grade 4 event of nephritis. There were no patients with Grade 5 events of nephritis.</p> <p>In the UC population in KN052 (n=370), there was 1 (0.3%) patient with a Grade 3 event and no patients with Grade 4 or 5 events of nephritis; in the UC population in KN045 (n=266), there were 2 (0.8%) patients with Grade 3 events and no patients with Grade 4 or 5 events of nephritis.</p> <p>In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), there was 1 (0.3%) patient with a Grade 3 event of nephritis. There were no patients with Grade 4 or 5 events of nephritis; in the pembrolizumab plus chemotherapy group (n=276), there were no patients with Grade 3-5 events of nephritis.</p> <p>In the melanoma population in KN054 (n=509), there were 2 (0.4%) patients with Grade 3 event of nephritis. There were no patients with Grade 4 or 5 events of nephritis.</p> <p>In the NSCLC population in KN407 and KN021 (Cohort A) (n=303), there were 2 (0.7%) patients with a Grade 3 events. There were no patients with Grade 4 or 5 events of nephritis.</p> <p>In the RCC population in KN426 pembrolizumab plus axitinib (n=429), there was 1 (0.3%) patient with Grade 3 event of nephritis. There were no patients with Grade 4 or 5 events of nephritis.</p> <p>In the CRC population in KN177 (n=153) there were no patients with Grade 3-5 events of nephritis reported.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), there were no patients with Grade 3-5 events of nephritis.</p>			

Table SVII.3.1.4: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Nephritis

	<p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), there were no patients with Grade 3-5 events of nephritis.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352) there were 3 (0.9%) patients with Grade 3, no patients with Grade 4, and 1 (0.3%) patient with a Grade 5 events of nephritis.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), there were 3 (0.6%) patients with Grade 3 events of nephritis. There were no patients with Grade 4 or 5 events of nephritis.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488) there was 1 (0.2%) patient with a Grade 3 event of nephritis. There were no patients with Grade 4 or 5 events of nephritis.</p> <p>In the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), there were no patients with Grade 3-5 events of nephritis.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), there were no patients with Grade 3-5 events of nephritis.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) there were 4 (0.5%) patients with Grade 3 events, and 2 (0.3%) patients with Grade 4 event of nephritis. There were no patients with Grade 5 events of nephritis reported.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), there were 2 (0.4%) patients with Grade 3 events of nephritis. There were no patients with Grade 4 or 5 events of nephritis.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), there was 1 (0.3%) patient with a Grade 3 event of nephritis. There were no patients with Grade 4 or 5 events of nephritis.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580) there were no patients with Grade 3, 4, or 5 events of nephritis.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), there were 4 (0.5%) patients with Grade 3 events of nephritis reported. There were no patients with Grade 4 or 5 events of nephritis.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), there were no patients with Grade 3, 4 or 5 events of nephritis.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), there were 3 (0.5%) patients with Grade 3 events of nephritis. There were no patients with Grade 4 or Grade 5 events of nephritis.</p> <p>In the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy group (n=382), there were no patients with Grade 3-5 events of nephritis.</p> <p>In the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528), there were 2 (0.4%) patients with a Grade 3 event of nephritis. There were no patients with Grade 4 or 5 events of nephritis.</p> <p>In the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), there were no patients with Grade 3, 4 or 5 events of nephritis.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), there was 1 (0.3%) patient with a Grade 3 event and 1 (0.3%) patient with Grade 4 event of nephritis. There were no patients with Grade 5 events of nephritis.</p> <p><u>Pembrolizumab SC</u></p> <p>There were no events of nephritis reported in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251).</p> <p>Review of other pembrolizumab clinical trial data from ongoing studies regarding immune-mediated nephritis is consistent with the risk as characterized in this RMP.</p>
<p><u>Risk Factors and Risk Groups:</u></p>	<p>Patients with severe renal dysfunction were excluded from clinical trials. No specific risk factors for nephritis associated with pembrolizumab have been identified.</p>

Table SVII.3.1.4: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Nephritis

<u>Preventability:</u>	Although the development of nephritis cannot be completely prevented; patients should be monitored for signs and symptoms of nephritis and if they develop, other causes excluded. Withholding/discontinuation of pembrolizumab and appropriate medical intervention including corticosteroids following the onset of nephritis may result in recovery.
<u>Impact on the Risk-Benefit Balance of the Product:</u>	Nephritis can potentially lead to severe or life threatening renal injury. Given the fatal outcome of untreated cancer, the risk of nephritis which generally can be managed is outweighed by the potential benefit.
<u>Public Health Impact:</u>	This risk has minimal public health impact outside its effect on individual patients.

Table SVII.3.1.5: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies – Hypophysitis (Including Hypopituitarism)

<u>Important Identified Risk:</u>	Immune-mediated Endocrinopathies- Hypophysitis (including hypopituitarism)
<u>Potential Mechanisms:</u>	Pembrolizumab is a potent and highly-selective humanized monoclonal antibody (mAb) of the Immunoglobulin G4 (IgG4)/kappa isotype directed to the programmed cell death-1 (PD-1) receptor and designed to directly block the interaction between the receptor and its ligands, PD-L1 and PD-L2. The PD-1 pathway represents a major immune control switch which may be engaged by tumor cells to overcome active T-cell immune surveillance. T-cell activation in proximity to normal tissue may lead to inflammation and injury to normal tissue.
<u>Evidence Source(s) and Strength of Evidence:</u>	Review of pembrolizumab clinical trial data, post-marketing experience and literature regarding endocrinopathies- hypophysitis represent sufficient evidence of a causal association with pembrolizumab exposure. CTD 2.7.4 Please reference evidence source cited in Table SVII.3.1.1; Details of Important Identified Risk: Immune-Mediated Pneumonitis
<u>Characterisation of the Risk:</u> Immune-mediated Endocrinopathies- Hypophysitis (including hypopituitarism)	

Table SVII.3.1.5: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies – Hypophysitis (Including Hypopituitarism)

Frequency with 95%CI	95% Confidence Interval for the Overall Incidence (%) of Hypophysitis KN001, KN002 and KN006 Melanoma and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab	
	Cumulative MEL 001/002/006 + NSCLC 001/010	
	n (%)	95% CI
	Hypophysitis*	17 (0.6) (0.4,1)
* includes hypopituitarism		
<u>Pembrolizumab IV</u>		
<p>The overall number and proportion of patients with hypophysitis in the NSCLC population in KN024 and KN042 (n=790) was 4 (0.5%) (95% CI-0.1, 1.3), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hypophysitis in the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488) was 3 (0.6%) (95% CI- 0.1, 1.8), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset for melanoma and NSCLC.</p> <p>There were no patients with events of patients with hypophysitis reported in the HL population in KN013, KN087, and KN204.</p> <p>The overall number and proportion of patients with hypophysitis in the UC population in KN052 (n=370) was 2 (0.5%) (95%CI-0.1,1.9) which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset for melanoma and NSCLC. There were no patients with events of hypophysitis reported in the UC population in KN045.</p> <p>There were no patients with events of hypophysitis reported in the HNSCC population in KN040, KN012 and KN055.</p> <p>The overall number and proportion of patients with hypophysitis in the HNSCC population in KN048 pembrolizumab monotherapy(n=300) and pembrolizumab plus chemotherapy (n=276) groups, were 1 (0.3%) (95% CI-0,1.8) and 1 (0.4%) (95% CI-0,2) respectively, which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hypophysitis in the melanoma population in KN054 (n=509) was 11 (2.2%) (95% CI-1.1,3.8). The proportion of patients with events of hypophysitis in KN054 although somewhat more common than in the reference safety dataset were generally consistent in nature with the proportion of events of hypophysitis previously reported for pembrolizumab.</p> <p>The overall number and proportion of patients with hypophysitis in the NSCLC population in KN407 and KN021 (Cohort A) (n=303), was 3 (1.0%) (95% CI-0.2, 2.9), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hypophysitis in the RCC population in KN426 pembrolizumab plus axitinib (n=429) was 5 (1.2%) (95% CI-0.4,2.7), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hypophysitis in the CRC population in KN177 (n=153) was 2 (1.3%) (95% CI-0.2,4.6) which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hypophysitis in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370) was 3 (0.8%) (95% CI-0.2,2.4), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset for melanoma and NSCLC.</p>		

Table SVII.3.1.5: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies – Hypophysitis (Including Hypopituitarism)

	<p>The overall number and proportion of patients with hypophysitis in the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596) was 1 (0.2%) (95% CI-0, 0.9), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hypophysitis in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352) was 3 (0.9%) (95% CI-0.2,2.5) which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hypophysitis in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530) was 3 (0.6%) (95% CI-0.1,1.6), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hypophysitis in the RCC population in KN564 pembrolizumab monotherapy (n=488) was 2 (0.4%) (95% CI-0,1.5), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset for melanoma and NSCLC.</p> <p>There were no patients with events of hypophysitis reported in the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B.</p> <p>The overall number and proportion of patients with hypophysitis in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307) was 1 (0.3%) (95% CI-0.0, 1.8), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hypophysitis in the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) was 15 (1.9%) (95% CI-1.1,3.1), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hypophysitis in the melanoma population in KN716 pembrolizumab monotherapy (n=483) was 10 (2.1%) (95% CI-1.0, 3.8), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hypophysitis in the gastric cancer population in KN811 pembrolizumab combination therapy (n=350) was 4 (1.1%) (95% CI-0.3,2.9), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hypophysitis in the NSCLC population in KN091 pembrolizumab monotherapy (n=580) was 7 (1.2%) (95% CI-0.5,2.5), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hypophysitis in the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785) was 3 (0.4%) (95% CI-0.1, 1.1), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hypophysitis in the BTC population in KN966 pembrolizumab plus chemotherapy (n=529) was 2 (0.4%) (95% CI-0.0, 1.4), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hypophysitis in the NSCLC population in KN671 pembrolizumab in combination with chemotherapy neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396) was 2 (0.5%) (95% CI-0.1, 1.8), which is generally similar to the overall proportion of patients with events of hypophysitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hypophysitis in the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564) was 5 (0.9%) (95% CI-0.3, 2.1), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset.</p>
--	--

Table SVII.3.1.5: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies – Hypophysitis (Including Hypopituitarism)

	<p>The overall number and proportion of patients with hypophysitis in the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382) was 2 (0.5%) (95% CI-0.1,1.9), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset.</p> <p>There were no events of hypophysitis reported in the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy.</p> <p>The overall number and proportion of patients with hypophysitis in the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260) was 1 (0.4%) (95% CI-0.0, 2.1), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hypophysitis in the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361) were 2 (0.6%) (95% CI-0.1,2.0), which is generally similar to the proportion of patients with the event of hypophysitis in the reference safety dataset.</p> <p><u>Pembrolizumab SC</u></p> <p>There were no events of hypophysitis reported in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy.</p>																															
<p>Outcomes</p>	<p>Percent overall, fatal, not resolved, resolved, and unknown for identified risk of hypopituitarism, hypophysitis - pembrolizumab. (AEs for 30 days after discontinuation were included)</p> <p style="text-align: center;">Hypopituitarism and Hypophysitis (AEOSI) KN001, KN002 and KN006 Melanoma and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment-Emergent)</p> <table border="1" data-bbox="496 1020 1414 1331"> <thead> <tr> <th colspan="2"></th> <th colspan="2">Cumulative NSCLC 001/010 + MEL 001/002/006</th> </tr> <tr> <th></th> <th>Outcome</th> <th>n</th> <th>(%)</th> </tr> </thead> <tbody> <tr> <td>Subjects in population</td> <td></td> <td>2,799</td> <td></td> </tr> <tr> <td rowspan="6">With one or more adverse events</td> <td>Overall</td> <td>17</td> <td>(0.6)</td> </tr> <tr> <td>Fatal</td> <td>0</td> <td>(0.0)</td> </tr> <tr> <td>Not Resolved</td> <td>10</td> <td>(0.4)</td> </tr> <tr> <td>Resolved</td> <td>5</td> <td>(0.2)</td> </tr> <tr> <td>Sequelae</td> <td>2</td> <td>(0.1)</td> </tr> <tr> <td>Unknown</td> <td>0</td> <td>(0.0)</td> </tr> </tbody> </table> <p>Every subject is counted once on each applicable row. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the percent incidence specified in the report title, after rounding. Outcome: Resolved = RECOVERED/RESOLVED, Not resolved = NOT RECOVERED/NOT RESOLVED.</p> <p><u>Pembrolizumab IV</u></p> <p>In the NSCLC population in KN024 and KN042 (n=790), the outcome of the 4 (0.5%) patients with events of hypophysitis. were as follows: 2 (0.3%) resolved, 2 (0.3%) not resolved. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), the outcome of the 3 patients with events of hypophysitis were as follows: 1 (0.2%) resolved, 1 (0.2%) resolving, and 1 (0.2%) resolved with sequelae. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the UC population in KN052 (n=370), the outcomes for the 2 patients with events of hypophysitis were as follows: 2 (0.5%) resolved. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), the outcome for the 1 (0.3%) patient with an event of hypophysitis was resolved with sequelae. There were no fatal</p>			Cumulative NSCLC 001/010 + MEL 001/002/006			Outcome	n	(%)	Subjects in population		2,799		With one or more adverse events	Overall	17	(0.6)	Fatal	0	(0.0)	Not Resolved	10	(0.4)	Resolved	5	(0.2)	Sequelae	2	(0.1)	Unknown	0	(0.0)
		Cumulative NSCLC 001/010 + MEL 001/002/006																														
	Outcome	n	(%)																													
Subjects in population		2,799																														
With one or more adverse events	Overall	17	(0.6)																													
	Fatal	0	(0.0)																													
	Not Resolved	10	(0.4)																													
	Resolved	5	(0.2)																													
	Sequelae	2	(0.1)																													
	Unknown	0	(0.0)																													

Table SVII.3.1.5: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies – Hypophysitis (Including Hypopituitarism)

	<p>outcomes. In the HNSCC population in KN048 pembrolizumab plus chemotherapy group (n=276), the outcome for the 1 (0.4%) patient with an event of hypophysitis was not resolved. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the melanoma population in KN054 (n=509), the outcomes for the 11 patients with events of hypophysitis were as follows: 1 (0.2%) resolved, 5 (1%) not resolved and 5 (1%) resolved with sequelae. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the NSCLC population in KN407 and KN021 (Cohort A) (n=303), the outcomes for the 3 patients with events of hypophysitis were as follows: 2 (0.7%) resolving, 1 (0.3%) resolved. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the RCC population in KN426 pembrolizumab plus axitinib (n=429) the outcomes for the 5 patients with events of hypophysitis were as follows: 2 (0.5%) resolved, 2 (0.5%) not resolved and 1 (0.2%) resolving. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the CRC population in KN177 (n=153), the outcomes for the 2 patients with events of hypophysitis were as follows: 2 (1.3%) not resolved. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), the outcomes for the 3 patients with events of hypophysitis were as follows: 2 (0.5%) resolved with sequelae and 1 (0.3%) not resolved. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), the outcome for the 1 (0.2%) patient with an event of hypophysitis was resolved.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), the outcomes for the 3 (0.9%) patients with events of hypophysitis were as follows: 2 (0.6%) not resolved, and 1 (0.3%) resolving. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530) the outcomes for the 3 patients with events of hypophysitis were as follows: 2 (0.4%) not resolved, and 1 (0.2%) resolved. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488), the outcomes for the 2 patients with events of hypophysitis were as follows: 2 (0.4%) resolved. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), the outcome for the 1 (0.3%) patient with an event of hypophysitis was resolving. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), the outcomes for the 15 patients with events of hypophysitis were as follows: 3 (0.4%) resolved, 6 (0.8%) not resolved, 5 (0.6%) resolved with sequelae, and 1 (0.1%) resolving. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), the outcomes for the 10 patients with events of hypophysitis were as follows: 5 (1%) not resolved, 3 (0.6%) resolving, 2 (0.4%) resolved. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), the outcome for the 4 patients with events of hypophysitis were as follows: 2 (0.6%) resolved, 1 (0.3%) not resolved and 1 (0.3%) resolving. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580), the outcomes for the 7 patients with events of hypophysitis were as follows: 6 (1%) not resolved and 1 (0.2%) resolved. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), the outcomes for the 3 patients with events of hypophysitis were as follows: 2 (0.3%) not resolved,</p>
--	--

Table SVII.3.1.5: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies – Hypophysitis (Including Hypopituitarism)

	<p>and 1 (0.1%) resolved with sequelae. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), the worst reported outcomes for the 2 (0.4%) patients with events of hypophysitis were not resolved. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy group (n=396), the outcomes for the 2 patients with events of hypophysitis were as follows: 1 (0.3%) resolved, and 1 (0.3%) resolving. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), the outcomes for the 5 patients with events of hypophysitis were as follows: 2 (0.4%) resolved and 3 (0.5%) not resolved. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382), the outcomes for patients with events of hypophysitis were not recorded per study protocol (managed by external collaboration).</p> <p>In the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), the reported outcomes for the 1 (0.4%) patient with an event of hypophysitis was not resolved. There were no patients with events of hypophysitis with a fatal outcome.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), the outcome for the 2 patients with events of hypophysitis were 1 (0.3%) resolved and 1 (0.3%) resolved with sequelae. There were no fatal outcomes.</p> <p><u>Pembrolizumab SC</u></p> <p>There were no events of hypophysitis reported in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy.</p> <p>Aggregate Review</p> <p>Review of the ASE dataset (n=7730) yielded no fatal cases of hypophysitis. Review of the global safety database also yielded no fatal cases from ongoing interventional monotherapy clinical trials and no fatal cases from the PM or non-interventional environment.</p>																	
<p>Seriousness</p>	<p>Percent serious for identified risk of hypopituitarism, hypophysitis – pembrolizumab (Serious AEs for 90 days after discontinuation were included.)</p> <p>Hypopituitarism and Hypophysitis - Serious (AEOSI)</p> <p>KN001, KN002 and KN006 Melanoma and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment-Emergent)</p> <table border="1" data-bbox="495 1495 1414 1717"> <thead> <tr> <th rowspan="2"></th> <th colspan="2">Cumulative NSCLC 001/010 + MEL 001/002/006</th> </tr> <tr> <th>n</th> <th>(%)</th> </tr> </thead> <tbody> <tr> <td>Subjects in population</td> <td>2,799</td> <td></td> </tr> <tr> <td>with one or more adverse events</td> <td>10</td> <td>(0.4)</td> </tr> <tr> <td>Hypophysitis</td> <td>5</td> <td>(0.2)</td> </tr> <tr> <td>Hypopituitarism</td> <td>5</td> <td>(0.2)</td> </tr> </tbody> </table> <p>Every subject is counted a single time for each applicable row and column. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns meets the incidence criterion in the report title, after rounding. Include all treated subjects in KN001 Part B1, B2, B3, D, C, F1, F2, F3 and all subjects in KN002 treated with Pembrolizumab in the original phase , and all subjects in KN006 and KN010 treated with Pembrolizumab.</p>		Cumulative NSCLC 001/010 + MEL 001/002/006		n	(%)	Subjects in population	2,799		with one or more adverse events	10	(0.4)	Hypophysitis	5	(0.2)	Hypopituitarism	5	(0.2)
	Cumulative NSCLC 001/010 + MEL 001/002/006																	
	n	(%)																
Subjects in population	2,799																	
with one or more adverse events	10	(0.4)																
Hypophysitis	5	(0.2)																
Hypopituitarism	5	(0.2)																

Table SVII.3.1.5: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies – Hypophysitis (Including Hypopituitarism)

	<p><u>Pembrolizumab IV</u></p> <p>The 4 patients with events of hypophysitis reported in the NSCLC population in KN024 and KN042 (n=790) experienced serious (0.5%) events of hypophysitis.</p> <p>There were no patients with serious events of hypophysitis reported in the NSCLC population in KN189 or KN021 (Cohorts C and G) pembrolizumab plus chemotherapy.</p> <p>The 2 (0.5%) patients with events of hypophysitis reported in the UC population in KN052 (n=370) experienced serious events of hypophysitis.</p> <p>The 1 (0.3%) patient with an event of hypophysitis reported in the HNSCC population in KN048 pembrolizumab monotherapy group (n=300) was serious. The 1 (0.4%) patient with an event of hypophysitis reported in the HNSCC population in KN048 pembrolizumab plus chemotherapy group (n=276) experienced a serious event of hypophysitis.</p> <p>Of the 11 patients with events of hypophysitis reported in the melanoma population in KN054 (n=509), 4 (0.8%) experienced serious events of hypophysitis.</p> <p>Of the 3 patients with events of hypophysitis reported in the NSCLC population in KN407 and KN021 (Cohort A) (n=303), 2 (0.7%) experienced serious events of hypophysitis.</p> <p>Of the 5 patients with events of hypophysitis in the RCC population in KN426 pembrolizumab plus axitinib (n=429), 2 (0.5%) experienced serious events of hypophysitis.</p> <p>There were no patients with serious adverse events of hypophysitis in the CRC population in KN177 (n=153).</p> <p>Of the 3 patients with events of hypophysitis in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), 2 experienced serious events of hypophysitis.</p> <p>There were no patients with serious events of hypophysitis in the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596).</p> <p>Of the 3 patients with events of hypophysitis in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), 2 (0.6%) experienced serious events of hypophysitis.</p> <p>Of the 3 patients with events of hypophysitis in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), 1 (0.2%) experienced a serious event of hypophysitis.</p> <p>Of the 2 patients with events of hypophysitis in the RCC population in KN564 pembrolizumab monotherapy (n=488), 1 (0.2%) experienced a serious event of hypophysitis.</p> <p>The 1 (0.3%) patient with an event of hypophysitis reported in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307) experienced a serious event of hypophysitis.</p> <p>Of the 15 patients with events of hypophysitis in the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), 10 (1.3%) experienced serious events of hypophysitis.</p> <p>Of the 10 patients with events of hypophysitis in the melanoma population in KN716 (n=483), 4 (0.8%) experienced serious events of hypophysitis.</p> <p>Of the 4 patients with events of hypophysitis in gastric cancer population in KN811 pembrolizumab combination therapy (n=350), 3 (0.9%) experienced serious events of hypophysitis.</p> <p>Of the 4 patients with events of hypophysitis in the NSCLC population in KN091 pembrolizumab monotherapy (n=580), 4 (0.7%) experienced serious events of hypophysitis.</p> <p>Of the 3 patients with events of hypophysitis in the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), 1 (0.1%) experienced a serious event of hypophysitis.</p> <p>Of the 2 patients with events of hypophysitis in the BTC population in KN966 (n=529), 1 (0.2%) experienced a serious event of hypophysitis.</p> <p>Of the 2 patients with events of hypophysitis in the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396), 1 (0.3%) experienced a serious event of hypophysitis.</p>
--	---

Table SVII.3.1.5: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies – Hypophysitis (Including Hypopituitarism)

	<p>There were no patients with serious adverse events of hypophysitis in the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564).</p> <p>Of the 2 patients with events of hypophysitis in the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382), 2 (0.5%) experienced serious events of hypophysitis. The 1 (0.4%) patient in the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260) experienced a serious event of hypophysitis.</p> <p>Of the 2 patients with events of hypophysitis reported in the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), 1 (0.3%) experienced a serious event of hypophysitis.</p> <p><u>Pembrolizumab SC</u></p> <p>There were no events of hypophysitis reported in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy.</p>																																			
<p>Severity and Nature of the Risk</p>	<p>Identified treatment emergent adverse events (AE) of hypopituitarism, hypophysitis (all events) in participants summarized by grade– pembrolizumab. (Table includes events for 30 days after discontinuation.)</p> <p style="text-align: center;">Hypopituitarism and Hypophysitis (AEOSI) KN001, KN002 and KN006 Melanoma and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment-Emergent)</p> <table border="1" data-bbox="495 911 1412 1276"> <thead> <tr> <th rowspan="2"></th> <th colspan="2">Cumulative NSCLC 001/010 + MEL 001/002/006</th> </tr> <tr> <th>n</th> <th>(%)</th> </tr> </thead> <tbody> <tr> <td>Subjects in population</td> <td>2,799</td> <td></td> </tr> <tr> <td>with one or more adverse events</td> <td>17</td> <td>(0.6)</td> </tr> <tr> <td>Hypophysitis</td> <td>9</td> <td>(0.3)</td> </tr> <tr> <td> Grade 2</td> <td>5</td> <td>(0.2)</td> </tr> <tr> <td> Grade 3</td> <td>3</td> <td>(0.1)</td> </tr> <tr> <td> Grade 4</td> <td>1</td> <td>(0.0)</td> </tr> <tr> <td>Hypopituitarism</td> <td>8</td> <td>(0.3)</td> </tr> <tr> <td> Grade 1</td> <td>2</td> <td>(0.1)</td> </tr> <tr> <td> Grade 2</td> <td>1</td> <td>(0.0)</td> </tr> <tr> <td> Grade 3</td> <td>5</td> <td>(0.2)</td> </tr> </tbody> </table> <p>Every subject is counted a single time for each applicable specific adverse event. A subject with multiple adverse events within a system organ class is counted a single time for that system organ class.</p> <p>A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the incidence specified in the report title, after rounding. Only the highest reported grade of a given adverse event is counted for the individual subject. Grades are based on NCI CTCAE version 4.0.</p> <p><u>Pembrolizumab IV</u></p> <p>In the NSCLC population in KN024 and KN042 (n=790), there were 4 patients with Grade 3 events of hypophysitis (0.5%). There were no patients with Grade 4 or 5 events of hypophysitis.</p> <p>In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), there were no patients with Grade 3, 4 or 5 events of hypophysitis.</p> <p>In the UC population in KN052 (n=370), there were 2 (0.5%) patients with Grade 3 events and no patients with Grade 4 or 5 events of hypophysitis.</p> <p>In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), there was 1 patient with a Grade 3 event of hypophysitis (0.3%). There were no patients with Grade 4 or 5 events of hypophysitis. In the HNSCC population in KN048 pembrolizumab plus chemotherapy group (n=276), there was 1 patient with a Grade 3 event of hypophysitis (0.4%). There were no patients with Grade 4 or 5 events of hypophysitis.</p>		Cumulative NSCLC 001/010 + MEL 001/002/006		n	(%)	Subjects in population	2,799		with one or more adverse events	17	(0.6)	Hypophysitis	9	(0.3)	Grade 2	5	(0.2)	Grade 3	3	(0.1)	Grade 4	1	(0.0)	Hypopituitarism	8	(0.3)	Grade 1	2	(0.1)	Grade 2	1	(0.0)	Grade 3	5	(0.2)
	Cumulative NSCLC 001/010 + MEL 001/002/006																																			
	n	(%)																																		
Subjects in population	2,799																																			
with one or more adverse events	17	(0.6)																																		
Hypophysitis	9	(0.3)																																		
Grade 2	5	(0.2)																																		
Grade 3	3	(0.1)																																		
Grade 4	1	(0.0)																																		
Hypopituitarism	8	(0.3)																																		
Grade 1	2	(0.1)																																		
Grade 2	1	(0.0)																																		
Grade 3	5	(0.2)																																		

Table SVII.3.1.5: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies – Hypophysitis (Including Hypopituitarism)

<p>In the melanoma population in KN054 (n=509), there were 3 (0.6%) patients with Grade 3 events and no patients with Grade 4 or 5 events of hypophysitis.</p> <p>In the NSCLC population in KN407 and KN021 (Cohort A) (n=303), there was 1 (0.3%) patient with a Grade 3 event and 1 (0.3%) patient with a Grade 4 event. There were no patients with Grade 5 events of hypophysitis.</p> <p>In the RCC population in KN426 pembrolizumab plus axitinib (n=429), there were 4 (0.9%) patients with Grade 3 events of hypophysitis. There were no patients with Grade 4 or 5 events of hypophysitis.</p> <p>In the CRC population in KN177 (n=153) there were no patients with Grade 3-5 events of hypophysitis reported.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), there was 1 (0.3%) patient with a Grade 3 event of hypophysitis. There were no patients with Grade 4 or 5 events of hypophysitis.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), there were no patients with Grade 3-5 events of hypophysitis.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), there were 2 (0.6%) patients with Grade 3 events of hypophysitis. There were no patients with Grade 4 or 5 events of hypophysitis.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), there were 2 (0.4%) patients with Grade 3 events of hypophysitis. There were no patients with Grade 4 or 5 events of hypophysitis.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488) there were 2 (0.4%) patients with Grade 3 events of hypophysitis. There were no patients with Grade 4 or 5 events of hypophysitis.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), there was 1 (0.3%) patient with a Grade 3 event of hypophysitis. There were no patients with Grade 4 or 5 events of hypophysitis.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) there were 10 (1.3%) patients with Grade 3 events of hypophysitis. There were no patients with Grade 4 or 5 events of hypophysitis reported.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), there were 3 (0.6%) patients with Grade 3 events of hypophysitis. There were no patients with Grade 4 or 5 events of hypophysitis.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), there were 2 (0.6%) patients with Grade 3 events of hypophysitis. There were no patients with Grade 4 or 5 events of hypophysitis.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580) there were 3 (0.5%) patients with Grade 3 events of hypophysitis. There were no patients with Grade 4 or 5 events of hypophysitis.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), there was 1 (0.1%) patient with a Grade 3 event of hypophysitis reported. There were no patients with Grade 4 or 5 events of hypophysitis.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), there was 1 (0.2%) patient with a Grade 3 event of hypophysitis reported. There were no patients with Grade 4 or 5 events of hypophysitis.</p> <p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396) there was 1 (0.3%) patient with a Grade 3 event of hypophysitis, there were no patients with Grade 4 or 5 events of hypophysitis reported.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), there were no patients with Grade 3-5 events of hypophysitis.</p>
--

Table SVII.3.1.5: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies – Hypophysitis (Including Hypopituitarism)

	<p>In the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382), there was 1 (0.3%) patient with a Grade 3 event, and 1 (0.3%) patient with a Grade 4 event of hypophysitis. There were no patients with Grade 5 events of hypophysitis.</p> <p>In the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), there were no patients with Grade 3, 4 or 5 events of hypophysitis.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), there was 1 patient with a Grade 3 event of hypophysitis (0.3%). There were no patients with Grade 4 or 5 events of hypophysitis.</p> <p><u>Pembrolizumab SC</u></p> <p>There were no events of hypophysitis reported in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy.</p>
<u>Risk Factors and Risk Groups:</u>	No specific risk factors for hypophysitis (including hypopituitarism) associated with pembrolizumab have been identified.
<u>Preventability:</u>	Although the development of hypophysitis (including hypopituitarism) cannot be completely prevented; patients should be monitored for signs and symptoms of hypophysitis (including hypopituitarism) and if they develop, other causes excluded. Withholding/discontinuation of pembrolizumab and appropriate medical intervention including corticosteroids following the onset of hypophysitis (including hypopituitarism) may result in recovery.
<u>Impact on the Risk-Benefit Balance of the Product:</u>	Hypophysitis and hypopituitarism have the potential to become severe or life-threatening due to hormone deficiencies in individual patients. Given the fatal outcome of untreated cancer, the risk of hypophysitis (including hypopituitarism) is outweighed by the potential benefit.
<u>Public Health Impact:</u>	This risk has minimal public health impact outside its effect on individual patients.

Table SVII.3.1.6: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies –Adrenal Insufficiency (Primary and Secondary)

<u>Important Identified Risk:</u>	Immune-mediated Endocrinopathies- Adrenal Insufficiency (primary and secondary)
<u>Potential Mechanisms:</u>	Pembrolizumab is a potent and highly-selective humanized monoclonal antibody (mAb) of the Immunoglobulin G4 (IgG4)/kappa isotype directed to the programmed cell death-1 (PD-1) receptor and designed to directly block the interaction between the receptor and its ligands, PD-L1 and PD-L2. The PD-1 pathway represents a major immune control switch which may be engaged by tumor cells to overcome active T-cell immune surveillance. T-cell activation in proximity to normal tissue may lead to inflammation and injury to normal tissue.
<u>Evidence Source(s) and Strength of Evidence:</u>	<p>Review of pembrolizumab clinical trial data, post-marketing experience and literature regarding endocrinopathies- Adrenal insufficiency (primary and secondary) represent sufficient evidence of a causal association with pembrolizumab exposure.</p> <p>CTD 2.7.4</p> <p>Please reference evidence source cited in Table SVII.3.1.1; Details of Important Identified Risk: Immune-Mediated Pneumonitis</p>

Table SVII.3.1.6: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies –Adrenal Insufficiency (Primary and Secondary)

Characterisation of the Risk: Immune-mediated Endocrinopathies- Adrenal Insufficiency (primary and secondary)		
Frequency with 95%CI	95% Confidence Interval for the Overall Incidence (%) of AEOSI- Adrenal Insufficiency KN001, KN002 and KN006 Melanoma and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab	
	Cumulative MEL 001/002/006 + NSCLC 001/010	
	n (%)	95% CI
	Adrenal Insufficiency*	22 (0.8) (0.5, 1.2)
*Includes adrenocortical insufficiency acute, secondary adrenocortical insufficiency, primary adrenal insufficiency and Addison's disease		
<u>Pembrolizumab IV</u>		
<p>The overall number and proportion of patients with adrenal insufficiency in the NSCLC population in KN024 and KN042 (n=790), was 4 (0.5%) (95% CI-0.1,1.3), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the NSCLC population in KN189 and KN021 (Cohorts C and G) (n=488) was 3 (0.6%) (95% CI- 0.1, 1.8), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with adrenal insufficiency reported in the HL population in KN013, KN087, and KN204 (n=389) was 1 (0.3%) (95%CI-0.0, 1.4), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the UC population in KN052 (n=370) was 6 (1.6%) (95%CI-0.6, 3.5) and in KN045 (n=266) was 1 (0.4%) (95%CI-0, 2.1), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the HNSCC population in KN040, KN012 and KN055 (n=609) was 4 (0.7%) (95%CI-0.2, 1.7), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the HNSCC population in KN048 pembrolizumab monotherapy (n=300) and pembrolizumab plus chemotherapy (n=276) groups, were 1 (0.3%) (95% CI-0,1.8) and 0 (0%) (95% CI-0,1.3) respectively, which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the melanoma population in KN054 (n=509) was 5 (1%) (95% CI-0.3, 2.3) which is generally similar to the overall frequency in the reference safety dataset for melanoma and NSCLC.</p> <p>There were no patients with events of adrenal insufficiency in the NSCLC population in KN407 and KN021 (Cohort A).</p> <p>The overall number and proportion of patients with adrenal insufficiency in the RCC population in KN426 pembrolizumab plus axitinib (n=429) was 13 (3.0%) (95% CI-1.6,5.1). The proportion of patients with events of adrenal insufficiency, in the RCC population in KN426 pembrolizumab plus axitinib, although somewhat more common than in the reference safety dataset were generally consistent in nature with the proportion of patients with events of adrenal insufficiency previously reported for pembrolizumab.</p>		

Table SVII.3.1.6: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies –Adrenal Insufficiency (Primary and Secondary)

	<p>The overall number and proportion of patients with adrenal insufficiency in the CRC population in KN177 (n=153) was 4 (2.6%) (95% CI-0.7,6.6) which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370) was 4 (1.1%) (95% CI-0.3,2.7), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596) was 7 (1.2%) (95% CI-0.5,2.4), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352) was 18 (5.1%) (95% CI-3.1,8.0). The proportion of patients with events of adrenal insufficiency, although higher than the reference safety dataset, were generally consistent in nature with the proportion of patients with events of adrenal insufficiency previously reported for pembrolizumab.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530) was 9 (1.7%) (95% CI-0.8, 3.2), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the RCC population in KN564 pembrolizumab monotherapy (n=488) was 10 (2%) (95% CI-1,3.7), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset for melanoma and NSCLC.</p> <p>There were no patients with events of adrenal insufficiency reported in the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307) was 4 (1.3%) (95% CI-0.4, 3.3), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) was 20 (2.6%) (95% CI-1.6,3.9), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the melanoma population in KN716 pembrolizumab monotherapy (n=483) was 11 (2.3%) (95% CI-1.1, 4.0), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the gastric cancer population in KN811 pembrolizumab combination therapy (n=350) was 4 (1.1%) (95% CI-0.3, 2.9), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the NSCLC population in KN091 pembrolizumab monotherapy (n=580) was 10 (1.7%) (95% CI- 0.8, 3.1), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785) was 10 (1.3%) (95% CI-0.6, 2.3), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the BTC population in KN966 pembrolizumab plus chemotherapy (n=529) was 3 (0.6%) (95% CI-0.1, 1.6), which is</p>
--	---

Table SVII.3.1.6: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies –Adrenal Insufficiency (Primary and Secondary)

	<p>generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the NSCLC population in KN671 pembrolizumab in combination with chemotherapy neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396) was 1 (0.3%) (95% CI-0.0, 1.4), which is generally similar to the overall proportion of patients with events of adrenal insufficiency in the reference safety dataset.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564) was 11 (2.0%) (95% CI-1.0, 3.5). which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382) was 4 (1.0%) (95% CI-0.3, 2.7), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset.</p> <p>The number and proportion of patients with adrenal insufficiency in the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528) was 2 (0.4%) (95% CI-0.0, 1.4), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260) was 2 (0.8%) (95% CI-0.1, 2.8), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset.</p> <p>The overall number and proportion of patients with adrenal insufficiency in the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361) were 7 (1.9%) (95% CI-0.8,4.0), which is generally similar to the proportion of patients with the event of adrenal insufficiency in the reference safety dataset.</p> <p><u>Pembrolizumab SC</u></p> <p>The overall number and proportion of patients with adrenal insufficiency in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251) was 5 (2%) (95% CI-0.6, 4.6), which is generally similar to the overall proportion of patients with events of adrenal insufficiency in the reference safety dataset.</p>
--	--

Table SVII.3.1.6: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies –Adrenal Insufficiency (Primary and Secondary)

Outcomes	Percent overall, fatal, not resolved, resolved, and unknown for identified risk of adrenal insufficiency (primary and secondary) - pembrolizumab. (AEs for 30 days after discontinuation were included)		
	Adrenal Insufficiency (AEOSI) KN001, KN002 and KN006 Melanoma and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment-Emergent)		
			Cumulative NSCLC 001/010 + MEL 001/002/006
		Outcome	n (%)
	Subjects in population		2,799
	With one or more adverse events	Overall	22 (0.8)
		Fatal	0 (0.0)
		Not Resolved	17 (0.6)
		Resolved	5 (0.2)
		Sequelae	0 (0.0)
	Unknown	0 (0.0)	
<p>Every subject is counted once on each applicable row. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the percent incidence specified in the report title, after rounding. Outcome: Resolved = RECOVERED/RESOLVED, Not resolved = NOT RECOVERED/NOT RESOLVED.</p>			
<p><u>Pembrolizumab IV</u></p> <p>In the NSCLC population in KN024 and KN042 (n=790), the outcome of the 4 (0.5%) patients with events of adrenal insufficiency were as follows: 2 (0.3%) resolved, 2 (0.3%) not resolved. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the NSCLC population in KN189 and KN021 (Cohorts C and G) (n=488), the outcome of the 3 patients with events of adrenal insufficiency were as follows: 1 (0.2%) resolved, 1 (0.2%) not resolved, and 1 (0.2%) unknown. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the HL population in KN013, KN087, and KN204 (n=389), the outcome for the 1 (0.3%) patient with an event of adrenal insufficiency was not recovered. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the UC population in KN052 (n=370), the outcomes for the 6 patients with events of adrenal insufficiency were as follows: 3 (0.8%) not resolved, 1 (0.3%) resolved with sequelae and 2 (0.5%) resolved and in the UC population in KN045 (n=266), the outcome for the 1 (0.4%) patient with an event of adrenal insufficiency was not resolved. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the HNSCC population in KN040, KN012 and KN055 (n=609), the outcomes for the 4 patients with events of adrenal insufficiency were as follows: 2 (0.3%) not resolved, 1 (0.2%) resolving and 1 (0.2%) resolved with sequelae. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), the outcome for the 1 (0.3%) patient with an event of adrenal insufficiency was resolved with sequelae. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the melanoma population in KN054 (n=509), the outcomes for the 5 patients with events of adrenal insufficiency were as follows: 3 (0.6%) not resolved and 2 (0.4%) resolved with sequelae. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the RCC population in KN426 pembrolizumab plus axitinib (n=429) the outcomes for the 13 patients with events of adrenal insufficiency were as follows: 2 (0.5%) resolved, 4 (0.9%) not resolved, 2 (0.5%) resolving and 5 (1.2%) resolved with sequelae. There were no patients with events of adrenal insufficiency with a fatal outcome.</p>			

Table SVII.3.1.6: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies –Adrenal Insufficiency (Primary and Secondary)

	<p>In the CRC population in KN177 (n=153), the outcomes for the 4 patients with events of adrenal insufficiency were as follows: 2 (1.3%) not resolved, 1 (0.7%) resolved, and 1 (0.7%) resolved with sequelae. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), the outcomes for the 18 patients with events of adrenal insufficiency were as follows: 15 (4.3%) not resolved, 2 (0.6%) resolving, and 1 (0.3%) resolved. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), the outcomes for the 4 patients with events of adrenal insufficiency were as follows: 1 (0.3%) resolved, 2 (0.5%) not resolved, and 1 (0.3%) resolving. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596) the outcomes for the 7 patients with events of adrenal insufficiency were as follows: 2 (0.3%) resolved, 4 (0.7%) not resolved, and 1 (0.2%) resolved with sequelae. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530) the outcomes for the 9 patients with events of adrenal insufficiency were as follows: 5 (0.9%) not resolved, 2 (0.4%) resolving, 1 (0.2%) resolved, and 1 (0.2%) resolved with sequelae. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488), the outcomes for the 10 patients with events of adrenal insufficiency were as follows: 1 (0.2%) resolved, 2 (0.4%) resolving, 3 (0.6%) not resolved and 4 (0.8%) resolved with sequelae. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), the outcomes for the 4 patients with events of adrenal insufficiency were as follows: 1 (0.3%) not resolved, 1 (0.3%) resolving, and 2 (0.7%) resolved with sequelae. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), the outcomes for the 20 patients with events of adrenal insufficiency were as follows: 1 (0.1%) resolved, 9 (1.1%) not resolved, 5 (0.6%) resolving, and 5 (0.6%) resolved with sequelae. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), the outcomes for the 11 patients with events of adrenal insufficiency were as follows: 5 (1%) not resolved, 3 (0.6%) resolving, and 3 (0.6%) resolved. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), the outcome for the 4 patients with events of adrenal insufficiency were as follows: 2 (0.6%) not resolved, 1 (0.3%) resolving and 1 (0.3%) resolved with sequelae. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580), the outcomes for the 10 patients with events of adrenal insufficiency were as follows: 1 (0.2%) resolved, 1 (0.2%) resolved with sequelae, and 8 (1.4%) not resolved. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), the outcomes for the 10 patients with events of adrenal insufficiency were as follows: 6 (0.8%) not resolved, and 4 (0.5%) resolving. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), the worst reported outcomes for the 3 patients with events of adrenal insufficiency were as follows: 1 (0.2%) resolving, 1 (0.2%) not resolved and 1 (0.2%) resolved with sequelae. There were no patients with events of adrenal insufficiency with a fatal outcome.</p>
--	--

Table SVII.3.1.6: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies –Adrenal Insufficiency (Primary and Secondary)

	<p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy group (n=396), the outcome for the 1 (0.3%) patient with an event of adrenal insufficiency was resolving. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), the outcomes for the 11 patients with events of adrenal insufficiency were as follows: 1 (0.2%) resolved and 10 (1.8%) not resolved. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382), the outcomes for patients with events of adrenal insufficiency were not recorded per study protocol (managed by external collaboration).</p> <p>In the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528), the outcome for the 2 (0.4%) patients with events of adrenal insufficiency was not resolved. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), the reported outcomes for the 2 patients with events of adrenal insufficiency were as follows: 1 (0.4%) resolved and 1 (0.4%) resolved with sequelae. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), the outcomes for the 7 patients with events of adrenal insufficiency were as follows: 4 (1.1%) resolved with sequelae, 2 (0.6%) resolved, and 1 (0.3%) not resolved. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p><u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251), the reported outcomes for the 5 patients with events of adrenal insufficiency were as follows: 3 (1.2%) not resolved and 2 (0.8%) resolved. There were no patients with events of adrenal insufficiency with a fatal outcome.</p> <p>Aggregate Review</p> <p>Review of the ASE dataset (n=7730) yielded no fatal cases of adrenal insufficiency. Review of the global safety database also yielded no fatal cases from ongoing interventional monotherapy clinical trials and no fatal cases from the PM or non-interventional environment.</p>
--	---

Table SVII.3.1.6: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies –Adrenal Insufficiency (Primary and Secondary)

Seriousness	Percent serious for identified risk of adrenal insufficiency (primary and secondary) – pembrolizumab (Serious AEs for 90 days after discontinuation were included.)	
	Adrenal Insufficiency - Serious (AEOSI)	
	KN001, KN002 and KN006 Melanoma and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab	
	(Treatment-Emergent)	
		Cumulative NSCLC 001/010 + MEL 001/002/006
		n (%)
	Subjects in population	2,799
	with one or more adverse events	10 (0.4)
	Adrenal insufficiency	8 (0.3)
	Adrenocortical insufficiency acute	1 (0.0)
Secondary adrenocortical insufficiency	1 (0.0)	
<p>Every subject is counted a single time for each applicable row and column. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns meets the incidence criterion in the report title, after rounding. Include all treated subjects in KN001 Part B1, B2, B3, D, C, F1, F2, F3 and all subjects in KN002 treated with Pembrolizumab in the original phase, and all subjects in KN006 and KN010 treated with Pembrolizumab.</p>		
<p><u>Pembrolizumab IV</u></p> <p>Of the 4 patients with events of adrenal insufficiency reported in the NSCLC population in KN024 and KN042 (n=790), 2 (0.3%) experienced serious events of adrenal insufficiency.</p> <p>Of the 3 patients with events of adrenal insufficiency reported in the NSCLC population in KN189 and KN021 (Cohorts C and G), 1 (0.2%) experienced a serious event of adrenal insufficiency.</p> <p>There were no patients with serious events of adrenal insufficiency reported in the HL population in KN013, KN087, and KN204 (n=389).</p> <p>Of the 6 (1.6%) patients with events of adrenal insufficiency reported in the UC population in KN052 (n=370), all 6 experienced serious events of adrenal insufficiency and the 1 (0.4%) patient with an event of adrenal insufficiency reported in the UC population in KN045 (n=266) experienced a serious event of adrenal insufficiency.</p> <p>Of the 4 patients with events of adrenal insufficiency reported in the HNSCC population in KN040, KN012 and KN055 (n=609), 1 (0.2%) experienced a serious event of adrenal insufficiency.</p> <p>The 1 (0.3%) patient with an event of adrenal insufficiency reported in the HNSCC population in KN048 pembrolizumab monotherapy group (n=300) experienced a serious event of adrenal insufficiency.</p> <p>Of the 5 patients with events of adrenal insufficiency reported in the melanoma population in KN054 (n=509), 1 (0.2%) experienced a serious event of adrenal insufficiency.</p> <p>Of the 13 patients with events of adrenal insufficiency in the RCC population in KN426 pembrolizumab plus axitinib (n=429), 5 (1.2%) experienced serious events of adrenal insufficiency.</p> <p>Of the 4 patients with events of adrenal insufficiency in the CRC population in KN177 (n=153), 2 (1.3%) experienced serious events of adrenal insufficiency.</p> <p>Of the 4 patients with events of adrenal insufficiency in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), 1 (0.3%) experienced a serious event of adrenal insufficiency.</p> <p>Of the 7 patients with events of adrenal insufficiency in the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), 3 (0.5%) experienced serious events of adrenal insufficiency.</p> <p>Of the 18 patients with events of adrenal insufficiency in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), 7 (2.0%) experienced serious events of adrenal insufficiency.</p>		

Table SVII.3.1.6: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies –Adrenal Insufficiency (Primary and Secondary)

	<p>Of the 9 patients with events of adrenal insufficiency in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), 6 (1.1%) experienced serious events of adrenal insufficiency.</p> <p>Of the 10 patients with events of adrenal insufficiency in the RCC population in KN564 pembrolizumab monotherapy (n=488), 6 (1.2%) experienced serious events of adrenal insufficiency.</p> <p>Of the 4 patients with events of adrenal insufficiency in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), 3 (1.0%) experienced serious events of adrenal insufficiency.</p> <p>Of the 20 patients with events of adrenal insufficiency in the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), 8 (1.0%) experienced serious events of adrenal insufficiency.</p> <p>Of the 11 patients with events of adrenal insufficiency in the melanoma population in KN716 pembrolizumab monotherapy (n=483), 4 (0.8%) experienced serious events of adrenal insufficiency.</p> <p>Of the 4 patients with events of adrenal insufficiency in gastric cancer population in KN811 pembrolizumab combination therapy (n=350), 1 (0.3%) experienced a serious event of adrenal insufficiency.</p> <p>Of the 10 patients with events of adrenal insufficiency in the NSCLC population of KN091 pembrolizumab monotherapy (n=580), 2 (0.3%) experienced serious events of adrenal insufficiency.</p> <p>Of the 10 patients with events of adrenal insufficiency in the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), 4 (0.5%) experienced serious events of adrenal insufficiency.</p> <p>Of the 3 patients with events of adrenal insufficiency in the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), 1 (0.2%) experienced a serious event of adrenal insufficiency.</p> <p>The 1 (0.3%) patient with an event of adrenal insufficiency in the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396), experienced a serious event of adrenal insufficiency.</p> <p>Of the 11 patients with events of adrenal insufficiency in the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), 2 (0.2%) experienced serious events of adrenal insufficiency.</p> <p>There were no patients with serious events of adrenal insufficiency reported in the endometrial carcinoma population in KN 868 pembrolizumab plus chemotherapy (n=382).</p> <p>There were no serious events of adrenal insufficiency in the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy.</p> <p>Of the 2 patients with events of adrenal insufficiency in the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), both experienced a serious event of adrenal insufficiency.</p> <p>Of the 7 patients with events of adrenal insufficiency reported in the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), 4 (1.1%) experienced a serious event of adrenal insufficiency.</p> <p><u>Pembrolizumab SC</u></p> <p>Of the 5 patients with an event of adrenal insufficiency in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251), 1 (0.4%) experienced a serious event of adrenal insufficiency.</p>
--	--

Table SVII.3.1.6: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies –Adrenal Insufficiency (Primary and Secondary)

Severity and Nature of the Risk	Identified treatment emergent adverse events (AE) of adrenal insufficiency (primary and secondary) (all events) in participants summarized by grade– pembrolizumab. (Table includes events for 30 days after discontinuation.)		
	Adrenal Insufficiency (AEOSI) KN001, KN002 and KN006 Melanoma and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment-Emergent)		
		Cumulative NSCLC 001/010 + MEL 001/002/006	
		n	(%)
	Subjects in population	2,799	
	with one or more adverse events	22	(0.8)
	Adrenal insufficiency	20	(0.7)
	Grade 1	3	(0.1)
	Grade 2	9	(0.3)
	Grade 3	7	(0.3)
	Grade 4	1	(0.0)
	Adrenocortical insufficiency acute	1	(0.0)
	Grade 3	1	(0.0)
	Secondary adrenocortical insufficiency	1	(0.0)
	Grade 3	1	(0.0)
	Every subject is counted a single time for each applicable specific adverse event. A subject with multiple adverse events within a system organ class is counted a single time for that system organ class. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the incidence specified in the report title, after rounding. Only the highest reported grade of a given adverse event is counted for the individual subject. Grades are based on NCI CTCAE version 4.0.		
	<u>Pembrolizumab IV</u>		
	In the NSCLC population in KN024 and KN042 (n=790), there was 1 (0.2%) patient with a Grade 3 event and 1 (0.2%) patient with a Grade 4 event of adrenal insufficiency. There were no patients with Grade 5 events of adrenal insufficiency.		
	In the NSCLC population in KN189 and KN021 (Cohorts C and G) (n=488), there was 1 (0.2%) patient with a Grade 3 event and no patients with Grade 4 or 5 events of adrenal insufficiency.		
	In the HL population in KN013, KN087, and KN204 (n=389), there were no patients with Grade 3-5 events of adrenal insufficiency.		
In the UC population in KN052 (n=370), there were 5 (1.4%) patients with Grade 3 events and 1 (0.3%) patient with a Grade 4 event of adrenal insufficiency. There were no patients with Grade 5 events of adrenal insufficiency. In the UC population in KN045 (n=266), there was 1 (0.4) patient with a Grade 3 event and patients with no Grade 4 or 5 events of adrenal insufficiency.			
In the HNSCC population in KN040, KN012 and KN055 (n=609), there was 1 (0.2%) patient with a Grade 3 event and no patients with Grade 4 or 5 events of adrenal insufficiency.			
In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), there was 1 patient with a Grade 3 event of adrenal insufficiency (0.3%). There were no patients with Grade 4 or 5 events of adrenal insufficiency.			
In the melanoma population in KN054 (n=509), there was 1 (0.2%) patient with a Grade 3 event and no patients with Grade 4 or 5 events of adrenal insufficiency.			
In the RCC population in KN426 pembrolizumab plus axitinib (n=429), there were 10 (2.3%) patients with Grade 2 events and 3 (0.7%) patients with Grade 3 events of adrenal insufficiency. There were no patients with Grade 4 or 5 events of adrenal insufficiency.			
In the CRC population in KN177 (n=153) there were 2 (1.3%) patients with Grade 3 events of adrenal insufficiency. There were no patients with Grade 4 or 5 events of adrenal insufficiency.			

Table SVII.3.1.6: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies –Adrenal Insufficiency (Primary and Secondary)

	<p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), there were 2 (0.5%) patients with Grade 3 events of adrenal insufficiency. There were no patients with Grade 4 or 5 events of adrenal insufficiency.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), there were 2 (0.3%) patients with Grade 3 events, and 1 (0.2%) patient with a Grade 4 event of adrenal insufficiency. There were no patients with Grade 5 events of adrenal insufficiency.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), there were 4 (1.1%) patients with Grade 3 events of adrenal insufficiency. There were no patients with Grade 4 or 5 events of adrenal insufficiency.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), there were 6 (1.1%) patients with Grade 3 events of adrenal insufficiency. There were no patients with Grade 4 or 5 events of adrenal insufficiency.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488) there were 6 (1.2%) patients with Grade 3 events of adrenal insufficiency. There were no patients with Grade 4 or 5 events of adrenal insufficiency.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), there were 3 (1.0%) patients with Grade 3 events of adrenal insufficiency. There were no patients with Grade 4 or 5 events of adrenal insufficiency.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) there were 7 (0.9%) patients with Grade 3 events and 1 (0.1%) patient with a Grade 4 event of adrenal insufficiency. There were no patients with Grade 5 events of adrenal insufficiency reported.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), there were 4 (0.8%) patients with Grade 3 events of adrenal insufficiency. There were no patients with Grade 4 or 5 events of adrenal insufficiency.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), there was 1 (0.3%) patient with a Grade 3 event of adrenal insufficiency. There were no patients with Grade 4 or 5 events of adrenal insufficiency.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580) there were 4 (0.7%) patients with Grade 3 events of adrenal insufficiency. There were no patients with Grade 4 or 5 events of adrenal insufficiency.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), there were 3 (0.4%) patients with a Grade 3, and 1 (0.1%) patient with a Grade 4 event of adrenal insufficiency reported. There were no patients with Grade 5 events of adrenal insufficiency.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), there was 1 (0.2%) patient with a Grade 3 event of adrenal insufficiency reported. There were no patients with Grade 4 or 5 events of adrenal insufficiency.</p> <p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396) there was 1 (0.3%) patient with a Grade 3 event of adrenal insufficiency, there were no patients with Grade 4 or 5 events of adrenal insufficiency reported.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), there were 2 (0.4%) patients with Grade 3 events of adrenal insufficiency. There were no patients with Grade 4 or 5 events of adrenal insufficiency.</p> <p>In the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382), there were no patients with Grade 3-5 events of adrenal insufficiency.</p> <p>In the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy, there were no Grade 3, 4 or 5 events of adrenal insufficiency.</p> <p>In the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), there were no patients with Grade 3, 4, or 5 events of adrenal insufficiency.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without</p>
--	--

Table SVII.3.1.6: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies –Adrenal Insufficiency (Primary and Secondary)

	<p>chemotherapy and then as monotherapy (n=361), there were 4 (1.1%) patients with a Grade 3 event of adrenal insufficiency. There were no patients with Grade 4 or 5 events of adrenal insufficiency.</p> <p><u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251) there was 1 (0.4%) patient with a Grade 3 event of adrenal insufficiency. There were no patients with Grade 4 or 5 events of adrenal insufficiency reported.</p>
<u>Risk Factors and Risk Groups:</u>	No specific risk factors for adrenal insufficiency (primary and secondary) associated with pembrolizumab have been identified.
<u>Preventability:</u>	<p>Although the development of adrenal insufficiency (primary and secondary) cannot be completely prevented; patients should be monitored for signs and symptoms of adrenal insufficiency (primary and secondary) and if they develop, other causes excluded.</p> <p>Withholding/discontinuation of pembrolizumab and appropriate medical intervention including corticosteroids following the onset of adrenal insufficiency (primary and secondary) may result in recovery.</p>
<u>Impact on the Risk-Benefit Balance of the Product:</u>	Adrenal insufficiency (primary and secondary) has the potential to become severe or life-threatening due to hormone deficiencies, in individual patients. Given the fatal outcome of untreated cancer, the risk of adrenal insufficiency (primary and secondary) is outweighed by the potential benefit.
<u>Public Health Impact:</u>	This risk has minimal public health impact outside its effect on individual patients.

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

<u>Important Identified Risk:</u>	Immune-mediated Endocrinopathies- Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)
<u>Potential Mechanisms:</u>	<p>Pembrolizumab is a potent and highly-selective humanized monoclonal antibody (mAb) of the Immunoglobulin G4 (IgG4)/kappa isotype directed to the programmed cell death-1 (PD-1) receptor and designed to directly block the interaction between the receptor and its ligands, PD-L1 and PD-L2. The PD-1 pathway represents a major immune control switch which may be engaged by tumor cells to overcome active T-cell immune surveillance. The specific mechanism of thyroid disorders associated with pembrolizumab is unknown, but it is likely that immune mediated adverse events, including thyroid disorders, is due to immune mediated reaction to certain normal tissues.</p>
<u>Evidence Source(s) and Strength of Evidence:</u>	<p>Review of pembrolizumab clinical trial data, post-marketing experience and literature regarding endocrinopathies- thyroid disorder represent sufficient evidence of a causal association with pembrolizumab exposure.</p> <p>CTD 2.7.4</p> <p>Please reference evidence source cited in Table SVII.3.1.1; Details of Important Identified Risk: Immune-Mediated Pneumonitis</p>
<u>Characterisation of the Risk:</u>	Immune-mediated Endocrinopathies- Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

Frequency with 95%CI	<p>95% Confidence Interval for the Overall Incidence (%) of Hypothyroidism, Hyperthyroidism, Thyroiditis KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th rowspan="2"></th> <th colspan="2" style="text-align: center;">Cumulative MEL 001/002/006 + NSCLC 001/010</th> </tr> <tr> <th style="text-align: center;">n (%)</th> <th style="text-align: center;">95% CI</th> </tr> </thead> <tbody> <tr> <td>Hypothyroidism</td> <td style="text-align: center;">237 (8.5)</td> <td style="text-align: center;">(7.5,9.6)</td> </tr> <tr> <td>Hyperthyroidism</td> <td style="text-align: center;">96 (3.4)</td> <td style="text-align: center;">(2.8,4.2)</td> </tr> <tr> <td>Thyroiditis</td> <td style="text-align: center;">16 (0.6)</td> <td style="text-align: center;">(0.3,0.9)</td> </tr> </tbody> </table>		Cumulative MEL 001/002/006 + NSCLC 001/010		n (%)	95% CI	Hypothyroidism	237 (8.5)	(7.5,9.6)	Hyperthyroidism	96 (3.4)	(2.8,4.2)	Thyroiditis	16 (0.6)	(0.3,0.9)
			Cumulative MEL 001/002/006 + NSCLC 001/010												
n (%)		95% CI													
Hypothyroidism	237 (8.5)	(7.5,9.6)													
Hyperthyroidism	96 (3.4)	(2.8,4.2)													
Thyroiditis	16 (0.6)	(0.3,0.9)													
	<p><u>Pembrolizumab IV</u></p> <p>In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), the overall number and proportion of patients with hypothyroidism was 39 (8%) (95% CI- 5.7, 10.8); of hyperthyroidism was 21 (4.3%) (95% CI- 2.7, 6.5); and of thyroiditis was 1 (0.2%) (95% CI- 0, 1.1), which was generally similar to the proportion of patients with the events of hypothyroidism, hyperthyroidism, and thyroiditis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hyperthyroidism in the HL population in KN013, KN087, and KN204 (n=389) was 17 (4.4%) (95%CI-2.6, 6.9); and of thyroiditis was 5 (1.3%) (95% CI-0.4, 3.0), which are generally similar to the proportion of patients with the events of hyperthyroidism and thyroiditis in the reference safety dataset for melanoma and NSCLC. The overall number and proportion of patients with hypothyroidism in the HL population in KN013, KN087, and KN204 (n=389) was 66 (17%) (95% CI-13.4, 21.1); all events of hypothyroidism were Grade 1 and 2. The rate of hypothyroidism seen in HL subjects is likely to be secondary to the significant proportion of subjects with a history of radiation to the neck and/or mediastinum, and to a higher rate of pre-existing hypothyroidism among participants in this group at baseline.</p> <p>The overall number and proportion of patients with hypothyroidism in the UC population in KN052 (n=370) was 24 (6.5%) (95%CI-4.2,9.5) and in KN045 (n=266) was 17 (6.4%) (95%CI-3.8,10); of hyperthyroidism in the UC population in KN052 (n=370) was 9 (2.4%) (95%CI-1.1,4.6) and in KN045 (n=266) was 10 (3.8%) (95%CI-1.8,6.8); and of thyroiditis in the UC population in KN052 (n=370) was 3 (0.8%) (95%CI-0.2,2.4) and in KN045 (n=266) was 2 (0.8%) (95%CI-0.1,2.7), which is generally similar to the proportion of patients with the events of hypothyroidism, hyperthyroidism, and thyroiditis in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hypothyroidism in the HNSCC population in KN040, KN012 and KN055 (n=609) was 92 (15.1%) (95% CI-12.4, 18.2); the majority of these events were Grade 1 and 2. In view of the history of these subjects and the known risk of hypothyroidism in the HNSCC population, the higher rate of hypothyroidism in the HNSCC population when compared with the reference safety dataset largely reflects disease-related risk factors. The overall number and proportion of patients with hyperthyroidism in the HNSCC population in KN040, KN012 and KN055 (n=609) was 10 (1.6%) (95% CI-0.8, 3), which is generally similar to the proportion of patients with the events of hyperthyroidism in the reference safety dataset for melanoma and NSCLC. The number and proportion of patients with thyroiditis in the HNSCC population in KN040, KN012 and KN055 (n=609) was 3 (0.5%) (95% CI-0.1, 1.4), which are generally similar to the proportion of patients with the events of hyperthyroidism in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with hypothyroidism in the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), was 54 (18%) (95% CI-13.8,22.8); of hyperthyroidism was 8 (2.7%) (95% CI-1.2,5.2); there were no cases of thyroiditis reported. The overall number and proportion of patients with hypothyroidism in the HNSCC population in KN048 pembrolizumab plus chemotherapy group (n=276), was 42 (15.2%) (95% CI-11.2,20); of</p>														

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

<p>hyperthyroidism was 13 (4.7%) (95% CI-2.5,7.9); and of thyroiditis was 1 (0.4%) (95% CI-0,2) in the HNSCC population in KN048 pembrolizumab plus chemotherapy group (n=276).The proportion of patients with events of hypothyroidism in KN048 although somewhat more common than in the reference safety dataset were generally consistent in nature with the proportion of events of hypothyroidism previously reported for pembrolizumab use in the HNSCC population.</p> <p>In the melanoma population in KN054 (n=509), the overall number and proportion of patients with events hypothyroidism was 75 (14.7%) (95% CI-11.8,18.1); of hyperthyroidism was 53 (10.4%) (95% CI-7.9,13.4); and of thyroiditis was 16 (3.1%) (95% CI-1.8, 5.1). The proportion of patients with events of hypothyroidism, hyperthyroidism and thyroiditis in KN054 although somewhat more common than in the reference safety dataset were generally consistent in nature with the proportion of events of hypothyroidism, hyperthyroidism and thyroiditis previously reported for pembrolizumab.</p> <p>The overall number and proportion of patients with hypothyroidism in the NSCLC population in KN407 and KN021 (Cohort A) (n=303), was 24 (7.9%) (95%CI-5.1, 11.6), the overall number and proportion of patients with hyperthyroidism was 20 (6.6%) (95% CI- 4.1,10), and the overall number and proportion of patients with thyroiditis was 3 (1%) (95%CI-0.2,2.9), which is generally similar to the proportion of patients with the events of hypothyroidism, hyperthyroidism, and thyroiditis in the reference safety dataset for melanoma and NSCLC.</p> <p>In the RCC population in KN426 pembrolizumab plus axitinib (n=429) the overall number and proportion of patients with hypothyroidism was 152 (35.4%) (95% CI-30.9,40.2); of hyperthyroidism was 55 (12.8%) (95% CI-9.8,16.4); and of thyroiditis was 12 (2.8%) (95% CI-1.5,4.8). Hypothyroidism is a known ADR of axitinib therapy, and axitinib contributes to the incidence of hypothyroidism in this combination. The proportion of patients with events of hypothyroidism, hyperthyroidism and thyroiditis in KN426 although more common than in the reference safety dataset were generally consistent in nature with the proportion of events of hypothyroidism, hyperthyroidism and thyroiditis previously reported for pembrolizumab.</p> <p>In the in the CRC population in KN177 (n=153) the overall number and proportion of patients with hypothyroidism was 19 (12.4%) (95% CI-7.6,18.7); of hyperthyroidism was 6 (3.9%) (95% CI-1.5,8.3); and of thyroiditis was 2 (1.3%) (95% CI-0.2,4.6). Hypothyroidism was reported more frequently in the KN177 pembrolizumab safety dataset compared with the RSD (12.4% vs 8.5%, respectively); however, when adjusted for exposure, the incidences were similar. The proportion of patients with events of hypothyroidism, hyperthyroidism and thyroiditis in KN177 were generally consistent in nature with the proportion of events of hypothyroidism, hyperthyroidism and thyroiditis previously reported for pembrolizumab.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), the overall number and proportion of patients with hypothyroidism was 40 (10.8%) (95% CI-7.8,14.4); of hyperthyroidism was 21 (5.7%) (95% CI-3.5,8.5); and of thyroiditis 1 (0.3%) (95% CI-0,1.5), which are generally similar to the proportion of patients with the events of hypothyroidism, hyperthyroidism, and thyroiditis in the reference safety dataset for melanoma and NSCLC.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596) the overall number and proportion of patients with hyperthyroidism was 31 (5.2%) (95% CI-3.6,7.3); and of thyroiditis was 8 (1.3%) (95% CI-0.6,2.6), which are generally similar to the proportion of patients with the events of hyperthyroidism and thyroiditis in the reference safety dataset for melanoma and NSCLC. The overall number and proportion of patients with hypothyroidism was 95 (15.9%) (95% CI-13.1,19.1), although higher than in the reference safety dataset, the events were generally consistent in nature with the proportion of events of hypothyroidism previously reported for pembrolizumab.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), the number and proportion of patients with thyroiditis was 2 (0.6%) (95% CI-0.1,2.0), which is generally similar to the proportion of patients with the events of thyroiditis in the reference safety dataset. The overall number and proportion of patients with hypothyroidism was 166 (47.2%) (95% CI- 41.8,52.5), which although higher than hypothyroidism in the reference safety dataset, was generally consistent in nature with the proportion of events of hypothyroidism previously reported for pembrolizumab. The overall number and proportion of patients with hyperthyroidism was 28 (8.0%) (95% CI-</p>

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

	<p>5.4,11.3), which although higher than hyperthyroidism in the reference safety dataset, was generally consistent in nature with the proportion of events of hyperthyroidism previously reported for pembrolizumab.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530) the overall number and proportion of patients with thyroiditis was 12 (2.3%) (95% CI-1.2, 3.9), which is generally similar to the proportion of patients with the event of thyroiditis in the reference safety dataset. The overall number and proportion of patients with hypothyroidism was 296 (55.8%) (95% CI-51.5, 60.1); and of hyperthyroidism was 54 (10.2%) (95% CI-7.7, 13.1). The proportion of patients with events of hypothyroidism and hyperthyroidism in KN146 and KN775, although higher in frequency than in the reference safety dataset, were generally consistent in nature with the proportion of events of hypothyroidism and hyperthyroidism previously reported for pembrolizumab.</p> <p>The overall number and proportion of patients with hypothyroidism in the RCC population in KN564 pembrolizumab monotherapy (n=488) was 103 (21.1%) (95% CI-17.6,25); of hyperthyroidism was 58 (11.9%) (95% CI- 9.1,15.1). The overall number and proportion of patients with thyroiditis was 6 (1.2%) (95% CI-0.5,2.7), which was generally consistent to the overall proportion of patients with the events of hypothyroidism, hyperthyroidism, and thyroiditis in the reference safety dataset. The proportion of patients with events of hypothyroidism and hyperthyroidism in KN564 although more common than in the reference safety dataset were generally consistent in nature with the proportion of events of hypothyroidism and hyperthyroidism previously reported for pembrolizumab.</p> <p>In the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), the overall number and proportion of patients with hypothyroidism was 53 (11.2%) (95% CI-8.5, 14.3) and of hyperthyroidism was 25 (5.3%) (95% CI-3.4, 7.7), which is generally similar to the proportion of patients with the events of hypothyroidism and hyperthyroidism in the reference safety dataset. There were no patients with events of thyroiditis reported in the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), the overall number and proportion of patients with hypothyroidism was 56 (18.2%) (95% CI-14.1, 23.0); of hyperthyroidism was 23 (7.5%) (95% CI-4.8, 11.0); and of thyroiditis 11 (3.6%) (95% CI-1.8, 6.3). The proportion of patients with events of hypothyroidism, hyperthyroidism and thyroiditis although more common than in the reference safety dataset were generally consistent in nature with the proportion of events of hypothyroidism, hyperthyroidism and thyroiditis previously reported for pembrolizumab.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) the number and proportion of patients with hyperthyroidism was 41 (5.2%) (95% CI-3.8,7.0); and of thyroiditis was 16 (2.0%) (95% CI- 1.2,3.3), which are generally consistent to the proportion of patients with events of hyperthyroidism and thyroiditis in the reference safety dataset. The overall number and proportion of patients with hypothyroidism was 118 (15.1%) (95% CI-12.6,17.8), although higher than in the reference safety dataset, the proportion of patients with events were generally consistent in nature with the proportion of events of hypothyroidism previously reported for pembrolizumab.</p> <p>In the melanoma population in KN716 (n=483), the overall number and proportion of patients with hypothyroidism was 76 (15.7%) (95% CI-12.6, 19.3); of hyperthyroidism was 50 (10.4%) (95% CI-7.8, 13.4); of thyroiditis was 8 (1.7%) (95% CI-0.7, 3.2). The proportion of patients with events of hypothyroidism and hyperthyroidism in KN716 although more common than in the reference safety dataset were generally consistent in nature with the proportion of events of hypothyroidism and hyperthyroidism previously reported for pembrolizumab in advanced cancer.</p> <p>The overall number and proportion of patients with hypothyroidism in the gastric cancer population in KN811 pembrolizumab combination therapy (n=350) was 37 (10.6%) (95% CI-7.6, 14.3); with hyperthyroidism was 14 (4.0%) (95% CI-2.2, 6.6); and with thyroiditis was 4 (1.1%) (95% CI-0.3, 2.9), which were generally similar to the proportion of patients with events of hypothyroidism, hyperthyroidism, and thyroiditis in the reference safety dataset.</p> <p>The overall number and proportion of patients with hypothyroidism in the NSCLC population in KN091 pembrolizumab monotherapy (n=580) was 120 (20.7%) (95% CI-17.5,24.2); of</p>
--	---

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

<p>hyperthyroidism was 62 (10.7%) (95% CI-8.3,13.5). The proportion of patients with events of hypothyroidism and hyperthyroidism in KN091 although more common than in the reference safety dataset, was generally consistent with the proportion of events of hypothyroidism and hyperthyroidism reported in adjuvant studies with pembrolizumab monotherapy. The proportion of patients with events of thyroiditis was 6 (1%) (95% CI-0.4,2.2) in KN091 which was similar to the proportion of patients with the events of thyroiditis in the reference safety dataset.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), the overall number and proportion of patients with hypothyroidism was 120 (15.3%) (95% CI-12.8, 18.0); of hyperthyroidism was 44 (5.6%) (95% CI-4.1, 7.5); and of thyroiditis was 9 (1.1%) (95% CI-0.5, 2.2). The proportion of patients with events of hypothyroidism, although more common than in the reference safety dataset, were generally consistent in nature with the proportion of patients with events of hypothyroidism previously reported for pembrolizumab. The proportions of patients with hyperthyroidism and thyroiditis were generally similar to the overall frequency in the reference safety dataset.</p> <p>In the BTC population in KN966 (n=529) pembrolizumab plus chemotherapy, the overall number and proportion of patients with hypothyroidism was 46 (8.7%) (95% CI-6.4, 11.4); of hyperthyroidism was 19 (3.6%) (95% CI 2.2, 5.6); and thyroiditis was 3 (0.6%) (95% CI-0.1, 1.6), which is generally similar to the proportion of patients with the events of hypothyroidism, hyperthyroidism, and thyroiditis in the reference safety dataset.</p> <p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396) the overall number and proportion of patients with hypothyroidism was 44 (11.1%) (95% CI-8.2, 14.6), with hyperthyroidism was 22 (5.6%) (95% CI-3.5, 8.3), and with thyroiditis was 4 (1.0) (95% CI 0.3, 2.6). The events of hypothyroidism and hyperthyroidism in KN671 were more common than in the reference safety dataset. However, after adjusting for exposure, the rates for hypothyroidism and hyperthyroidism were similar to those reported in the reference dataset. The overall number and proportion of patients with thyroiditis were similar to the reference safety dataset.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), the overall number and proportion of patients with hypothyroidism was 60 (10.6%) (95% CI-8.2, 13.5); of hyperthyroidism was 25 (4.4%) (95% CI-2.9, 6.5); and of thyroiditis was 5 (0.9%) (95% CI-0.3, 2.1). The proportions of patients with hypothyroidism, hyperthyroidism and thyroiditis were generally similar to the overall frequency in the reference safety dataset.</p> <p>In the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), the overall number and proportion of patients with hypothyroidism was 24 (9.2%) (95% CI-6, 13.4); of hyperthyroidism was 4 (1.5%) (95% CI-0.4, 3.9), which is generally similar to the proportion of patients with the events of hypothyroidism and hyperthyroidism in the reference safety dataset. There were no patients with events of thyroiditis reported in the mesothelioma population in KN483 and KNA17.</p> <p>The overall number and proportion of patients with hypothyroidism in the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), was 89 (24.7%) (95% CI-20.3,29.4); of hyperthyroidism was 32 (8.9%) (95% CI-6.1,12.3); and of thyroiditis was 2 (0.6%) (95% CI-0.1,2.0). The proportion of patients with events of hypothyroidism and hyperthyroidism in KN689 although somewhat more common than in the reference safety dataset were generally consistent with the proportion of events of hypothyroidism and hyperthyroidism previously reported for pembrolizumab use in the HNSCC population. The rate of thyroiditis was similar to the reference safety dataset.</p> <p><u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251) the overall number and proportion of patients with hypothyroidism was 35 (13.9%) (95% CI-9.9, 18.9), with hyperthyroidism was 20 (8%) (95% CI-4.9, 12.0), and with thyroiditis was 1 (0.4) (95% CI 0.0, 2.2). The events of hypothyroidism and hyperthyroidism in MK-3475A-D77 were more common than in the reference safety dataset. However, after adjusting for exposure, the rates for hypothyroidism and</p>
--

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

	hyperthyroidism were similar to those reported in the reference dataset. The overall number and proportion of patients with thyroiditis were similar to the reference safety dataset.																																				
Outcomes	<p>Percent overall, fatal, not resolved, resolved, and unknown for identified risk of thyroid disorder– pembrolizumab (Table includes events for 30 days after discontinuation.)</p> <p style="text-align: center;">Hypothyroidism (AEOSI)</p> <p style="text-align: center;">KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment Emergent).</p> <table border="1" data-bbox="483 562 1412 877"> <thead> <tr> <th></th> <th></th> <th colspan="2">Cumulative NSCLC 001/010 + MEL 001/002/006</th> </tr> <tr> <th></th> <th>Outcome</th> <th>n</th> <th>(%)</th> </tr> </thead> <tbody> <tr> <td>Subjects in population</td> <td></td> <td>2,799</td> <td></td> </tr> <tr> <td rowspan="7">With one or more adverse events</td> <td>Overall</td> <td>237</td> <td>(8.5)</td> </tr> <tr> <td>Fatal</td> <td>0</td> <td>(0.0)</td> </tr> <tr> <td>Not Resolved</td> <td>176</td> <td>(6.3)</td> </tr> <tr> <td>Resolved</td> <td>46</td> <td>(1.6)</td> </tr> <tr> <td>Resolving</td> <td>6</td> <td>(0.2)</td> </tr> <tr> <td>Sequelae</td> <td>2</td> <td>(0.1)</td> </tr> <tr> <td>Unknown</td> <td>7</td> <td>(0.3)</td> </tr> </tbody> </table> <p>Every subject is counted once on each applicable row. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the percent incidence specified in the report title, after rounding. Outcome: Resolved = RECOVERED/RESOLVED, Not resolved = NOT RECOVERED/NOT RESOLVED.</p> <p><u>Pembrolizumab IV</u></p> <p>In the NSCLC population in KN024 and KN042 (n=790), the outcomes of the 93 patients with events of hypothyroidism were as follows: 25 (3.2%) resolved, 12 (1.5%) resolving, 52 (6.6%) not resolved, 2 (0.3%) resolved with sequelae, 2 (0.3%) unknown. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), the outcomes of the 39 patients with events of hypothyroidism were as follows: 12 (2.5%) resolved, 5 (1%) resolving, 21 (4.3%) not resolved, 1 (0.2%) unknown. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the HL population in KN013, KN087, and KN204 (n=389), the outcomes of the 66 patients with events of hypothyroidism were as follows: 27 (6.9%) resolved, 8 (2.1%) resolving, 27 (6.9%) not resolved, and 4 (1.0%) resolved with sequelae. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the UC population in KN052 (n=370), the outcomes for the 24 patients with events of hypothyroidism were as follows: 1 (0.3%) resolved, 3(0.8%) resolving, 17 (4.6%) not resolved, and 3 (0.8%) unknown and in KN045 (n=266), the outcomes for the 17 patients with events of hypothyroidism were as follows: 4 (1.5%) resolved, 2 (0.8%) resolved with sequelae, 11 (4.1%) not resolved. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the HNSCC population in KN040, KN012 and KN055 (n=609), the outcomes of the 92 patients with events of hypothyroidism were as follows: 12 (2.0%) resolved, 8 (1.3%) resolving, 66 (10.8%) not resolved and 6 (1.0%) unknown. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), the outcomes of the 54 (18%) patients with events of hypothyroidism were as follows: 14 (4.7%) resolved, 12 (4.0%) resolving, 27 (9.0%) not resolved and 1 (0.3%) unknown. There were no fatal outcomes. In the HNSCC population in KN048 pembrolizumab plus chemotherapy group (n=276), the outcomes of the 42 (15.2%) patients with events of hypothyroidism were as follows: 10 (3.6%) resolved, 5</p>					Cumulative NSCLC 001/010 + MEL 001/002/006			Outcome	n	(%)	Subjects in population		2,799		With one or more adverse events	Overall	237	(8.5)	Fatal	0	(0.0)	Not Resolved	176	(6.3)	Resolved	46	(1.6)	Resolving	6	(0.2)	Sequelae	2	(0.1)	Unknown	7	(0.3)
		Cumulative NSCLC 001/010 + MEL 001/002/006																																			
	Outcome	n	(%)																																		
Subjects in population		2,799																																			
With one or more adverse events	Overall	237	(8.5)																																		
	Fatal	0	(0.0)																																		
	Not Resolved	176	(6.3)																																		
	Resolved	46	(1.6)																																		
	Resolving	6	(0.2)																																		
	Sequelae	2	(0.1)																																		
	Unknown	7	(0.3)																																		

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

	<p>(1.8%) resolving and 27 (9.8%) not resolved. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the melanoma population in KN054 (n=509), the outcomes of the 75 patients with events of hypothyroidism were as follows: 20 (3.9%) resolved, 2 (0.4%) resolved with sequelae, and 53 (10.4%) not resolved. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the NSCLC population in KN407 and KN021 (Cohort A) (n=303), the outcomes of the 24 patients with events of hypothyroidism were as follows: 13 (4.3%) not resolved, 8 (2.6%) resolving, and 3 (1.0%) resolved. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the RCC population in KN426 pembrolizumab plus axitinib (n=429), the outcomes for the 152 patients with events of hypothyroidism were as follows: 38 (8.9%) resolved, 95 (22.1%) not resolved, and 19 (4.4%) resolving. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the CRC population in KN177 (n=153), the outcomes for the 19 patients with events of hypothyroidism were as follows: 5 (3.3%) resolved, 1 (0.7%) resolving and 13 (8.5%) not resolved. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), the outcomes for the 40 patients with events of hypothyroidism were as follows: 11 (3%) resolved, 23 (6.2%) not resolved, 5 (1.4%) resolving and 1 (0.3%) resolved with sequelae. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), the outcomes for the 95 patients with events of hypothyroidism were as follows: 26 (4.4%) resolved, 47 (7.9%) not resolved, 21 (3.5%) resolving, and 1 (0.2%) resolved with sequelae. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), the outcomes for the 166 patients with events of hypothyroidism were as follows: 118 (33.5%) not resolved, 33 (9.4%) resolved, 14 (4.0%) resolving, and 1 (0.3%) resolved with sequelae. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), the outcomes for the 296 patients with events of hypothyroidism were as follows: 197 (37.2%) not resolved, 54 (10.2%) resolved, 38 (7.2%) resolving, 6 (1.1%) resolved with sequelae, and 1 (0.2%) unknown. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488), the outcomes for the 103 patients with events of hypothyroidism were as follows: 15 (3.1%) resolved, 60 (12.3%) not resolved, 26 (5.3%) resolving, and 2 (0.4%) resolved with sequelae. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), the outcomes for the 53 (11.2%) patients with events of hypothyroidism were as follows: 15 (3.2%) resolved, 8 (1.7%) resolving, 1 (0.2%) unknown and 29 (6.1%) not resolved. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), the outcomes for the 56 patients with events of hypothyroidism were as follows: 32 (10.4%) not resolved, 16 (5.2%) resolving, and 8 (2.6%) resolved. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), the outcomes for the 118 patients with events of hypothyroidism were as follows: 33 (4.2%) resolved, 64 (8.2%) not resolved, 19 (2.4%) resolving, and 2 (0.3%) resolved with sequelae. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), the outcomes for the 76 patients with events of hypothyroidism were as follows: 43 (8.9%) not resolved, 24 (5%) resolving, and 9 (1.9%) resolved. There were no patients with events of hypothyroidism with a fatal outcome.</p>
--	--

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

	<p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), the outcome for the 37 patients with events of hypothyroidism were as follows: 17 (4.9%) not resolved, 12 (3.4%) resolved, and 8 (2.3%) resolving. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580), the outcomes for the 120 patients with events of hypothyroidism were as follows: 21 (3.6%) resolved and 99 (17.1%) not resolved. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), the outcomes for the 120 patients with events of hypothyroidism were as follows: 69 (8.8%) not resolved, 28 (3.6%) resolving, 20 (2.5%) resolved, and 3 (0.4%) resolved with sequelae. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), the worst reported outcomes for the 46 patients with events of hypothyroidism were as follows: 26 (4.9%) not resolved, 14 (2.6%) resolving and 6 (1.1%) resolved. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy group (n=396), the outcome for the 44 patients with events of hypothyroidism were as follows: 26 (6.6%) not resolved, 8 (2.0%) resolving, 9 (2.3%) resolved, and 1 (0.3%) resolved with sequelae. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), the outcomes for the 60 patients with events of hypothyroidism were as follows: 47 (8.3%) not resolved, 1 (0.2%) resolving, 11 (2.0%) resolved, and 1 (0.2%) resolved with sequelae. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), the reported outcomes for the 24 patients with events of hypothyroidism were as follows: 18 (6.9%) not resolved and 6 (2.3%) resolved. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), the outcomes of the 89 patients with events of hypothyroidism were as follows: 47 (13%) not resolved, 19 (5.26%) resolving, 17 (4.7%) resolved, 5 (5.6%) resolved with sequelae, and 1 (1.4%) unknown. There were no patients with events of hypothyroidism with a fatal outcome.</p> <p><u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251), the outcome for the 35 patients with events of hypothyroidism were as follows: 19 (7.6%) not resolved, 10 (4.0%) resolving, and 5 (2.0%) resolved. There were no patients with events of hypothyroidism with a fatal outcome.</p>
--	---

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

Hyperthyroidism (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment Emergent)			
		Cumulative NSCLC 001/010 + MEL 001/002/006	
	Outcome	n	(%)
Subjects in population		2,799	
With one or more adverse events	Overall	96	(3.4)
	Fatal	0	(0.0)
	Not Resolved	23	(0.8)
	Resolved	71	(2.5)
	Unknown	2	(0.1)
<p>Every subject is counted once on each applicable row. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the percent incidence specified in the report title, after rounding. Outcome: Resolved = RECOVERED/RESOLVED, Not resolved = NOT RECOVERED/NOT RESOLVED.</p>			
<p><u>Pembrolizumab IV</u></p> <p>In the NSCLC population in KN024 and KN042 (n=790), the outcomes of the 50 patients with events of hyperthyroidism were as follows: 40 (5.1%) resolved, 3 (4%) resolving (0.9%) not resolved. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), the outcomes of the 21 patients with events of hyperthyroidism were as follows: 16 (3.3%) resolved, 3 (0.6%) resolving, 2 (0.4%) not resolved. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the HL population in KN013, KN087, and KN204 (n=389), the outcomes of the 17 patients with events of hyperthyroidism were as follows: 15 (3.9%) resolved, 1 (0.3%) not resolved, and 1 (0.3%) resolved with sequelae. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the UC population in KN052 (n=370), the outcomes for the 9 patients with events of hyperthyroidism were as follows: 6 (1.6%) resolved, 1 (0.3%) resolving, 1 (0.3%) not resolved, and 1 (0.3%) unknown and in KN045 (n=266), the outcomes for the 10 patients with events of hyperthyroidism were as follows: 7(2.6%) resolved, 2 (0.8%) not resolved, and 1 (0.4%) unknown. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the HNSCC population in KN040, KN012 and KN055 (n=609), the outcomes of the 10 patients with events of hyperthyroidism were as follows: 6 (1.0%) resolved, 2 (0.3%) resolving, 1 (0.2%) not resolved and 1(0.2%) unknown. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), the outcomes of the 8 (2.7%) patients with events of hyperthyroidism were as follows: 4 (1.3%) resolved, 1 (0.3%) resolving, 1 (0.3%) resolved with sequelae, and 2 (0.7%) not resolved. There were no fatal outcomes. In the HNSCC population in KN048 pembrolizumab plus chemotherapy group (n=276), the outcomes of the 13 (4.7%) patients with events of hyperthyroidism were as follows: 9 (3.3%) resolved, 2 (0.7%) resolving and 2 (0.7%) not resolved. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the melanoma population in KN054 (n=509), the outcomes of the 53 patients with events of hyperthyroidism were as follows: 44 (8.6%) resolved, 6 (1.2%) not resolved and 3 (0.6%) recovered with sequelae. There were no patients with events of hyperthyroidism with a fatal outcome.</p>			

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

	<p>In the NSCLC population in KN407 and KN021 (Cohort A) (n=303), the outcomes of the 20 patients with events of hyperthyroidism were as follows: 12 (4.0%) resolved, 5 (1.7%) not resolved, 3 (1.0%) resolving. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the RCC population in KN426 pembrolizumab plus axitinib (n=429) the outcomes for the 55 patients with events of hyperthyroidism were as follows: 47 (11.0%) resolved, 7 (1.6%) not resolved, and 1 (0.2%) resolving. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the CRC population in KN177 (n=153), the outcomes for the 6 patients with events of hyperthyroidism were as follows: 5 (3.3%) resolved and 1 (0.7%) not resolved. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), the outcomes for the 21 patients with events of hyperthyroidism were as follows: 14 (3.8%) resolved, 6 (1.6%) not resolved, and 1 (0.3%) resolving. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596) the outcomes for the 31 patients with events of hyperthyroidism were as follows: 21 (3.5%) resolved, 7 (1.2%) not resolved, 1 (0.2%) resolving, 1 (0.2%) resolved with sequelae and 1 (0.2%) unknown. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), the outcomes for the 28 patients with events of hyperthyroidism were as follows: 20 (5.7%) resolved, 4 (1.1%) resolving, 3 (0.9%) not resolved, 1 (0.3%) resolved with sequelae. There were no events of hyperthyroidism with a fatal outcome.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530) the outcomes for the 54 patients with events of hyperthyroidism were as follows: 46 (8.7%) resolved, 7 (1.3%) not resolved, and 1 (0.2%) resolved with sequelae. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488), the outcomes for the 58 patients with events of hyperthyroidism were as follows: 47 (9.6%) resolved, 3 (0.6%) not resolved, 3 (0.6%) resolving, and 5 (1%) resolved with sequelae. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), the outcomes for the 25 (5.3%) patients with events of hyperthyroidism were as follows: 21 (4.4%) resolved, 1 (0.2%) resolved with sequelae, and 3 (0.6%) not resolved. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), the outcomes for the 23 patients with events of hyperthyroidism were as follows: 19 (6.2%) resolved, 3 (1.0%) not resolved and 1 (0.3%) resolved with sequelae. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), the outcomes for the 41 patients with events of hyperthyroidism were as follows: 35 (4.5%) resolved, 3 (0.4%) not resolved, 2 (0.3%) resolving, and 1 (0.1%) resolved with sequelae. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), the outcomes for the 50 patients with events of hyperthyroidism were as follows: 43 (8.9%) resolved, 5 (1%) not resolved, 1 (0.2%) resolving, and 1 (0.2%) resolved with sequelae. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), the outcome for the 14 patients with events of hyperthyroidism were as follows: 6 (1.7%) resolved, 6 (1.7%) not resolved and 2 (0.6%) resolving. There were no patients with events of hyperthyroidism with a fatal outcome.</p>
--	--

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

	<p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580), the outcomes for the 62 patients with events of hyperthyroidism were as follows: 60 (10.3%) resolved and 2 (0.3%) not resolved. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), the outcomes for the 44 patients with events of hyperthyroidism were as follows: 33 (4.2%) resolved, 7 (0.9%) not resolved, 2 (0.3%) resolving, and 2 (0.3%) resolved with sequelae. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), the worst reported outcomes for the 19 patients with events of hyperthyroidism were as follows: 15 (2.8%) resolved, 2 (0.4%) not resolved and 2 (0.4%) resolving. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy group (n=396), the outcome for the 22 patients with events of hyperthyroidism were as follows: 3 (0.8%) not resolved, 2 (0.5%) resolving, and 17 (4.3%) resolved. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), the outcome for the 25 patients with events of hyperthyroidism were as follows: 20 (3.5%) resolved, 4 (0.7%) not resolved and 1 (0.2%) resolved with sequelae. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), the reported outcomes for the 4 patients with events of hyperthyroidism were as follows: 3 (1.2%) resolved and 1 (0.4%) not resolved. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), the outcomes of the 32 patients with events of hyperthyroidism were as follows: 23 (6.4%) resolved, 6 (1.7%) not resolved, and 3 (0.8%) resolving. There were no patients with events of hyperthyroidism with a fatal outcome.</p> <p><u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251), the outcome for the 20 patients with events of hyperthyroidism were as follows: 14 (5.6%) resolved, 4 (1.6%) not resolved, and 2 (0.8%) resolving. There were no patients with events of hyperthyroidism with a fatal outcome.</p>
--	---

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

Thyroiditis (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment Emergent)			
		Cumulative NSCLC 001/010 + MEL 001/002/006	
	Outcome	n	(%)
Subjects in population		2,799	
With one or more adverse events	Overall	16	(0.6)
	Fatal	0	(0.0)
	Not Resolved	8	(0.3)
	Resolved	7	(0.3)
	Resolving	1	(0.0)
	Unknown	0	(0.0)

Every subject is counted once on each applicable row.
A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the percent incidence specified in the report title, after rounding.
Outcome: Resolved = RECOVERED/RESOLVED, Not resolved = NOT RECOVERED/NOT RESOLVED.

Pembrolizumab IV

In the NSCLC population in KN024 and KN042 (n=790), the outcomes of the 14 patients with events of thyroiditis were as follows: 7 (0.9%) not resolved, 6 (0.8%) resolved and 1 (0.1%) recovered with sequelae. There were no patients with events of thyroiditis with a fatal outcome.

In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), the outcome of the 1 (0.2%) patient with an event of thyroiditis was not resolved.

In the HL population in KN013, KN087, and KN204 (n=389), the outcomes of the 5 patients with events of thyroiditis were as follows: 2 (0.5%) not resolved, 2 (0.5%) resolved, and 1 (0.3%) recovered with sequelae. There were no patients with events of thyroiditis with a fatal outcome.

In the UC population in KN052 (n=370), the outcomes for the 3 patients with events of thyroiditis were as follows: 1 (0.3%) resolving and 2 (0.5%) not resolved and in KN045 (n=266), the outcomes for the 2 patients with events of thyroiditis were as follows: 2 (0.8%) resolving. There were no patients with events of thyroiditis with a fatal outcome.

In the HNSCC population in KN040, KN012 and KN055 (n=609), the outcomes of the 3 patients with events of thyroiditis were as follows: 1 (0.2%) resolved, 1 (0.2%) resolving and 1 (0.2%) not resolved. There were no patients with events of thyroiditis with a fatal outcome.

In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), there were no events of thyroiditis. In the HNSCC population in KN048 pembrolizumab plus chemotherapy group (n=276), the outcome of the 1 (0.4%) patient with an event of thyroiditis was resolved. There were no patients with events of thyroiditis with a fatal outcome.

In the melanoma population in KN054 (n=509), the outcomes of the 16 patients with events of thyroiditis were as follows: 6 (1.2%) not resolved, 9 (1.8%) resolved, and 1(0.2%) recovered with sequelae. There were no patients with events of thyroiditis with a fatal outcome.

In the NSCLC population in KN407 and KN021 (Cohort A) (n=303), the outcomes of the 3 patients with events of thyroiditis were as follows: 2 (0.7%) not resolved, 1 (0.3%) resolved. There were no patients with events of thyroiditis with a fatal outcome.

In the RCC population in KN426 pembrolizumab plus axitinib (n=429) the outcomes for the 12 patients with events of thyroiditis were as follows: 4 (0.9%) resolved, 7 (1.6%) not resolved, and 1 (0.2%) resolving. There were no patients with events of thyroiditis with a fatal outcome.

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

	<p>In the CRC population in KN177 (n=153), the outcomes for the 2 patients with events of thyroiditis were as follows: 1 (0.7%) resolved and 1 (0.7%) resolved with sequelae. There were no patients with events of thyroiditis with a fatal outcome.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), the outcome for the 1 patient with an event of thyroiditis was as follows: 1 (0.3%) not resolved. There were no patients with events of thyroiditis with a fatal outcome.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), the outcomes for the 8 patients with events of thyroiditis were as follows: 4 (0.7%) resolved, 2 (0.3%) not resolved, and 2 (0.3%) resolving. There were no fatal outcomes.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), the outcomes for the 2 patients with events of thyroiditis there were as follows: 1 (0.3%) resolved, and 1 (0.3%) not resolved. There were no patients with events of thyroiditis with a fatal outcome.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530) the outcomes for the 12 patients with events of thyroiditis were as follows: 6 (1.1%) not resolved, 4 (0.8%) resolved, 1 (0.2%) resolved with sequelae, and 1 (0.2%) resolving. There were no patients with events of thyroiditis with a fatal outcome.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488), the outcomes for the 6 patients with events of thyroiditis were as follows: 3 (0.6%) resolved and 3 (0.6%) not resolved. There were no patients with events of thyroiditis with a fatal outcome.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), the outcomes for the 11 patients with events of thyroiditis were as follows: 6 (2.0%) not resolved, 4 (1.3%) resolved and 1 (0.3%) resolving. There were no patients with events of thyroiditis with a fatal outcome.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), the outcomes for the 16 patients with events of thyroiditis were as follows: 7 (0.9%) resolved, 8 (1.0%) not resolved, and 1 (0.1%) resolved with sequelae. There were no patients with events of thyroiditis with a fatal outcome.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), the outcome for the 8 patients with events of thyroiditis was as follows: 5 (1%) not resolved, 2 (0.4%) resolving, and 1 (0.2%) resolved. There were no patients with events of thyroiditis with a fatal outcome.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), the outcome for the 4 patients with events of thyroiditis were as follows: 2 (0.6%) not resolved, 1 (0.3%) resolving and 1 (0.3%) resolved. There were no patients with events of thyroiditis with a fatal outcome.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580), the outcomes for the 6 patients with events of thyroiditis were as follows: 1 (0.2%) resolved, 1 (0.2%) resolved with sequelae, and 4 (0.7%) not resolved. There were no patients with events of thyroiditis with a fatal outcome.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), the outcomes for the 9 patients with events of thyroiditis were as follows: 3 (0.4%) not resolved, 3 (0.4%) resolving, 2 (0.3%) resolved, and 1 (0.1%) resolved with sequelae. There were no patients with events of thyroiditis with a fatal outcome.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), the worst reported outcomes for the 3 (0.6%) patients with events of thyroiditis were not resolved. There were no patients with events of thyroiditis with a fatal outcome.</p> <p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy group (n=396), the outcome for the 4 patients with events of thyroiditis were as follows: 3 (0.8%) not resolved, and 1 (0.3%) resolved. There were no patients with events of thyroiditis with a fatal outcome.</p>
--	---

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

	<p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), the outcomes for the 5 patients with events of thyroiditis were as follows: 4 (0.7%) not resolved and 1 (0.2%) resolved. There were no patients with events of thyroiditis with a fatal outcome.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), the outcomes of the 2 patients with events of thyroiditis was 1 (0.3%) resolved and 1 (0.3%) resolving. There were no patients with events of thyroiditis with a fatal outcome.</p> <p><u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251), the outcome for the 1 (0.4%) patient with thyroiditis was not resolved. There were no patients with events of thyroiditis with a fatal outcome.</p> <p>Aggregate Review</p> <p>Review of the ASE dataset (n=7730) yielded no fatal cases of thyroid disorders. Review of the global safety database yielded the following: no fatal cases from ongoing interventional monotherapy clinical trials and 1 fatal case of thyroiditis and 1 fatal case of hypothyroidism reported from the PM or non-interventional environment. In both cases, the thyroid disorder event was the only Grade 5 event.</p>																		
<p>Seriousness</p>	<p>Percent serious for identified risk of thyroid disorder - pembrolizumab. (Serious AEs for 90 days after discontinuation were included.)</p> <p style="text-align: center;">Hypothyroidism - Serious (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment-Emergent)</p> <table border="1" data-bbox="483 1136 1406 1381"> <thead> <tr> <th></th> <th colspan="2">Cumulative NSCLC 001/010 + MEL 001/002/006</th> </tr> <tr> <th></th> <th>n</th> <th>(%)</th> </tr> </thead> <tbody> <tr> <td>Subjects in population</td> <td>2,799</td> <td></td> </tr> <tr> <td>with one or more adverse events</td> <td>5</td> <td>(0.2)</td> </tr> <tr> <td></td> <td></td> <td></td> </tr> <tr> <td>Hypothyroidism</td> <td>5</td> <td>(0.2)</td> </tr> </tbody> </table> <p>Every subject is counted a single time for each applicable row and column. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns meets the incidence criterion in the report title, after rounding. Include all treated subjects in KN001 Part B1, B2, B3, D, C, F1, F2, F3 and all subjects in KN002 treated with Pembrolizumab in the original phase and all subjects in KN006 and KN010 treated with Pembrolizumab.</p> <p><u>Pembrolizumab IV</u></p> <p>Of the 93 patients with events of hypothyroidism in the NSCLC population in KN024 and KN042 (n=790), 1 (0.1%) experienced a serious event of hypothyroidism.</p> <p>Of the 39 patients with events of hypothyroidism in the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), 1 (0.2%) of the events experienced a serious event of hypothyroidism.</p> <p>Of the 66 patients with events of hypothyroidism in the HL population in KN013, KN087, and KN204 (n=389), none of the patients experienced events serious events of hypothyroidism.</p>		Cumulative NSCLC 001/010 + MEL 001/002/006			n	(%)	Subjects in population	2,799		with one or more adverse events	5	(0.2)				Hypothyroidism	5	(0.2)
	Cumulative NSCLC 001/010 + MEL 001/002/006																		
	n	(%)																	
Subjects in population	2,799																		
with one or more adverse events	5	(0.2)																	
Hypothyroidism	5	(0.2)																	

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

	<p>Of the 24 patients with events of hypothyroidism in the UC population in KN052 (n=370) and the 17 events of hypothyroidism in the UC population in KN045 (n=266), none of the patients experienced serious events of hypothyroidism.</p> <p>Of the 92 patients with events of hypothyroidism in the HNSCC population in KN040, KN012 and KN055 (n=609), none of the patients experienced serious events of hypothyroidism.</p> <p>Of the 54 patients with events of hypothyroidism in the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), none of the patients experienced serious events of hypothyroidism. Of the 42 patients with events of hypothyroidism in the HNSCC population in KN048 pembrolizumab plus chemotherapy group (n=276), none of the patients experienced serious events of hypothyroidism.</p> <p>Of the 75 patients with events of hypothyroidism in the melanoma population in KN054 (n=509), none of the patients experienced serious events of hypothyroidism.</p> <p>Of the 24 patients with events of hypothyroidism in the NSCLC population in KN407 and KN021 (Cohort A) (n=303), 1 (0.3%) experienced a serious event of hypothyroidism.</p> <p>Of the 152 patients with events of hypothyroidism in the RCC population in KN426 pembrolizumab plus axitinib (n=429), 1 (0.2%) experienced a serious event of hypothyroidism.</p> <p>Of the 19 patients with events of hypothyroidism in the CRC population in KN177 (n=153), none of the patients experienced serious events of hypothyroidism.</p> <p>There were no patients with serious events of hypothyroidism in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370).</p> <p>Of the 95 patients with events of hypothyroidism in the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), 1 (0.2%) experienced a serious event of hypothyroidism.</p> <p>Of the 166 patients with events of hypothyroidism in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), 3 (0.9%) experienced serious events of hypothyroidism.</p> <p>Of the 296 patients with events of hypothyroidism in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), 3 (0.6%) experienced serious events of hypothyroidism.</p> <p>Of the 103 patients with events of hypothyroidism in the RCC population in KN564 pembrolizumab monotherapy (n=488), none of the patients experienced serious events of hypothyroidism.</p> <p>Of the 53 patients with events of hypothyroidism in the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), none of the patients experienced serious events of hypothyroidism.</p> <p>There were no patients with serious events of hypothyroidism in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307).</p> <p>Of the 118 patients with events of hypothyroidism in the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), 3 (0.4%) experienced serious events of hypothyroidism.</p> <p>There were no patients with serious events of hypothyroidism in the melanoma population in KN716 pembrolizumab monotherapy (n=483).</p> <p>Of the 37 patients with events of hypothyroidism in gastric cancer population in KN811 pembrolizumab combination therapy (n=350), 1 (0.3%) experienced a serious event of hypothyroidism.</p> <p>Of the 120 patients with events of hypothyroidism in the NSCLC population in KN091 pembrolizumab monotherapy (n=580), 2 (0.3%) experienced serious events of hypothyroidism.</p> <p>There were no serious events of hypothyroidism in the gastric population in KN859 pembrolizumab monotherapy (n=785).</p> <p>Of the 46 patients with events of hypothyroidism in the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), 1 (0.2%) experienced a serious event of hypothyroidism.</p>
--	---

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

	<p>Of the 44 patients with events of hypothyroidism in the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396), 1 (0.3%) experienced a serious event of hypothyroidism.</p> <p>Of the 60 patients with events of hypothyroidism in the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), 1 (0.2%) experienced a serious event of hypothyroidism.</p> <p>Of the 89 patients with events of hypothyroidism in the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), none of the patients experienced serious events of hypothyroidism.</p> <p><u>Pembrolizumab SC</u></p> <p>There were no serious events of hypothyroidism in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251).</p> <p style="text-align: center;">Hyperthyroidism - Serious (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment-Emergent)</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th colspan="2" style="text-align: center;">Cumulative NSCLC 001/010 + MEL 001/002/006</th> </tr> <tr> <th></th> <th style="text-align: center;">n</th> <th style="text-align: center;">(%)</th> </tr> </thead> <tbody> <tr> <td>Subjects in population</td> <td style="text-align: center;">2,799</td> <td></td> </tr> <tr> <td>with one or more adverse events</td> <td style="text-align: center;">6</td> <td style="text-align: center;">(0.2)</td> </tr> <tr> <td>Hyperthyroidism</td> <td style="text-align: center;">6</td> <td style="text-align: center;">(0.2)</td> </tr> </tbody> </table> <p>Every subject is counted a single time for each applicable row and column. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns meets the incidence criterion in the report title, after rounding. Include all treated subjects in KN001 Part B1, B2, B3, D, C, F1, F2, F3 and all subjects in KN002 treated with Pembrolizumab in the original phase and all subjects in KN006 and KN010 treated with Pembrolizumab.</p> <p><u>Pembrolizumab IV</u></p> <p>Of the 50 patients with events of hyperthyroidism in the NSCLC population in KN024 and KN042 (n=790), 2 (0.3%) experienced serious events of hyperthyroidism.</p> <p>Of the 21 patients with events of hyperthyroidism in the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), none of the patients experienced serious events of hyperthyroidism.</p> <p>Of the 17 patients with events of hyperthyroidism in the HL population in KN013, KN087, and KN204 (n=389), none of the patients experienced serious events of hyperthyroidism.</p> <p>Of the 9 patients with events of hyperthyroidism in the UC population in KN052 (n=370) and the 10 events of hyperthyroidism in the UC population in KN045 (n=266), none of the of the patients experienced serious events of hyperthyroidism.</p> <p>Of the 10 patients with events of hyperthyroidism in the HNSCC population in KN040, KN012 and KN055 (n=609), 1 (0.2%) experienced a serious event of hyperthyroidism.</p> <p>Of the 8 patients with events of hyperthyroidism in the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), 1 (0.3%) experienced a serious event of hyperthyroidism. Of the 13 patients with events of hyperthyroidism in the HNSCC pembrolizumab plus chemotherapy group (n=276), none of the patients experienced serious events of hyperthyroidism.</p> <p>Of the 53 patients with events of hyperthyroidism in the melanoma population in KN054 (n=509), 1 (0.2%) experienced a serious event of hyperthyroidism.</p>		Cumulative NSCLC 001/010 + MEL 001/002/006			n	(%)	Subjects in population	2,799		with one or more adverse events	6	(0.2)	Hyperthyroidism	6	(0.2)
	Cumulative NSCLC 001/010 + MEL 001/002/006															
	n	(%)														
Subjects in population	2,799															
with one or more adverse events	6	(0.2)														
Hyperthyroidism	6	(0.2)														

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

	<p>Of the 20 patients with events of hyperthyroidism in the NSCLC population in KN407 and KN021 (Cohort A) (n=303), 1 (0.3%) experienced a serious event of hyperthyroidism.</p> <p>Of the 55 patients with events of hyperthyroidism in the RCC population in KN426 pembrolizumab plus axitinib (n=429), 2 (0.5%) experienced serious events of hyperthyroidism.</p> <p>Of the 6 patients with events of hyperthyroidism in the CRC population in KN177 (n=153), none of the patients experienced serious events of hyperthyroidism.</p> <p>Of the 21 patients with events of hyperthyroidism in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), 2 (0.5%) experienced serious events of hyperthyroidism.</p> <p>Of the 31 patients with events of hyperthyroidism in the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), none of the patients experienced serious events of hyperthyroidism.</p> <p>Of the 28 patients with events of hyperthyroidism in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), none of the patients experienced serious events of hyperthyroidism.</p> <p>Of the 54 patients with events of hyperthyroidism in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), 3 (0.6%) experienced serious events of hyperthyroidism.</p> <p>Of the 58 patients with events of hyperthyroidism in the RCC population in KN564 pembrolizumab monotherapy (n=488), 2 (0.4%) experienced serious events of hyperthyroidism.</p> <p>Of the 25 patients with events of hyperthyroidism in the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), 1 (0.2%) experienced a serious event of hyperthyroidism.</p> <p>There were no patients with serious events of hyperthyroidism in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307).</p> <p>Of the 41 patients with events of hyperthyroidism in the TNBC population in KN522 pembrolizumab combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), 2 (0.3%) experienced serious events of hyperthyroidism.</p> <p>There were no patients with serious events of hyperthyroidism in the melanoma population in KN716 pembrolizumab monotherapy (n=483).</p> <p>Of the 14 patients with events of hyperthyroidism in the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), none of the patients experienced serious events of hyperthyroidism.</p> <p>Of the 62 patients with events of hyperthyroidism in the NSCLC population in KN091 pembrolizumab monotherapy (n=580), 2 (0.3%) experienced serious events of hyperthyroidism.</p> <p>Of the 44 patients with events of hyperthyroidism in the gastric population in KN859 pembrolizumab monotherapy (n=785), none of the patients experienced serious events of hyperthyroidism.</p> <p>Of the 19 patients with events of hyperthyroidism in the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), 1 (0.2%) experienced a serious event of hyperthyroidism.</p> <p>Of the 22 patients with events of hyperthyroidism in the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396), 1 (0.3%) experienced a serious event of hyperthyroidism.</p> <p>There were no patients with serious events of hyperthyroidism in the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564).</p> <p>There were no serious events of hyperthyroidism in the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260).</p>
--	--

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

	<p>Of the 32 patients with events of hyperthyroidism in the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), none of the patients experienced serious events of hyperthyroidism.</p> <p><u>Pembrolizumab SC</u></p> <p>There were no serious events of hyperthyroidism in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251).</p> <p style="text-align: center;">Thyroiditis - Serious (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment-Emergent)</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th colspan="2" style="text-align: center;">Cumulative NSCLC 001/010 + MEL 001/002/006</th> </tr> <tr> <th></th> <th style="text-align: center;">n</th> <th style="text-align: center;">(%)</th> </tr> </thead> <tbody> <tr> <td>Subjects in population</td> <td style="text-align: center;">2,799</td> <td></td> </tr> <tr> <td>with one or more adverse events</td> <td style="text-align: center;">1</td> <td style="text-align: center;">(0.0)</td> </tr> <tr> <td>Thyroiditis</td> <td style="text-align: center;">1</td> <td style="text-align: center;">(0.0)</td> </tr> </tbody> </table> <p>Every subject is counted a single time for each applicable row and column. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns meets the incidence criterion in the report title, after rounding. Include all treated subjects in KN001 Part B1, B2, B3, D, C, F1, F2, F3 and all subjects in KN002 treated with Pembrolizumab in the original phase and all subjects in KN006 and KN010 treated with Pembrolizumab.</p> <p><u>Pembrolizumab IV</u></p> <p>The 1 patient with an event of thyroiditis in the NSCLC population in KN024 and KN042 (n=790), did not experience a serious event of thyroiditis.</p> <p>The 1 patient with an event of thyroiditis in the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), did not experience a serious event of thyroiditis.</p> <p>Of the 5 patients with events of thyroiditis in the HL population in KN013, KN087, and KN204 (n=389), none of the patients experienced serious events of thyroiditis.</p> <p>Of the 3 patients with events of thyroiditis in the UC population in KN052 (n=370), 1 (0.3%) was serious. There were no patients with serious events of thyroiditis reported in the UC population in KN045.</p> <p>Of the 3 patients with events of thyroiditis in the HNSCC population in KN040, KN012 and KN055 (n=609), none of the patients experienced serious events of thyroiditis.</p> <p>There were no patients with events of thyroiditis in the HNSCC population in KN048 monotherapy group (n=300). The 1 patient with an event of thyroiditis in the HNSCC population in KN048 pembrolizumab plus chemotherapy group (n=278) did not experience a serious event of thyroiditis.</p> <p>Of the 16 patients with events of thyroiditis in the melanoma population in KN054 (n=509), 2 (0.4%) experienced serious events of thyroiditis.</p> <p>Of the 3 patients with events of thyroiditis in the NSCLC population in KN407 and KN021 (Cohort A) (n=303), 1 (0.3%) experienced a serious event of thyroiditis.</p> <p>Of the 12 patients with events of thyroiditis in the RCC population in KN426 pembrolizumab plus axitinib (n=429), none of the patients experienced serious events of thyroiditis.</p> <p>Of the 2 patients with events of thyroiditis in the CRC population in KN177 (n=153), 1 (0.7%) experienced a serious event of thyroiditis.</p> <p>There were no patients with serious events of thyroiditis in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370).</p>		Cumulative NSCLC 001/010 + MEL 001/002/006			n	(%)	Subjects in population	2,799		with one or more adverse events	1	(0.0)	Thyroiditis	1	(0.0)
	Cumulative NSCLC 001/010 + MEL 001/002/006															
	n	(%)														
Subjects in population	2,799															
with one or more adverse events	1	(0.0)														
Thyroiditis	1	(0.0)														

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

	<p>Of the 8 patients with events of thyroiditis in the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), none of the patients experienced serious events of thyroiditis.</p> <p>Of the 2 patients with events of thyroiditis in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), none of the patients experienced serious events of thyroiditis.</p> <p>Of the 12 patients with events of thyroiditis in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), none of the patients experienced serious events of thyroiditis.</p> <p>Of the 6 patients with events of thyroiditis in the RCC population in KN564 pembrolizumab monotherapy (n=488), 2 (0.4%) experienced serious events of thyroiditis.</p> <p>There were no patients with serious events of thyroiditis in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307).</p> <p>Of the 16 patients with events of thyroiditis in the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), 2 (0.3%) experienced serious events of thyroiditis.</p> <p>There were no patients with serious events of thyroiditis in the melanoma population in KN716 pembrolizumab monotherapy (n=483).</p> <p>Of the 4 patients with events of thyroiditis in the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), none of the patients experienced serious events of thyroiditis.</p> <p>Of the 6 patients with events of thyroiditis in the NSCLC population in KN091 pembrolizumab monotherapy (n=580), none of the patients experienced serious events of thyroiditis.</p> <p>Of the 9 patients with events of thyroiditis in the gastric population in KN859 pembrolizumab monotherapy (n=785), none of the patients experienced serious events of thyroiditis.</p> <p>Of the 3 patients with events of thyroiditis in the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), none of the patients experienced serious events of thyroiditis.</p> <p>Of the 4 patients with events of thyroiditis in the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396), there were no serious events of thyroiditis.</p> <p>There were no patients with serious events of thyroiditis in the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564).</p> <p>Of the 2 patients with events of thyroiditis in the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), none of the patients experienced serious events of thyroiditis.</p> <p><u>Pembrolizumab SC</u></p> <p>There were no serious events of thyroiditis in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251).</p>
--	---

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

Severity and Nature of the Risk	Identified treatment emergent adverse events (AE) of thyroid disorder (all events) in participants summarized by grade– pembrolizumab (AEs for 30 days after discontinuation were included)	
	Hypothyroidism (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment-Emergent)	
		Cumulative NSCLC 001/010 + MEL 001/002/006
		n (%)
	Subjects in population	2,799
	with one or more adverse events	237 (8.5)
	Hypothyroidism	236 (8.4)
	Grade 1	60 (2.1)
	Grade 2	173 (6.2)
	Grade 3	3 (0.1)
	Myxoedema	1 (0.0)
	Grade 2	1 (0.0)
	Primary hypothyroidism	1 (0.0)
	Grade 2	1 (0.0)
	<p>Every subject is counted a single time for each applicable specific adverse event. A subject with multiple adverse events within a system organ class is counted a single time for that system organ class.</p> <p>A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the incidence specified in the report title, after rounding.</p> <p>Only the highest reported grade of a given adverse event is counted for the individual subject.</p> <p>Grades are based on NCI CTCAE version 4.0</p>	
<u>Pembrolizumab IV</u>		
<p>In the NSCLC population in KN024 and KN042 (n=790), there was 1 (0.1%) patient with a Grade 3 event and no patients with Grade 4-5 events of hypothyroidism.</p> <p>In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), there were 2 (0.4%) patients with Grade 3 events and no patients with Grade 4-5 events of hypothyroidism.</p> <p>In the HL population in KN013, KN087, and KN204 (n=389), there were no patients with Grade 3-5 events of hypothyroidism.</p> <p>In the UC population in KN052 (n=370) and KN045 (n=266), there were no patients with Grade 3-5 events of hypothyroidism.</p> <p>In the HNSCC population in KN040, KN012 and KN055 (n=609), there were 3 (0.5%) patients with Grade 3 events and no patients with Grade 4-5 events of hypothyroidism.</p> <p>In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), there were no patients with Grade 3-5 events of hypothyroidism. In the HNSCC population in KN048 pembrolizumab plus chemotherapy group (n=276), there were no patients with Grade 3-5 events of hypothyroidism.</p> <p>In the melanoma population in KN054 (n=509), there were no patients with Grade 3-5 events of hypothyroidism.</p> <p>In the NSCLC population in KN407 and KN021 (Cohort A) (n=303), there was 1 (0.3%) patient with a Grade 3 event of hypothyroidism. There were no patients with Grade 4 or Grade 5 events of hypothyroidism.</p> <p>In the RCC population in KN426 pembrolizumab plus axitinib (n=429), there was 1 (0.2%) patient with a Grade 3 event of hypothyroidism. There were no patients with Grade 4 or 5 events of hypothyroidism.</p>		

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

	<p>In the CRC population in KN177 (n=153) there were no patients with Grade 3-5 events of hypothyroidism reported.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), there were no patients with Grade 3-5 events of hypothyroidism.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), there were 2 (0.3%) patients with Grade 3 events of hypothyroidism. There were no patients with Grade 4 or 5 events of hypothyroidism.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), there were 5 (1.4%) patients with Grade 3 events of hypothyroidism. There were no patients with Grade 4 or 5 events of hypothyroidism.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), there were 5 (0.9%) patients with Grade 3, and 1 (0.2%) patient with a Grade 4 events of hypothyroidism. There were no patients with Grade 5 events of hypothyroidism.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488) there was 1 (0.2%) patient with a Grade 3 event of hypothyroidism. There were no patients with Grade 4 or 5 events of hypothyroidism.</p> <p>In the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), there were no patients with Grade 3-5 events of hypothyroidism reported.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), there were 4 (1.3%) patients with Grade 3 events of hypothyroidism. There were no patients with Grade 4 or 5 events of hypothyroidism reported.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) there were 4 (0.5%) patients with Grade 3 events of hypothyroidism. There were no patients with Grade 4 or 5 events of hypothyroidism reported.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), there were no patients with Grade 3-5 events of hypothyroidism.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), there was 1 (0.3%) patient with a Grade 3 event of hypothyroidism. There were no patients with Grade 4 or 5 events of hypothyroidism.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580) there was 1 (0.2%) patient with a Grade 3 event of hypothyroidism. There were no patients with Grade 4 or 5 events of hypothyroidism.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), there was 1 (0.2%) patient with a Grade 3 event of hypothyroidism reported. There were no patients with Grade 4 or 5 events of hypothyroidism.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), there was 1 (0.1%) patient with a Grade 3 event of hypothyroidism reported. There were no patients with Grade 4 or 5 events of hypothyroidism reported.</p> <p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396) there were no patients with Grade 3, 4, or 5 events of hypothyroidism reported.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), there were 2 (0.4%) patients with a Grade 3 event of hypothyroidism reported. There were no patients with Grade 4 or 5 events of hypothyroidism reported.</p> <p>In the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), there were no patients with Grade 3-5 events of hypothyroidism.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), there were no patients with Grade 3-5 events of hypothyroidism.</p>
--	---

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

<p><u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251) there were no patients with Grade 3, 4, or 5 events of hypothyroidism reported.</p> <p style="text-align: center;">Hyperthyroidism (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment-Emergent)</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th rowspan="2"></th> <th colspan="2" style="text-align: center;">Cumulative NSCLC 001/010 + MEL 001/002/006</th> </tr> <tr> <th style="text-align: center;">n</th> <th style="text-align: center;">(%)</th> </tr> </thead> <tbody> <tr> <td>Subjects in population</td> <td style="text-align: center;">2,799</td> <td></td> </tr> <tr> <td>with one or more adverse events</td> <td style="text-align: center;">96</td> <td style="text-align: center;">(3.4)</td> </tr> <tr> <td>Hyperthyroidism</td> <td style="text-align: center;">96</td> <td style="text-align: center;">(3.4)</td> </tr> <tr> <td> Grade 1</td> <td style="text-align: center;">70</td> <td style="text-align: center;">(2.5)</td> </tr> <tr> <td> Grade 2</td> <td style="text-align: center;">22</td> <td style="text-align: center;">(0.8)</td> </tr> <tr> <td> Grade 3</td> <td style="text-align: center;">4</td> <td style="text-align: center;">(0.1)</td> </tr> </tbody> </table> <p>Every subject is counted a single time for each applicable specific adverse event. A subject with multiple adverse events within a system organ class is counted a single time for that system organ class. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the incidence specified in the report title, after rounding. Only the highest reported grade of a given adverse event is counted for the individual subject. Grades are based on NCI CTCAE version 4.0.</p>				Cumulative NSCLC 001/010 + MEL 001/002/006		n	(%)	Subjects in population	2,799		with one or more adverse events	96	(3.4)	Hyperthyroidism	96	(3.4)	Grade 1	70	(2.5)	Grade 2	22	(0.8)	Grade 3	4	(0.1)
	Cumulative NSCLC 001/010 + MEL 001/002/006																								
	n	(%)																							
Subjects in population	2,799																								
with one or more adverse events	96	(3.4)																							
Hyperthyroidism	96	(3.4)																							
Grade 1	70	(2.5)																							
Grade 2	22	(0.8)																							
Grade 3	4	(0.1)																							
<p><u>Pembrolizumab IV</u></p> <p>In the NSCLC population in KN024 and KN042 (n=790), there was 1 (0.1%) patient with a Grade 3 and no patients with Grade 4-5 events of hyperthyroidism.</p> <p>In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), there were no patients with Grade 3-5 events of hyperthyroidism.</p> <p>In the HL population in KN013, KN087, and KN204 (n=389), there were no patients with Grade 3-5 events of hyperthyroidism reported.</p> <p>In the UC population in KN052 (n=370) and KN045 (n=266), there were no patients with Grade 3-5 events of hyperthyroidism.</p> <p>In the HNSCC population in KN040, KN012 and KN055 (n=609), there were no patients with Grade 3-5 events of hyperthyroidism.</p> <p>In the HNSCC population in KN048 pembrolizumab monotherapy group (n=300), there was 1 (0.3%) patient with a Grade 3 event of hyperthyroidism. There were no patients with Grade 4 or 5 events of hyperthyroidism.</p> <p>In the HNSCC population in KN048 pembrolizumab plus chemotherapy group (n=276), there were no patients with Grade 3-5 events of hyperthyroidism.</p> <p>In the melanoma population in KN054 (n=509), there was 1 patient with a Grade 3 (0.2%) event of hyperthyroidism. There were no patients with Grade 4 or 5 events of hyperthyroidism.</p> <p>In the NSCLC population in KN407 and KN021 (Cohort A) (n=303), there was 1 (0.3%) patient with a Grade 3 event of hyperthyroidism. There were no patients with Grade 4 or 5 events of hyperthyroidism.</p> <p>In the RCC population in KN426 pembrolizumab plus axitinib (n=429), there were 5 (1.2%) patients with Grade 3 events of hyperthyroidism. There were no patients with Grade 4 or 5 events of hyperthyroidism.</p>																									

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

	<p>In the CRC population in KN177 (n=153) there were no patients with Grade 3-5 events of hyperthyroidism. reported.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), there was 1 (0.3%) patient with a Grade 3 event of hyperthyroidism. There were no patients with Grade 4 or 5 events of hyperthyroidism.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), there was 1 (0.2%) patient with a Grade 3 event of hyperthyroidism. There were no patients with Grade 4 or 5 events of hyperthyroidism.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), there were no patients with Grade 3-5 events of hyperthyroidism.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), there were 3 (0.6%) patients with Grade 3 events of hyperthyroidism. There were no patients with Grade 4 or 5 events of hyperthyroidism.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488) there was 1 (0.2%) patient with a Grade 3 event of hyperthyroidism. There were no patients with Grade 4 or 5 events of hyperthyroidism.</p> <p>In the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), there was 1 (0.2%) patient with a Grade 3 event of hyperthyroidism. There were no patients with Grade 4 or 5 events of hyperthyroidism reported.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), there were no patients with Grade 3-5 events of hyperthyroidism reported.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) there were 2 (0.3%) patients with Grade 3 events of hyperthyroidism. There were no patients with Grade 4 or 5 events of hyperthyroidism reported.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), there was 1 (0.2%) patient with a Grade 3 event of hyperthyroidism. There were no patients with Grade 4 or 5 events of hyperthyroidism.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), there were no patients with Grade 3-5 events of hyperthyroidism.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580) there was 1 (0.2%) patient with a Grade 3 event of hyperthyroidism. There were no patients with Grade 4 or 5 events of hyperthyroidism.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), there were no patients with Grade 3-5 events of hyperthyroidism reported.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), there was 1 (0.2%) patient with a Grade 3 event of hyperthyroidism reported. There were no patients with Grade 4 or 5 events of hyperthyroidism.</p> <p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396) there were no patients with Grade 3, 4, or 5 events of hyperthyroidism reported.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), there was 1 (0.2%) patient with a Grade 3 event of hyperthyroidism. There were no patients with Grade 4 or 5 events of hyperthyroidism.</p> <p>In the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260), there were no patients with Grade 3-5 events of hyperthyroidism reported.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), there were no patients with Grade 3-5 events of hyperthyroidism.</p>
--	---

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

<p><u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251) there were no patients with Grade 3, 4, or 5 events of hyperthyroidism reported.</p> <p style="text-align: center;">Thyroiditis (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and KN010 NSCLC Subjects Treated with Pembrolizumab (Treatment-Emergent)</p> <table border="1"> <thead> <tr> <th rowspan="2"></th> <th colspan="2">Cumulative NSCLC 001/010 + MEL 001/002/006</th> </tr> <tr> <th>n</th> <th>(%)</th> </tr> </thead> <tbody> <tr> <td>Subjects in population</td> <td>2,799</td> <td></td> </tr> <tr> <td>with one or more adverse events</td> <td>16</td> <td>(0.6)</td> </tr> <tr> <td>Autoimmune thyroiditis</td> <td>5</td> <td>(0.2)</td> </tr> <tr> <td> Grade 1</td> <td>2</td> <td>(0.1)</td> </tr> <tr> <td> Grade 2</td> <td>3</td> <td>(0.1)</td> </tr> <tr> <td>Thyroiditis</td> <td>11</td> <td>(0.4)</td> </tr> <tr> <td> Grade 1</td> <td>7</td> <td>(0.3)</td> </tr> <tr> <td> Grade 2</td> <td>4</td> <td>(0.1)</td> </tr> </tbody> </table> <p>Every subject is counted a single time for each applicable specific adverse event. A subject with multiple adverse events within a system organ class is counted a single time for that system organ class. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the incidence specified in the report title, after rounding. Only the highest reported grade of a given adverse event is counted for the individual subject. Grades are based on NCI CTCAE version 4.0.</p>				Cumulative NSCLC 001/010 + MEL 001/002/006		n	(%)	Subjects in population	2,799		with one or more adverse events	16	(0.6)	Autoimmune thyroiditis	5	(0.2)	Grade 1	2	(0.1)	Grade 2	3	(0.1)	Thyroiditis	11	(0.4)	Grade 1	7	(0.3)	Grade 2	4	(0.1)
	Cumulative NSCLC 001/010 + MEL 001/002/006																														
	n	(%)																													
Subjects in population	2,799																														
with one or more adverse events	16	(0.6)																													
Autoimmune thyroiditis	5	(0.2)																													
Grade 1	2	(0.1)																													
Grade 2	3	(0.1)																													
Thyroiditis	11	(0.4)																													
Grade 1	7	(0.3)																													
Grade 2	4	(0.1)																													
<p><u>Pembrolizumab IV</u></p> <p>In the NSCLC population in KN024 and KN042 (n=790), there were no patients with Grade 3-5 events of thyroiditis.</p> <p>In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), there were no patients with Grade 3-5 events of thyroiditis.</p> <p>In the HL population in KN013, KN087, and KN204 (n=389), there were no patients with Grade 3-5 events of thyroiditis reported.</p> <p>In the UC population in KN052 (n=370), there was 1 (0.3%) patient with a Grade 3 event of thyroiditis and no patients with Grade 4 or Grade 5 events and in the UC population in KN045 (n=266), there were no patients with Grade 3-5 events of thyroiditis.</p> <p>In the HNSCC population in KN040, KN012 and KN055 (n=609), there were no patients with Grade 3-5 events of thyroiditis.</p> <p>In the HNSCC population in KN048 within both the pembrolizumab monotherapy (n=300) and pembrolizumab plus chemotherapy (n=276) groups, there were no patients with Grade 3-5 events of thyroiditis.</p> <p>In the melanoma population in KN054 (n=509), there were no patients with Grade 3-5 events of thyroiditis.</p> <p>In the NSCLC population in KN407 and KN021 (Cohort A) (n=303), there was 1 (0.3%) patient with a Grade 3 event of thyroiditis reported. There were no patients with Grade 4 or Grade 5 events of thyroiditis.</p> <p>In the RCC population in KN426 pembrolizumab plus axitinib (n=429), there was 1 (0.2%) patient with a Grade 3 event of thyroiditis. There were no patients with Grade 4 or 5 events of thyroiditis.</p>																															

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

	<p>In the CRC population in KN177 (n=153) there were no patients with Grade 3-5 events of thyroiditis reported.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), there were no patients with Grade 3-5 events of thyroiditis.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), there was 1 (0.2%) patient with a Grade 3 event of thyroiditis. There were no patients with Grade 4 or 5 events of thyroiditis.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), there were no patients with Grade 3-5 events of thyroiditis.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), there were no patients with Grade 3-5 events of thyroiditis reported.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488) there were 2 (0.4%) Grade 3 events of thyroiditis. There were no patients with Grade 4 or 5 events of thyroiditis.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), there were 2 (0.7%) patients with Grade 3 and no patients with Grade 4 or 5 events of thyroiditis reported.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) there were 2 (0.3%) patients with Grade 3 events of thyroiditis. There were no patients with Grade 4 or 5 events of thyroiditis reported.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), there were no patients with Grade 3-5 events of thyroiditis.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), there were no patients with Grade 3-5 events of thyroiditis.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580) there were no patients with Grade 3, 4 or 5 events of thyroiditis.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), there were no patients with Grade 3-5 events of thyroiditis reported.</p> <p>In the BTC population in KN966 pembrolizumab plus chemotherapy (n=529), there were no patients with Grade 3, 4, or 5 events of thyroiditis.</p> <p>In the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396) there were no patients with Grade 3, 4, or 5 events of thyroiditis reported.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), there were no patients with Grade 3-5 events of thyroiditis reported.</p> <p>In the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361), there were no patients with Grade 3-5 events of thyroiditis.</p> <p><u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251) there were no patients with Grade 3, 4, or 5 events of thyroiditis reported.</p> <p>In the global safety database through 31-DEC-2017, (pembrolizumab monotherapy trials only), there were no Grade 5 clinical trial reports of thyroid disorders.</p> <p>Review of pembrolizumab clinical trial data from ongoing studies and postmarketing data regarding immune-mediated endocrinopathies- thyroid disorder (hypothyroidism, hyperthyroidism, thyroiditis), including reports of fatal cases of hypothyroidism and thyroiditis, is consistent with the risk as characterized in this RMP.</p>
<p><u>Risk Factors and Risk Groups:</u></p>	<p>No specific risk factors for thyroid disorders (hyperthyroidism, hypothyroidism, thyroiditis) associated with pembrolizumab have been identified.</p>

Table SVII.3.1.7: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies - Thyroid Disorder (Hypothyroidism, Hyperthyroidism, Thyroiditis)

<u>Preventability:</u>	Although the development of thyroid disorders (hypothyroidism, hyperthyroidism, thyroiditis) cannot be completely prevented; patients should be monitored for signs and symptoms of thyroid disorders (hypothyroidism, hyperthyroidism, thyroiditis) and if they develop, other causes excluded. Withholding/discontinuation of pembrolizumab and appropriate medical intervention including corticosteroids following the onset of thyroid disorders (hypothyroidism, hyperthyroidism, thyroiditis) may result in recovery.
<u>Impact on the Risk-Benefit Balance of the Product:</u>	Severe thyroid disorders (hypothyroidism, hyperthyroidism, and thyroiditis) may be reported to be life-threatening or fatal in individual patients. However, given the fatal outcome of untreated cancer, the risk of thyroid disorders which generally can be managed, is outweighed by the potential benefit of pembrolizumab treatment.
<u>Public Health Impact:</u>	This risk has minimal public health impact outside its effect on individual patients.

Table SVII.3.1.8: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies -Type 1 Diabetes Mellitus

<u>Important Identified Risk:</u>	Immune-mediated Endocrinopathies-Type 1 Diabetes Mellitus (T1DM)									
<u>Potential Mechanisms:</u>	Pembrolizumab is a potent and highly-selective humanized monoclonal antibody (mAb) of the Immunoglobulin G4 (IgG4)/kappa isotype directed to the programmed cell death-1 (PD-1) receptor and designed to directly block the interaction between the receptor and its ligands, PD-L1 and PD-L2. The PD-1 pathway represents a major immune control switch which may be engaged by tumor cells to overcome active T-cell immune surveillance. T-cell activation in proximity to normal tissue may lead to inflammation and injury to normal tissue.									
<u>Evidence Source(s) and Strength of Evidence:</u>	Review of pembrolizumab clinical trial data, post-marketing experience and literature regarding endocrinopathies- type 1 diabetes mellitus represent sufficient evidence of a causal association with pembrolizumab exposure. CTD 2.7.4 Please reference evidence source cited in Table SVII.3.1.1; Details of Important Identified Risk: Immune-Mediated Pneumonitis									
<u>Characterisation of the Risk:</u> Immune-mediated Endocrinopathies-Type 1 Diabetes Mellitus (T1DM)										
<u>Frequency with 95%CI</u>	<p>95% Confidence Interval for the Overall Incidence (%) of Type 1 Diabetes Mellitus KN001, KN002 and KN006 Melanoma Subjects and KN001 and 010 NSCLC Subjects Treated with Pembrolizumab</p> <table border="1"> <thead> <tr> <th></th> <th colspan="2">Cumulative MEL 001/002/006 + NSCLC 001/010</th> </tr> <tr> <th></th> <th>n (%)</th> <th>95% CI</th> </tr> </thead> <tbody> <tr> <td>Type 1 Diabetes Mellitus</td> <td>6 (0.2)</td> <td>(0.1,0.5)</td> </tr> </tbody> </table> <p><u>Pembrolizumab IV:</u> The overall number and proportion of patients with T1DM in the NSCLC population in KN024 and KN042 (n=790) was 1 (0.1%) (95%CI-0, 0.7), which is generally similar to the proportion of patients with the event of T1DM in the reference safety dataset for melanoma and NSCLC. The overall number and proportion of patients with T1DM in the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488) was 1 (0.2%) (95% CI-0, 1.1), which is generally similar to the proportion of patients with the event of T1DM in the reference safety dataset for melanoma and NSCLC. There were no patients with events of T1DM reported in the HL population in KN013, KN087, and KN204 (n=389).</p>		Cumulative MEL 001/002/006 + NSCLC 001/010			n (%)	95% CI	Type 1 Diabetes Mellitus	6 (0.2)	(0.1,0.5)
	Cumulative MEL 001/002/006 + NSCLC 001/010									
	n (%)	95% CI								
Type 1 Diabetes Mellitus	6 (0.2)	(0.1,0.5)								

Table SVII.3.1.8: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies -Type 1 Diabetes Mellitus

	<p>The overall number and proportion of patients with T1DM in the UC population in KN052 (n=370) was 4 (1.1%) (95%CI-0.3,2.7), which is generally similar to the proportion of patients with the event of T1DM in the reference safety dataset for melanoma and NSCLC. There were no patients with events of T1DM in the UC population in KN045.</p> <p>The overall number and proportion of patients with T1DM in the HNSCC population in KN040, KN012 and KN055 (n=609) was 3 (0.5%) (95%CI-0.1,1.4), which is generally similar to the proportion of patients with the event of T1DM in the reference safety dataset for melanoma and NSCLC.</p> <p>There were no patients with events of T1DM reported in the HNSCC population in KN048 pembrolizumab monotherapy group (n=300) or pembrolizumab plus chemotherapy group (n=276).</p> <p>The overall number and proportion of patients with T1DM in the melanoma population in KN054 (n=509) was 5 (1.0%) (95%CI-0.3,2.3). The proportion of patients with events of T1DM in KN054 although somewhat more common than in the reference safety dataset were generally consistent in nature with the proportion of events of T1DM previously reported for pembrolizumab.</p> <p>There were no patients with events of T1DM reported in the NSCLC population in KN407 and KN021 (Cohort A) (n=303).</p> <p>The overall number and proportion of patients with T1DM in the RCC population in KN426 pembrolizumab plus axitinib (n=429) was 1 (0.2%) (95% CI-0,1.3), which is generally similar to the overall frequency in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with T1DM in the CRC population in KN177 (n=153) was 1 (0.7%) (95% CI-0, 3.6) which is generally similar to the overall frequency in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with T1DM in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370) was 1 (0.3%) (95% CI-0,1.5), which is generally similar to the proportion of patients with the event of T1DM in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with T1DM in the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596) was 2 (0.3%) (95% CI-0,1.2), which is generally similar to the proportion of patients with the event of T1DM in the reference dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with T1DM in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352) was 2 (0.6%), which is generally similar to the proportion of patients with the event of T1DM in the reference safety dataset.</p> <p>The overall number and proportion of patients with T1DM in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530) was 4 (0.8%) (95% CI-0.2, 1.9), which is generally similar to the proportion of patients with the event of T1DM in the reference safety dataset.</p> <p>The overall number and proportion of patients with T1DM in the RCC population in KN564 pembrolizumab monotherapy (n=488) was 9 (1.8%) (95% CI-0.8,3.5). The proportion of patients with events of T1DM in KN564 although somewhat more common than in the reference safety dataset were generally consistent in nature with the proportion of patients with events of T1DM previously reported for pembrolizumab.</p> <p>The overall number and proportion of patients with T1DM in the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475) was 2 (0.4%) (95% CI-0.1,1.5) which is generally similar to the proportion of patients with the event of T1DM in the reference safety dataset.</p> <p>The overall number and proportion of patients with T1DM in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307) was 2 (0.7%) (95% CI-0.1, 2.3), which is generally similar to the proportion of patients with the event of T1DM in the reference safety dataset.</p> <p>The overall number and proportion of patients with T1DM in the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) was 4 (0.5%) (95% CI-0.1,1.3), which is generally similar to the proportion of patients with the event of T1DM in the reference safety dataset.</p>
--	---

Table SVII.3.1.8: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies -Type 1 Diabetes Mellitus

	<p>The overall number and proportion of patients with T1DM in the melanoma population in KN716 pembrolizumab monotherapy (n=483) was 2 (0.4%) (95% CI-0.1, 1.5), which is generally similar to the proportion of patients with the event of T1DM in the reference safety dataset for melanoma and NSCLC.</p> <p>The overall number and proportion of patients with T1DM in the gastric cancer population in KN811 pembrolizumab combination therapy (n=350) was 1 (0.3%) (95% CI-0.0, 1.6), which is generally similar to the proportion of patients with events of T1DM in the reference safety dataset.</p> <p>The overall number and proportion of patients with T1DM in the NSCLC population in KN091 pembrolizumab monotherapy (n=580) was 1 (0.2%) (95% CI-0.0,1.0), which is generally similar to the proportion of patients with the event of T1DM in the reference safety dataset.</p> <p>The overall number and proportion of patients with T1DM in the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785) was 5 (0.6%) (95% CI-0.2, 1.5), which is generally similar to the proportion of patients with the events of T1DM in the reference safety dataset.</p> <p>There were no patients with events of T1DM reported in the BTC population in KN966 pembrolizumab plus chemotherapy group (n=529).</p> <p>There were no events of T1DM reported in the NSCLC population in KN671 pembrolizumab in combination with chemotherapy as neoadjuvant treatment followed by continued adjuvant pembrolizumab monotherapy (n=396).</p> <p>The overall number and proportion of patients with T1DM in the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564) was 1 (0.2%) (95% CI-0.0%, 1.0%), which is generally similar to the proportion of patients with the events of T1DM in the reference safety dataset.</p> <p>The overall number and proportion of patients with T1DM in the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382) was 2 (0.5%) (95% CI-0.1, 1.9), which is generally similar to the proportion of patients with the event of T1DM in the reference safety dataset.</p> <p>The overall number and proportion of patients with T1DM in the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528) was 1 (0.2%) (95% CI-0.0, 1.1), which is generally similar to the proportion of patients with events of T1DM in the reference safety dataset.</p> <p>There were no patients with events of T1DM reported in the mesothelioma population in KN483 and KNA17 pembrolizumab plus chemotherapy (n=260).</p> <p>There were no patients with events of T1DM reported in the HNSCC population in KN689 who received pembrolizumab monotherapy as neoadjuvant treatment continued as adjuvant treatment in combination with radiation with or without chemotherapy and then as monotherapy (n=361).</p> <p><u>Pembrolizumab SC</u></p> <p>The overall number and proportion of patients with events of T1DM reported in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251) was 1 (0.4%) (95% CI-0.0, 2.2), which is generally similar to the proportion of patients with events of T1DM in the reference safety dataset.</p>
--	--

Table SVII.3.1.8: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies -Type 1 Diabetes Mellitus

Outcomes	Percent overall, fatal, not resolved, resolved, and unknown for identified risk of Type 1 Diabetes Mellitus - pembrolizumab (Table includes events for 30 days after discontinuation.)		
	Type 1 Diabetes Mellitus (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and 010 NSCLC Subjects Treated with Pembrolizumab (Treatment Emergent)		
			Cumulative NSCLC 001/010 + MEL 001/002/006
		Outcome	n (%)
	Subjects in population		2,799
	With one or more adverse events	Overall	6 (0.2)
		Fatal	0 (0.0)
		Not Resolved	4 (0.1)
		Resolved	2 (0.1)
		Unknown	0 (0.0)
	Every subject is counted once on each applicable row. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the percent incidence specified in the report title, after rounding. Outcome: Resolved = RECOVERED/RESOLVED, Not resolved = NOT RECOVERED/NOT RESOLVED.		
	<u>Pembrolizumab IV</u>		
	In the NSCLC population in KN024 and KN042 (n=790), the outcome of the 1 (0.1%) patient with an event of T1DM was resolved. There were no patients with events of T1DM with a fatal outcome.		
	In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), the outcome of the 1 (0.2%) patient with an event of T1DM was not resolved. There were no patients with events of T1DM with a fatal outcome.		
	In the UC population in KN052 (n=370), the outcomes for the 4 patients with events of T1DM were as follows: 1 (0.3%) resolved, 1 (0.3%) resolved with sequelae, and 2 (0.5%) not resolved. There were no patients with events of T1DM with a fatal outcome.		
In the HNSCC population in KN040, KN012 and KN055 (n=609), the outcomes of the 3 patients with events of T1DM were as follows: 3 (0.5%) not resolved. There were no fatal outcomes.			
In the melanoma population in KN054 (n=509), the outcome of the 5 patients with events of T1DM were as follows: 5 (1.0%) not resolved. There were no patients with events of T1DM with a fatal outcome.			
In the RCC population in KN426 pembrolizumab plus axitinib (n=429), the outcome for the 1 (0.2%) patient with an event of T1DM was resolved with sequelae. There were no fatal outcomes.			
In the CRC population in KN177 (n=153), the outcome for the 1 patient with an event of T1DM was as follows: 1 (0.7%) not resolved. There were no patients with events of T1DM with a fatal outcome.			
In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), the outcome for the 1 (0.3%) patient with an event of T1DM was not resolved. There were no patients with events of T1DM with a fatal outcome.			
In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), the outcome for the 2 patients with events of T1DM were as follows: 1 (0.2%) not resolved, and 1 (0.2%) resolved with sequelae. There were no patients with events of T1DM with a fatal outcome.			
In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), the outcome for the 2 patients with events of T1DM were as follows: 2 (0.6%) not resolved. There were no patients with events of T1DM with a fatal outcome.			
In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), the outcome for the 4 (0.8%) patients with events of T1DM were as follows: 1 (0.2%) resolved, 2 (0.4%) resolved with sequelae, and 1 (0.2%) resolving. There were no patients with events of T1DM with a fatal outcome.			

Table SVII.3.1.8: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies -Type 1 Diabetes Mellitus

	<p>In the RCC population in KN564 pembrolizumab monotherapy (n=488), the outcomes for the 9 patients with events of T1DM were as follows: 8 (1.6%) resolved with sequelae and 1 (0.2%) resolving. There were no patients with events of T1DM with a fatal outcome.</p> <p>In the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), the outcomes for the 2 (0.4%) patients with events of T1DM were as follows: 1 (0.2%) resolved and 1 (0.2%) not resolved. There were no patients with events of T1DM with a fatal outcome.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), the outcomes for the 2 patients with events of T1DM were as follows: 2 (0.7%) resolved. There were no patients with events of T1DM with a fatal outcome.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), the outcomes for the 4 patients with events of T1DM were as follows: 3 (0.4%) not resolved, and 1 (0.1%) resolved with sequelae. There were no patients with events of T1DM with a fatal outcome.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), the outcomes of the 2 patients with events of T1DM were as follows: 1 (0.2%) not resolved, and 1 (0.2%) resolved with sequelae. There were no patients with events of T1DM with a fatal outcome.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), the outcome for the 1 (0.3%) patient with the event of T1DM was resolved. There were no patients with events of T1DM with a fatal outcome.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580), the outcomes for the 1 patient with an event of T1DM was as follows: 1 (0.2%) not resolved. There were no patients with events of T1DM with a fatal outcome.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), the outcomes for the 5 patients with events of T1DM were as follows: 2 (0.3%) not resolved, 2 (0.3%) resolved, and 1 (0.1%) resolved with sequelae. There were no patients with events of T1DM with a fatal outcome.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), the outcome for the 1 patient with the event of T1DM was as follows: 1 (0.2%) not resolved. There were no patients with events of T1DM with a fatal outcome.</p> <p>In the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382) the outcomes for patients with events of T1DM were not recorded per study protocol (managed by external collaboration).</p> <p>In the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528), the outcome for the 1 (0.2%) patient with the event of T1DM was not resolved. There were no patients with an event of T1DM with a fatal outcome.</p> <p><u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251), the outcome for the 1 (0.4%) patient with the event of T1DM was resolved. There were no patients with an event of T1DM with a fatal outcome.</p> <p><u>Aggregate Review</u></p> <p>Review of the ASE dataset (n=7730) yielded no fatal cases of T1DM. Review of the global safety database yielded the following: 1 fatal case from ongoing interventional monotherapy clinical trials and no fatal cases from the PM or non-interventional environment.</p>
--	--

Table SVII.3.1.8: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies -Type 1 Diabetes Mellitus

Seriousness	Percent serious for identified risk of Type 1 Diabetes Mellitus - Pembrolizumab (Serious AEs for 90 days after discontinuation were included.)	
	Type 1 Diabetes Mellitus - Serious (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and 010 NSCLC Subjects Treated with Pembrolizumab (Treatment Emergent)	
		Cumulative NSCLC 001/010 + MEL 001/002/006
		n (%)
	Subjects in population	2,799
	with one or more adverse events	6 (0.2)
	Diabetic ketoacidosis	2 (0.1)
	Type 1 diabetes mellitus	4 (0.1)
	Every subject is counted a single time for each applicable row and column. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns meets the incidence criterion in the report title, after rounding. Include all treated subjects in KN001 Part B1, B2, B3, D, C, F1, F2, F3 and all subjects in KN002 treated with Pembrolizumab in the original phase and all subjects in KN006 and KN010 treated with Pembrolizumab.	
	<u>Pembrolizumab IV</u>	

The 1 patient with an event of T1DM reported in the NSCLC population in KN024 and KN042 (n=790) experienced a serious event of T1DM (0.1%).

The 1 patient with an event of T1DM reported in the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488) experienced a serious event of T1DM (0.2%).

Of the 4 patients with events of T1DM reported in the UC population in KN052 (n=370), all experienced serious events of T1DM (1.1%).

Of the 3 patients with events of T1DM reported in the HNSCC population in KN040, KN012 and KN055 (n=609), all experienced serious events of T1DM (0.5%).

Of the 5 patients with events of T1DM reported in the melanoma population in KN054 (n=509), all experienced serious events of T1DM (1.0%).

The 1 patient with an event of T1DM reported in the RCC population in KN426 pembrolizumab plus axitinib (n=429) experienced a serious event of T1DM (0.2%).

There were no patients with serious events of T1DM reported in the CRC population in KN177 (n=153).

There were no patients with serious events of T1DM in the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370).

Of the 2 patients with events of T1DM reported in the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), 1 (0.2%) experienced a serious event of T1DM.

Of the 2 patients with events of T1DM reported in the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), 1 (0.3%) experienced a serious event of T1DM.

All 4 patients with events of T1DM reported in the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530) experienced serious events of T1DM (0.8%).

Of the 9 patients with events of T1DM in the RCC population in KN564 pembrolizumab monotherapy (n=488), all experienced serious events of T1DM (1.8%).

Of the 2 patients with events of T1DM in the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), 1 (0.2%) experienced a serious event of T1DM.

The 2 patients with events of T1DM reported in the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307) experienced serious events of T1DM (0.7%).

Of the 4 patients with events of T1DM in the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783), 3 (0.4%) experienced serious events of T1DM.

Table SVII.3.1.8: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies -Type 1 Diabetes Mellitus

	<p>Of the 2 patients with events of T1DM reported in the melanoma population in KN716 pembrolizumab monotherapy (n=483), both experienced serious events of T1DM (0.4%).</p> <p>The 1 patient with the event of T1DM reported in the gastric cancer population in KN811 pembrolizumab combination therapy (n=350) experienced a serious (0.3%) event.</p> <p>The 1 patient with an event of T1DM reported in the NSCLC population in KN091 pembrolizumab monotherapy (n=580) experienced a serious event of T1DM (0.2%).</p> <p>Of the 5 patients with events of T1DM in the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), all 5 (0.6%) experienced a serious event of T1DM.</p> <p>There were no patients with serious events of T1DM in the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564).</p> <p>The 2 patients with events of T1DM reported in the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382) experienced serious events of T1DM (0.5%).</p> <p>The 1 patient with an event of T1DM reported in the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528) experienced a serious (0.2%) event.</p> <p><u>Pembrolizumab SC</u></p> <p>The 1 patient with an event of T1DM reported in the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251) experienced a serious (0.4%) event.</p>																																	
<p>Severity and Nature of the Risk</p>	<p>Identified treatment emergent adverse events (AE) of Type 1 Diabetes Mellitus (all events) in participants summarized by grade–pembrolizumab (AEs for 30 days after discontinuation were included)</p> <p style="text-align: center;">Type 1 Diabetes Mellitus (AEOSI) KN001, KN002 and KN006 Melanoma Subjects and KN001 and 010 NSCLC Subjects Treated with Pembrolizumab (Treatment Emergent)</p> <table border="1" data-bbox="492 1098 1414 1409"> <thead> <tr> <th></th> <th colspan="2">Cumulative NSCLC 001/010 + MEL 001/002/006</th> </tr> <tr> <th></th> <th>n</th> <th>(%)</th> </tr> </thead> <tbody> <tr> <td>Subjects in population</td> <td>2,799</td> <td></td> </tr> <tr> <td>with one or more adverse events</td> <td>6</td> <td>(0.2)</td> </tr> <tr> <td>Diabetic ketoacidosis</td> <td>2</td> <td>(0.1)</td> </tr> <tr> <td> Grade 3</td> <td>1</td> <td>(0.0)</td> </tr> <tr> <td> Grade 4</td> <td>1</td> <td>(0.0)</td> </tr> <tr> <td>Type 1 diabetes mellitus</td> <td>5</td> <td>(0.2)</td> </tr> <tr> <td> Grade 2</td> <td>2</td> <td>(0.1)</td> </tr> <tr> <td> Grade 3</td> <td>2</td> <td>(0.1)</td> </tr> <tr> <td> Grade 4</td> <td>1</td> <td>(0.0)</td> </tr> </tbody> </table> <p>Every subject is counted a single time for each applicable specific adverse event. A subject with multiple adverse events within a system organ class is counted a single time for that system organ class. A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns is greater than or equal to the incidence specified in the report title, after rounding. Only the highest reported grade of a given adverse event is counted for the individual subject. Grades are based on NCI CTCAE version 4.0.</p> <p><u>Pembrolizumab IV</u></p> <p>In the NSCLC population in KN024 and KN042 (n=790), there was 1 (0.1%) patient with a Grade 4 event of T1DM. There were no patients with Grade 3 or 5 events of T1DM.</p> <p>In the NSCLC population in KN189 and KN021 (Cohorts C and G) pembrolizumab plus chemotherapy (n=488), there was 1 (0.2%) patient with a Grade 4 event of T1DM. There were no patients with Grade 3 or 5 events of T1DM.</p> <p>In the UC population in KN052 (n=370), there were 2 (0.5%) patients with Grade 3 events and 2 (0.5%) patients with Grade 4 events of T1DM.</p> <p>In the HNSCC population in KN040, KN012 and KN055 (n=609), there were 3 (0.5%) patients with Grade 4 events of T1DM.</p>		Cumulative NSCLC 001/010 + MEL 001/002/006			n	(%)	Subjects in population	2,799		with one or more adverse events	6	(0.2)	Diabetic ketoacidosis	2	(0.1)	Grade 3	1	(0.0)	Grade 4	1	(0.0)	Type 1 diabetes mellitus	5	(0.2)	Grade 2	2	(0.1)	Grade 3	2	(0.1)	Grade 4	1	(0.0)
	Cumulative NSCLC 001/010 + MEL 001/002/006																																	
	n	(%)																																
Subjects in population	2,799																																	
with one or more adverse events	6	(0.2)																																
Diabetic ketoacidosis	2	(0.1)																																
Grade 3	1	(0.0)																																
Grade 4	1	(0.0)																																
Type 1 diabetes mellitus	5	(0.2)																																
Grade 2	2	(0.1)																																
Grade 3	2	(0.1)																																
Grade 4	1	(0.0)																																

Table SVII.3.1.8: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies -Type 1 Diabetes Mellitus

	<p>In the melanoma population in KN054 (n=509), there were 4 (0.8%) patients with Grade 3 events and 1 (0.2%) patient with a Grade 4 event of T1DM.</p> <p>In the RCC population in KN426 pembrolizumab plus axitinib (n=429), there was 1 (0.2%) patient with a Grade 4 event of T1DM. There were no patients with Grade 5 events of T1DM.</p> <p>In the CRC population in KN177 (n=153) there was 1 (0.7%) patient with a Grade 3 event of T1DM. There were no patients with Grade 4 or 5 events of T1DM reported.</p> <p>In the esophageal cancer population in KN590 pembrolizumab plus chemotherapy (n=370), there was 1 (0.3%) patient with a Grade 3 event of T1DM. There were no patients with Grade 4 or 5 events of T1DM.</p> <p>In the TNBC population in KN355 pembrolizumab plus chemotherapy group (n=596), there was 1 (0.2%) patient with a Grade 3 event of T1DM. There were no patients with Grade 4 or 5 events of T1DM.</p> <p>In the RCC population in KN581 pembrolizumab plus lenvatinib (n=352), there was 1 (0.3%) patient with a Grade 4 event of T1DM. There were no patients with Grade 3 or 5 events of T1DM.</p> <p>In the endometrial carcinoma population in KN146 and KN775 pembrolizumab plus lenvatinib (n=530), there were 2 (0.4%) patients with Grade 3 and 2 (0.4%) patients with Grade 4 events of T1DM. There were no patients with Grade 5 events of T1DM.</p> <p>In the RCC population in KN564 pembrolizumab monotherapy (n=488) there were 7 (1.4%) patients with Grade 3 and 2 (0.4%) patients with Grade 4 events of T1DM. There were no patients with Grade 5 events of T1DM.</p> <p>In the MSI-H population in KN158 Cohort K and KN164 Cohorts A and B (n=475), there was 1 (0.2%) patient with a Grade 3 event of T1DM. There were no patients with Grade 4 or 5 events of T1DM reported.</p> <p>In the cervical cancer population in KN826 pembrolizumab plus chemotherapy (n=307), there were 2 (0.7%) patients with Grade 4 events of T1DM. There were no patients with Grade 3 or 5 events of T1DM.</p> <p>In the TNBC population in KN522 pembrolizumab in combination with neoadjuvant chemotherapy followed by continued adjuvant pembrolizumab monotherapy group (n=783) there were 2 (0.3%) patients with Grade 3 events and 2 (0.3%) patients with Grade 4 events of T1DM. There were no patients with Grade 5 events of T1DM reported.</p> <p>In the melanoma population in KN716 pembrolizumab monotherapy (n=483), there was 1 (0.2%) patient with a Grade 3 and 1 (0.2%) patient with a Grade 4 event of T1DM. There were no patients with Grade 5 events of T1DM.</p> <p>In the gastric cancer population in KN811 pembrolizumab combination therapy (n=350), there was 1 (0.3%) patient with a Grade 3 event of T1DM. There were no patients with Grade 4 or Grade 5 events of T1DM.</p> <p>In the NSCLC population in KN091 pembrolizumab monotherapy (n=580) there was 1 (0.2%) patient with a Grade 3 event of T1DM. There were no patients with Grade 4 or 5 events of T1DM.</p> <p>In the gastric cancer population in KN859 pembrolizumab plus chemotherapy (n=785), there were 4 (0.5%) patients with a Grade 3 and 1 (0.1%) patient with a Grade 4 event of T1DM reported. There were no patients with a Grade 5 events of T1DM.</p> <p>In the UC population in KN869 and KNA39 pembrolizumab + EV combination therapy (n=564), there was 1 (0.2%) patient with a Grade 3 event of T1DM. There were no patients with Grade 4 or 5 events of T1DM.</p> <p>In the endometrial carcinoma population in KN868 pembrolizumab plus chemotherapy (n=382), there were 2 (0.5%) patients with Grade 4 events. There were no patients with Grade 3 or 5 events of T1DM.</p> <p>In the cervical cancer population in KNA18 pembrolizumab in combination with chemoradiotherapy (n=528), there was 1 (0.2%) patient with a Grade 4 event of T1DM. There were no patients with a Grade 3 or Grade 5 event of T1DM. <u>Pembrolizumab SC</u></p> <p>In the NSCLC population in MK-3475A-D77 pembrolizumab administered as MK-3475A subcutaneously in combination with chemotherapy (n=251), there was 1 (0.4%) patient with a Grade 3 event of T1DM. There were no patients with a Grade 4 or Grade 5 event of T1DM.</p>
--	--

Table SVII.3.1.8: Details of Important Identified Risk: Immune-Mediated Adverse Reactions- Endocrinopathies -Type 1 Diabetes Mellitus

	<p>In the global safety database through 31-DEC-2017, (pembrolizumab monotherapy trials only), there was 1 Grade 5 clinical trial report of T1DM.</p> <p>Review of pembrolizumab clinical trial data from ongoing studies and postmarketing data regarding immune-mediated endocrinopathies- Type I diabetes mellitus, including fatal cases, is consistent with the risk as characterized in this RMP.</p>
<u>Risk Factors and Risk Groups:</u>	No specific risk factors for Type I DM associated with pembrolizumab have been identified.
<u>Preventability:</u>	Although the development of Type I DM cannot be completely prevented; patients should be monitored for hyperglycemia or other signs and symptoms of diabetes. Insulin should be administered for type 1 diabetes, and pembrolizumab should be withheld in cases of Grade 3 hyperglycaemia until metabolic control is achieved.
<u>Impact on the Risk-Benefit Balance of the Product:</u>	Diabetes mellitus and diabetic ketoacidosis can be life-threatening or fatal in individual patients. Given the fatal outcome of untreated cancer, the risk of T1DM is outweighed by the potential benefit of pembrolizumab treatment.
<u>Public Health Impact:</u>	This risk has minimal public health impact outside its effect on individual patients.

Table SVII.3.1.9: Details of Important Potential Risk: For Hematologic Malignancies: Increased Risk of Severe Complications of Allogeneic Stem Cell Transplantation (SCT) in Patients Who Have Previously Received Pembrolizumab

Important Potential Risk:	For hematologic malignancies: increased risk of severe complications of allogeneic stem cell transplantation (SCT) in patients who have previously received pembrolizumab
<u>Potential Mechanisms:</u>	The safety and efficacy of HSCT may be different in patients previously exposed to PD-1 inhibitors, given their immunomodulatory mechanism and their prolonged clinical activity. Specifically, residual PD-1 inhibition peri- and post-SCT could enhance allogeneic T-cell responses, which could augment the graft-versus-tumor (GVT) effect but also increase the incidence or severity of graft-versus-host disease (GVHD) and other immune complications of SCT.
<u>Evidence Source(s) and Strength of Evidence:</u>	Review of pembrolizumab literature regarding increased risk of severe complications of allogeneic stem cell transplantation in patients who have previously received pembrolizumab represents scientific evidence of a possible causal association with pembrolizumab exposure. Abstract [Ref. 5.4: 04FG2Z] CTD 2.7.4
Characterisation of the Risk: For hematologic malignancies: increased risk of severe complications of allogeneic stem cell transplantation (SCT) in patients who have previously received pembrolizumab	
Frequency with 95%CI	Unknown
Seriousness / Outcomes	A case series has been presented in the literature [Ref. 5.4: 04FG2Z] that describes a cohort of patients who received PD-1 inhibitors and went on to receive allogeneic SCT. An increased risk of severe complications, including graft versus host disease (GVHD) and veno-occlusive disease of the liver was noted. In addition, the US product circular for nivolumab [Ref. 5.4: 04GJ9J] describes severe complications of allogeneic SCT following prior nivolumab therapy. Fatal outcomes have been reported.
Severity and Nature of the Risk	Hepatic VOD and severe GVHD (which can include skin, liver and gastrointestinal symptoms), including fatal reports, have been reported in patients who have undergone allogeneic stem cell transplant following pembrolizumab therapy. Review of the reports does not provide sufficient evidence for a causal relationship between an increased risk of severe complications of allogeneic transplant and prior pembrolizumab therapy. Therefore, this is presented as a potential risk in this RMP.
<u>Risk Factors and Risk Groups:</u>	Patients with hematologic malignancies undergoing allogeneic SCT who were previously treated with a PD-inhibitor.
<u>Preventability:</u>	These complications are preventable if patients do not undergo allogeneic SCT following PD-1 therapy; however, hematologic malignancy patients who are candidates for allogeneic SCT have generally exhausted other treatment options.
<u>Impact on the Risk-Benefit Balance of the Product:</u>	These complications may be fatal in individual patients. However, given the fatal outcome of untreated lymphoma in patients who have exhausted other treatment alternatives and that the limited data available suggests that these complications can be medically managed in the majority of cases, the risk is outweighed by the potential benefit.
<u>Public Health Impact:</u>	This risk has minimal public health impact outside its effect on individual patients.

Table SVII.3.1.10: Details of Important Potential Risk: Graft Versus Host Disease (GVHD) After Pembrolizumab Administration in Patients With a History of Allogeneic Stem Cell Transplant (SCT)

Important Potential Risk:	GVHD after pembrolizumab administration in patients with a history of allogeneic SCT
<u>Potential Mechanisms:</u>	The anti-tumor effect of allogeneic SCT is based upon donor stem cells reconstituting an immune system, thereby allowing for a potent graft versus tumor (GVT) effect, but also creating a significant risk of GVHD. GVHD occurs due to the donor's immune system recognizing recipient major (MHC) or minor histocompatibility antigens, exacerbated by tissue damage caused by conditioning regimens. One of the major physiologic mechanisms responsible for maintaining immune tolerance in this setting is the attenuation of donor T-cell response caused by ligation of the PD1 receptor on donor T lymphocytes to its ligands on recipient tissue. Disruption of this immune axis by PD1 inhibition, while promoting GVT effect, may also increase the risk of GVHD.
<u>Evidence Source(s) and Strength of Evidence:</u>	Published literature [Ref. 5.4: 04PSFD] Postmarketing data
<u>Characterisation of the Risk:</u> GVHD after pembrolizumab administration in patients with a history of allogeneic SCT	
Frequency with 95%CI	Unknown
Seriousness / Outcomes	Published literature has suggested that a PD-1 inhibitor (nivolumab) may have induced fatal and serious GVHD given post-allogeneic SCT. Two case series, primarily involving nivolumab, have been published in the literature, suggesting that PD-1 therapy in patients who have undergone allogeneic SCT are at risk of developing GVHD. Importantly, these case series identify a risk factor of a history of GVHD after allogeneic SCT. Furthermore, it was observed that those patients who developed GVHD following PD-1-inhibitor therapy were, on average, closer to their allogeneic SCT than those who did not develop GVHD following treatment with PD-1-inhibitors.
Severity and Nature of the Risk	Currently there is very little direct experience with pembrolizumab in this setting by which to estimate any potential effect. In the clinical trial database, there are 5 subjects who received an allogeneic SCT 5 or more years before pembrolizumab treatment. Three of these had a history of GVHD with the allogeneic SCT; however, none developed GVHD after receiving pembrolizumab. Therefore, this is presented as a potential risk in this RMP.
<u>Risk Factors and Risk Groups:</u>	Patients with a history of allogeneic SCT treated with a PD-1 inhibitor.
<u>Preventability:</u>	These complications are preventable if patients do not undergo allogeneic SCT prior to PD-1 therapy. However, hematologic malignancy patients who have undergone allogeneic SCT and subsequently relapse or develop a secondary malignancy and may be candidates for pembrolizumab therapy have generally exhausted other treatment options.
<u>Impact on the Risk-Benefit Balance of the Product:</u>	These complications may be fatal in individual patients. However, given the fatal outcome of untreated lymphoma in patients who have exhausted other treatment alternatives and that the limited data available suggests that these complications can be medically managed in the majority of cases, as well as the need for therapeutic alternatives in patients who have a medical history of allogeneic SCT and then develop secondary malignancy, the risk is outweighed by the potential benefit.
<u>Public Health Impact:</u>	This risk has minimal public health impact outside its effect on individual patients.

SVII.3.2 Presentation of the Missing Information

Not applicable.

PART II: MODULE SVIII - SUMMARY OF THE SAFETY CONCERNS

Table SVIII.1: Summary of Safety Concerns

Summary of safety concerns	
Important identified risks	Immune-mediated adverse reactions
Important potential risks	For hematologic malignancies: increased risk of severe complications of allogeneic stem cell transplantation (SCT) in patients who have previously received pembrolizumab Graft versus host disease (GVHD) after pembrolizumab administration in patients with a history of allogeneic stem cell transplant (SCT)
Missing information	None

PART III: PHARMACOVIGILANCE PLAN (INCLUDING POST-AUTHORISATION SAFETY STUDIES)

III.1 Routine Pharmacovigilance Activities

The MAH maintains systems and standard practices for routine pharmacovigilance activities to collect reports of suspected adverse reactions (including spontaneous reports, reports from clinical studies, reports of pregnancy/lactation exposures, overdoses and medication errors); prepare reports for regulatory authorities (e.g. individual case safety reports, Periodic Safety Update Reports (PSURs), etc.), and maintain continuous monitoring of the safety profile of approved products (including signal detection, issue evaluation, updating of labeling, and liaison with regulatory authorities). The MAH maintains a Pharmacovigilance System Master File which contains details of these systems and standard practices.

MAH-sponsored clinical trials are being conducted in multiple different tumor types including but not limited to: melanoma, non-small cell lung cancer, hematologic malignancies, bladder tumors, and multiple other solid tumor types that will continue to inform the safety profile of pembrolizumab. All of these on-going and planned clinical trials will continue to provide valuable data and help to inform the MAH regarding ongoing safety concerns and to identify future safety signals.

Routine Pharmacovigilance Activities Beyond Adverse Reactions Reporting and Signal Detection:

Table III.1.1: Overview of Pharmacovigilance Actions

Areas Requiring Further Investigation	Proposed Routine and Additional Pharmacovigilance Activities	Objectives
Important Identified risk: Immune-Mediated Adverse Reactions		
In order to monitor for and better characterize the occurrence of immune-mediated adverse reactions the MAH monitors and evaluates reports of immune-mediated adverse reactions received in the postmarketing and clinical environment.	Routine pharmacovigilance Additional pharmacovigilance including: <ul style="list-style-type: none"> • Safety monitoring in all ongoing MAH-sponsored clinical trials for pembrolizumab in various tumor types 	To monitor, identify and evaluate reports of immune-mediated adverse reactions in patients treated with pembrolizumab
Important Potential Risk: For hematologic malignancies: increased risk of severe complications of allogeneic SCT in patients who have previously received pembrolizumab		
In order to monitor for and better characterize the occurrence (for hematologic malignancies) of an increased risk of severe complications of allogeneic SCT in patients who have previously received pembrolizumab, the MAH monitors and evaluates reports of severe complications of allogeneic SCT in patients who have previously received pembrolizumab from both the postmarketing and clinical environment.	Routine pharmacovigilance Additional pharmacovigilance including: <ul style="list-style-type: none"> • Safety monitoring in the ongoing HL trial (KN204) 	To monitor, identify and evaluate for hematologic malignancies: reports of severe complications of allogeneic SCT in patients who have previously received pembrolizumab
Important Potential Risk: Graft versus host disease (GVHD) after pembrolizumab administration in patients with a history of allogeneic stem cell transplant (SCT)		
In order to monitor for and better characterize the occurrence of GVHD after pembrolizumab administration in patients with a history of allogeneic SCT, the MAH monitors and evaluates reports of GVHD after pembrolizumab administration in patients with a history of allogeneic SCT from both the postmarketing and clinical trial environment.	Routine pharmacovigilance Additional pharmacovigilance including: <ul style="list-style-type: none"> ▪ Safety monitoring in all ongoing MAH-sponsored clinical trials for pembrolizumab in various tumor types 	To monitor, identify and evaluate reports of GVHD after pembrolizumab administration in patients with a history of allogeneic SCT

Other Forms of Routine Pharmacovigilance Activities for Safety Concerns:

Not applicable.

III.2 Additional Pharmacovigilance Activities

All ongoing and planned additional pharmacovigilance activities are described below.

III.3 Summary Table of Additional Pharmacovigilance Activities

On-Going and Planned Additional Pharmacovigilance Activities

There are no ongoing or planned additional pharmacovigilance studies that are required for pembrolizumab.

PART IV: PLANS FOR POST-AUTHORISATION EFFICACY STUDIES

Table IV.1: Planned and On-Going Post-Authorisation Efficacy Studies that are Conditions of the Marketing Authorisation or that are Specific Obligations

Study Status	Summary of Objectives	Efficacy Uncertainties Addressed	Milestones	Due Date
Efficacy studies which are conditions of the marketing authorization				
A Phase III, Randomized, Open-label, Clinical Trial to Compare Pembrolizumab with Brentuximab Vedotin in Subjects with Relapsed or Refractory Classical Hodgkin Lymphoma (KN204) (on-going)	To compare overall survival (OS), progression free survival (PFS) and objective response rate (ORR) of pembrolizumab when compared to Brentuximab Vedotin in subjects with relapsed or refractory cHL and to examine the safety and tolerability between treatment groups.	Long term efficacy and safety	Final study report	4Q 2026
Adjuvant immunotherapy with anti-PD-1 monoclonal antibody Pembrolizumab (MK-3475) versus placebo after complete resection of high-risk Stage III melanoma: A randomized, double-blind Phase 3 trial of the EORTC Melanoma Group (KN054) (on-going)	To prospectively assess whether post-operative adjuvant therapy with pembrolizumab improves recurrence-free survival (RFS) as compared to placebo in high-risk patients with complete resection of Stage IIIA (> 1 mm metastasis), IIIB and IIIC melanoma. To prospectively assess whether in the subgroup of patients with PD-L1- positive tumor expression, pembrolizumab improves recurrence-free survival as compared to placebo (primary endpoints); distant metastasis free survival (DMFS) and overall survival (OS) in all-subjects and subjects with PD-L1-positive tumors (secondary endpoints for final study report)	Long term efficacy and safety	Final Study Report	4Q 2027
A randomized, placebo-controlled, parallel-group, crossover/rechallenge, multi-center study of adjuvant pembrolizumab in participants 12 years of age and older with resected Stage IIB or IIC cutaneous melanoma (KN716) (on-going)	To compare RFS, DMFS and OS between pembrolizumab and placebo in patients with resected Stage IIB or IIC cutaneous melanoma	Long term efficacy and safety	Clinical Study Report	4Q 2028

Table IV.1: Planned and On-Going Post-Authorisation Efficacy Studies that are Conditions of the Marketing Authorisation or that are Specific Obligations

Study Status	Summary of Objectives	Efficacy Uncertainties Addressed	Milestones	Due Date
A randomized, Phase 3 trial with pembrolizumab versus placebo for patients with early stage NSCLC after resection and completion of standard adjuvant therapy (KN091) (on-going)	To compare OS between pembrolizumab and placebo in patients with early stage NSCLC after resection and completion of standard adjuvant therapy To assess data on treatment post-progression, and particularly on the uptake and activity of anti-PD(L)1 in patients previously treated with adjuvant pembrolizumab	Long term efficacy and safety	Final Study Report	3Q2026

PART V: RISK MINIMISATION MEASURES (INCLUDING EVALUATION OF THE EFFECTIVENESS OF RISK MINIMISATION ACTIVITIES)

Risk Minimisation Plan

V.1 Routine Risk Minimisation Measures

Table V.1.1: Description of Routine Risk Minimisation Measures by Safety Concern

Safety Concern	Routine Risk Minimisation Activities
Immune-Mediated Adverse Reactions	<p><u>Objective of the Risk Minimisation Measures:</u> The objective is to inform prescribers of the risk of immune-mediated adverse reactions and to provide appropriate advice to minimize the risk in clinical practice.</p> <p><u>Text in SmPC:</u> Sections 4.2, 4.4, 4.8 of the SmPC: Immune-mediated adverse reactions</p>
For Hematologic Malignancies: Increased risk of severe complications of allogeneic stem cell transplantation (SCT) in patients who have previously received pembrolizumab	<p><u>Objective of the Risk Minimisation Measures:</u> The objective is to inform prescribers of the potential risk of: For hematologic malignancies: Increased risk of severe complications of allogeneic SCT in patients who have previously received pembrolizumab and to provide appropriate advice to minimize the risk in clinical practice.</p> <p><u>Text in SmPC:</u> Sections 4.4, 4.8 of the SmPC</p>
Graft versus host disease (GVHD) after pembrolizumab administration in patients with a history of allogeneic stem cell transplantation (SCT)	<p><u>Objective of the Risk Minimisation Measures:</u> The objective is to inform prescribers of the potential risk of GVHD after pembrolizumab administration in patients with a history of allogeneic SCT and to provide appropriate information to assist in early recognition and treatment in clinical practice.</p> <p><u>Text in SmPC:</u> Section 4.4 of the SmPC</p>

V.2 Additional Risk Minimisation Measures

Table V.2.1: Description of Additional Risk Minimisation Measures by Safety Concern

Safety Concern	Additional Risk Minimisation Activities
<p>Immune-Mediated Adverse Reactions</p>	<p>Objective and Rationale: Receipt of educational material will reinforce communication of the information in the SmPC from prescribers to their patients and are intended to help minimize the risk of immune-mediated reactions in clinical practice. Patient educational material in the form of patient card will enable patient awareness of side effects and early recognition of immune mediated events supporting risk Minimisation.</p> <p>Proposed Actions: Educational materials:</p> <ul style="list-style-type: none"> • Patient Card <p>Healthcare providers will distribute the Patient Card to the patients/caregivers to educate them about symptoms of immune-mediated adverse reactions.</p> <p>Patients will receive:</p> <ul style="list-style-type: none"> • Information about key symptoms that need to be reported immediately to the physician/nurse. • Instruction on the importance of carrying the wallet-sized Patient Card at all times, to show it at medical visits to healthcare professionals other than the prescriber.
	<p>How Effectiveness of the Risk Minimisation Measures for the Safety Concern Will be Measured: Implementation:</p> <ul style="list-style-type: none"> • Communication via direct mail to oncologists, oncology nurses and oncology pharmacists who treat patients with the approved indication(s). The records of the mailer will be used to track the risk Minimisation activity. • Training of the local country staff on the patient card and on the need to provide the material to new prescribers will be completed. Records of the training will be maintained. <p>Clinical outcomes:</p> <ul style="list-style-type: none"> • The MAH's educational plan will encourage patients to report signs and symptoms associated with immune-mediated reactions to their HCPs. • Information from ongoing pharmacovigilance activities will assess appropriate recognition and management of immune-mediated adverse reactions and that the events observed in real world use are consistent with the cases observed in the clinical trial program and characterized in the SmPC and patient card. <p>Criteria for Judging the Success of the Proposed Risk Minimisation Measure(s): Clinical features of reported Adverse Drug Reactions (ADRs) are reviewed to ensure consistency with both the known safety profile of pembrolizumab and also with appropriate medical management guidelines in the SmPC. Upon review of the data, appropriate measures will be considered if new information is obtained that indicates that additional risk minimisation measures are warranted.</p> <p>Planned Date(s) for Assessment: Proposed review period:</p> <ul style="list-style-type: none"> • Data will be reviewed as part of routine pharmacovigilance processes and will be presented in the PSUR if clinically meaningful new safety information is identified.

Table V.2.1: Description of Additional Risk Minimisation Measures by Safety Concern

Safety Concern	Additional Risk Minimisation Activities
For Hematologic Malignancies: Increased risk of severe complications of allogeneic stem cell transplantation (SCT) in patients who have previously received pembrolizumab	Not applicable
Graft versus host disease (GVHD) after pembrolizumab administration in patients with a history of allogeneic stem cell transplant (SCT)	Not applicable

V.3 Summary of Risk Minimisation Measures

Table V.3.1: Summary Table of Pharmacovigilance Activities and Risk Minimisation Activities by Safety Concern

Safety Concern	Risk minimisation Measures	Pharmacovigilance Activities
Important Identified Risks: Immune-Mediated Adverse Reactions		
Immune-mediated adverse reactions	Routine risk minimisation measures: <ul style="list-style-type: none"> ▪ The risk of the immune-mediated adverse reactions associated with the use of pembrolizumab is described in the SmPC, Section 4.2, 4.4, 4.8 and appropriate advice is provided to the prescriber to minimize the risk. 	Routine pharmacovigilance activities
	Additional risk minimisation measures: <ul style="list-style-type: none"> ▪ Patient card 	Additional pharmacovigilance including: <ul style="list-style-type: none"> • Safety monitoring in all ongoing MAH-sponsored clinical trials for pembrolizumab in various tumor types

Table V.3.1: Summary Table of Pharmacovigilance Activities and Risk Minimisation Activities by Safety Concern

Safety Concern	Risk minimisation Measures	Pharmacovigilance Activities
Important Potential Risks		
<p>For hematologic malignancies: increased risk of severe complications of allogeneic SCT in patients who have previously received pembrolizumab</p>	<p>Routine risk minimisation measures:</p> <ul style="list-style-type: none"> ▪ For Hematologic malignancies: the increased risk of severe complications of allogeneic SCT in patients who have previously received pembrolizumab is described in the SmPC, Section 4.4, 4.8 and appropriate advice is provided to the prescriber to minimize the risk. <p>No additional risk minimisation measures warranted</p>	<p>Routine pharmacovigilance activities</p> <p>Additional pharmacovigilance including:</p> <ul style="list-style-type: none"> ▪ Safety monitoring in the ongoing HL trial (KN204).
<p>GVHD after pembrolizumab administration in patients with a history of allogeneic SCT</p>	<p>Routine risk minimisation measures:</p> <ul style="list-style-type: none"> ▪ GVHD after pembrolizumab administration in patients with a history of allogeneic SCT is described in the SmPC, Section 4.4 and appropriate advice is provided to the prescriber to minimize the risk. <p>No additional risk minimisation measures warranted</p>	<p>Routine pharmacovigilance activities</p> <p>Additional pharmacovigilance including:</p> <ul style="list-style-type: none"> • Safety monitoring in all ongoing MAH-sponsored clinical trials for pembrolizumab in various tumor types

PART VI: SUMMARY OF THE RISK MANAGEMENT PLAN BY PRODUCT

Summary of risk management plan for pembrolizumab

This is a summary of the risk management plan (RMP) for pembrolizumab. The RMP details important risks of pembrolizumab, how these risks can be minimised, and how more information will be obtained about pembrolizumab's risks and uncertainties (missing information).

Pembrolizumab's Summary of product characteristics (SmPC) and its package leaflet give essential information to healthcare professionals and patients on how pembrolizumab should be used.

I. The Medicine and What it is Used For

Pembrolizumab Intravenous (IV)

Melanoma

Pembrolizumab as monotherapy is indicated for the treatment of adults and adolescents aged 12 years and older with advanced (unresectable or metastatic) melanoma.

Pembrolizumab as monotherapy is indicated for the adjuvant treatment of adults and adolescents aged 12 years and older with Stage IIB, IIC or III melanoma and who have undergone complete resection.

Non-small cell lung carcinoma (NSCLC)

Pembrolizumab in combination with platinum-containing chemotherapy as neoadjuvant treatment, and then continued as monotherapy as adjuvant treatment, is indicated for the treatment of resectable non-small cell lung carcinoma at high risk of recurrence in adults.

Pembrolizumab as monotherapy is indicated for the adjuvant treatment of adults with non-small cell lung carcinoma who are at high risk of recurrence following complete resection and platinum-based chemotherapy.

Pembrolizumab as monotherapy is indicated for the first-line treatment of metastatic non-small cell lung carcinoma (NSCLC) in adults whose tumours express PD-L1 with a $\geq 50\%$ tumour proportion score (TPS) with no EGFR or ALK positive tumour mutations.

Pembrolizumab, in combination with pemetrexed and platinum chemotherapy, is indicated for the first-line treatment of metastatic non-squamous NSCLC in adults whose tumours have no EGFR or ALK positive mutations.

Pembrolizumab, in combination with carboplatin and either paclitaxel or nab-paclitaxel, is indicated for the first-line treatment of metastatic squamous NSCLC in adults.

Pembrolizumab as monotherapy is indicated for the treatment of locally advanced or metastatic NSCLC in adults whose tumours express PD-L1 with a $\geq 1\%$ TPS and who have received at least one prior chemotherapy regimen. Patients with EGFR or ALK positive tumour mutations should also have received targeted therapy before receiving pembrolizumab.

Malignant pleural mesothelioma (MPM)

Pembrolizumab, in combination with pemetrexed and platinum chemotherapy, is indicated for the first line treatment of adults with unresectable non-epithelioid malignant pleural mesothelioma.

Classical Hodgkin lymphoma (cHL)

Pembrolizumab as monotherapy is indicated for the treatment of adult and paediatric patients aged 3 years and older with relapsed or refractory classical Hodgkin Lymphoma (cHL) who have failed autologous stem cell transplant (ASCT) or following at least two prior therapies when ASCT is not a treatment option.

Urothelial carcinoma

Pembrolizumab as monotherapy is indicated for the treatment of locally advanced or metastatic urothelial carcinoma in adults who have received prior platinum-containing chemotherapy.

Pembrolizumab as monotherapy is indicated for the treatment of locally advanced or metastatic urothelial carcinoma in adults who are not eligible for cisplatin-containing chemotherapy and whose tumours express PD-L1 with a combined positive score (CPS) ≥ 10 .

Pembrolizumab, in combination with enfortumab vedotin is indicated for the first-line treatment of unresectable or metastatic urothelial carcinoma in adults.

Head and neck squamous cell carcinoma (HNSCC)

Pembrolizumab as monotherapy is indicated for the treatment of resectable locally advanced head and neck squamous cell carcinoma (HNSCC) as neoadjuvant treatment, continued as adjuvant treatment in combination with radiation therapy with or without concomitant cisplatin and then as monotherapy in adults whose tumours express PD-L1 with a CPS ≥ 1 .

Pembrolizumab, as monotherapy or in combination with platinum and 5-fluorouracil (5-FU) chemotherapy, is indicated for the first-line treatment of metastatic or unresectable recurrent HNSCC in adults whose tumours express PD-L1 with a CPS ≥ 1 .

Pembrolizumab as monotherapy is indicated for the treatment of recurrent or metastatic HNSCC in adults whose tumours express PD-L1 with a $\geq 50\%$ TPS and progressing on or after platinum-containing chemotherapy.

Renal cell carcinoma (RCC)

Pembrolizumab, in combination with axitinib, is indicated for the first-line treatment of advanced renal cell carcinoma (RCC) in adults.

Pembrolizumab, in combination with lenvatinib, is indicated for the first-line treatment of advanced renal cell carcinoma (RCC) in adults.

Pembrolizumab as monotherapy is indicated for the adjuvant treatment of adults with renal cell carcinoma at increased risk of recurrence following nephrectomy, or following nephrectomy and resection of metastatic lesions.

Microsatellite instability high (MSI-H) or mismatch repair deficient (dMMR) cancers

Colorectal cancer (CRC)

Pembrolizumab as monotherapy is indicated for adults with MSI-H or dMMR colorectal cancer in the following settings:

- first-line treatment of metastatic colorectal cancer;
- treatment of unresectable or metastatic colorectal cancer after previous fluoropyrimidine-based combination therapy.

Non-colorectal cancers

Pembrolizumab as monotherapy is indicated for the treatment of the following MSI-H or dMMR tumours in adults with:

- advanced or recurrent endometrial carcinoma, who have disease progression on or following prior treatment with a platinum-containing therapy in any setting and who are not candidates for curative surgery or radiation;
- unresectable or metastatic gastric, small intestine, or biliary cancer, who have disease progression on or following at least one prior therapy.

Oesophageal carcinoma

Pembrolizumab, in combination with platinum and fluoropyrimidine based chemotherapy, is indicated for the first-line treatment of locally advanced unresectable or metastatic carcinoma of the oesophagus in adults whose tumours express PD-L1 with a CPS \geq 10

Triple negative breast cancer (TNBC)

Pembrolizumab, in combination with chemotherapy as neoadjuvant treatment, and then continued as monotherapy as adjuvant treatment after surgery, is indicated for the treatment of

adults with locally advanced, or early-stage triple-negative breast cancer (TNBC) at high risk of recurrence.

Pembrolizumab, in combination with chemotherapy, is indicated for the treatment of locally recurrent unresectable or metastatic triple-negative breast cancer (TNBC) in adults whose tumors express PD-L1 with a CPS ≥ 10 and who have not received prior chemotherapy for metastatic disease.

Endometrial carcinoma (EC)

Pembrolizumab, in combination with carboplatin and paclitaxel, is indicated for the first-line treatment of primary advanced or recurrent endometrial carcinoma in adults who are candidates for systemic therapy.

Pembrolizumab, in combination with lenvatinib, is indicated for the treatment of advanced or recurrent endometrial carcinoma in adults who have disease progression on or following prior treatment with a platinum-containing therapy in any setting and who are not candidates for curative surgery or radiation.

Cervical cancer

Pembrolizumab, in combination with chemoradiotherapy (external beam radiation therapy followed by brachytherapy), is indicated for the treatment of FIGO 2014 Stage III - IVA locally advanced cervical cancer in adults who have not received prior definitive therapy.

Pembrolizumab, in combination with chemotherapy with or without bevacizumab, is indicated for the treatment of persistent, recurrent, or metastatic cervical cancer in adults whose tumours express PD-L1 with a CPS ≥ 1 .

Gastric or gastro-oesophageal junction adenocarcinoma

Pembrolizumab, in combination with trastuzumab, fluoropyrimidine- and platinum-containing chemotherapy, is indicated for the first-line treatment of locally advanced unresectable or metastatic HER2-positive gastric or gastro-oesophageal junction adenocarcinoma in adults whose tumours express PD-L1 with a CPS ≥ 1 .

Pembrolizumab, in combination with fluoropyrimidine- and platinum-containing chemotherapy, is indicated for the first-line treatment of locally advanced unresectable or metastatic HER2-negative gastric or gastro-oesophageal junction adenocarcinoma in adults whose tumours express PD-L1 with a CPS ≥ 1 .

Biliary tract carcinoma (BTC)

Pembrolizumab, in combination with gemcitabine and cisplatin, is indicated for the first-line treatment of locally advanced unresectable or metastatic biliary tract carcinoma in adults.

It contains 100 mg liquid pembrolizumab in a 10 mL single-use vial as the active substance and it is given by intravenous infusion.

The Recommended Dose of Pembrolizumab IV is:

- 200 mg every 3 weeks or 400 mg every 6 weeks administered as an intravenous infusion over 30 minutes.
- The recommended dose of pembrolizumab as monotherapy in paediatric patients aged 3 years and older with cHL or patients aged 12 years and older with melanoma is 2 mg/kg bodyweight (bw) (up to a maximum of 200 mg), every 3 weeks administered as an intravenous infusion over 30 minutes

Pembrolizumab Subcutaneous (SC)

Melanoma

Pembrolizumab as monotherapy is indicated for the treatment of adults with advanced (unresectable or metastatic) melanoma.

Pembrolizumab as monotherapy is indicated for the adjuvant treatment of adults with Stage IIB, IIC or III melanoma and who have undergone complete resection.

Non-small cell lung carcinoma (NSCLC)

Pembrolizumab, in combination with platinum-containing chemotherapy as neoadjuvant treatment, and then continued as monotherapy as adjuvant treatment, is indicated for the treatment of resectable non-small cell lung carcinoma at high risk of recurrence in adults.

Pembrolizumab as monotherapy is indicated for the adjuvant treatment of adults with non-small cell lung carcinoma who are at high risk of recurrence following complete resection and platinum-based chemotherapy.

Pembrolizumab as monotherapy is indicated for the first line treatment of metastatic non-small cell lung carcinoma in adults whose tumours express PD L1 with a $\geq 50\%$ tumour proportion score (TPS) with no EGFR or ALK positive tumour mutations.

Pembrolizumab, in combination with pemetrexed and platinum chemotherapy, is indicated for the first-line treatment of metastatic non squamous non-small cell lung carcinoma in adults whose tumours have no EGFR or ALK positive mutations.

Pembrolizumab, in combination with carboplatin and either paclitaxel or nab paclitaxel, is indicated for the first line treatment of metastatic squamous non-small cell lung carcinoma in adults.

Pembrolizumab as monotherapy is indicated for the treatment of locally advanced or metastatic non-small cell lung carcinoma in adults whose tumours express PD L1 with a $\geq 1\%$ TPS and who have received at least one prior chemotherapy regimen. Patients with EGFR or ALK positive tumour mutations should also have received targeted therapy before receiving Pembrolizumab.

Classical Hodgkin lymphoma (cHL)

Pembrolizumab as monotherapy is indicated for the treatment of adults with relapsed or refractory classical Hodgkin lymphoma who have failed autologous stem cell transplant (ASCT) or following at least two prior therapies when ASCT is not a treatment option.

Urothelial carcinoma

Pembrolizumab, in combination with enfortumab vedotin, is indicated for the first-line treatment of unresectable or metastatic urothelial carcinoma in adults.

Pembrolizumab as monotherapy is indicated for the treatment of locally advanced or metastatic urothelial carcinoma in adults who have received prior platinum containing chemotherapy.

Pembrolizumab as monotherapy is indicated for the treatment of locally advanced or metastatic urothelial carcinoma in adults who are not eligible for cisplatin containing chemotherapy and whose tumours express PD L1 with a combined positive score (CPS) ≥ 10 .

Head and neck squamous cell carcinoma (HNSCC)

Pembrolizumab as monotherapy is indicated for the treatment of resectable locally advanced head and neck squamous cell carcinoma (HNSCC) as neoadjuvant treatment, continued as adjuvant treatment in combination with radiation therapy with or without concomitant cisplatin and then as monotherapy in adults whose tumours express PD-L1 with a CPS ≥ 1 .

Pembrolizumab, as monotherapy or in combination with platinum and 5 fluorouracil (5 FU) chemotherapy, is indicated for the first line treatment of metastatic or unresectable recurrent head and neck squamous cell carcinoma in adults whose tumours express PD L1 with a CPS ≥ 1 .

Pembrolizumab as monotherapy is indicated for the treatment of recurrent or metastatic head and neck squamous cell carcinoma in adults whose tumours express PD L1 with a $\geq 50\%$ TPS and progressing on or after platinum containing chemotherapy.

Renal cell carcinoma (RCC)

Pembrolizumab, in combination with axitinib, is indicated for the first line treatment of advanced renal cell carcinoma in adults.

Pembrolizumab, in combination with lenvatinib, is indicated for the first line treatment of advanced renal cell carcinoma in adults.

Pembrolizumab as monotherapy is indicated for the adjuvant treatment of adults with renal cell carcinoma at increased risk of recurrence following nephrectomy, or following nephrectomy and resection of metastatic lesions.

Microsatellite instability high (MSI-H) or mismatch repair deficient (dMMR) cancers

Colorectal cancer (CRC)

Pembrolizumab as monotherapy is indicated for adults with MSI-H or dMMR colorectal cancer in the following settings:

- first line treatment of metastatic colorectal cancer;
- treatment of unresectable or metastatic colorectal cancer after previous fluoropyrimidine based combination therapy.

Non-colorectal cancers

Pembrolizumab as monotherapy is indicated for the treatment of the following MSI-H or dMMR tumours in adults with:

- advanced or recurrent endometrial carcinoma, who have disease progression on or following prior treatment with a platinum containing therapy in any setting and who are not candidates for curative surgery or radiation;
- unresectable or metastatic gastric, small intestine, or biliary cancer, who have disease progression on or following at least one prior therapy.

Oesophageal carcinoma

Pembrolizumab, in combination with platinum and fluoropyrimidine based chemotherapy, is indicated for the first-line treatment of locally advanced unresectable or metastatic carcinoma of the oesophagus in adults whose tumours express PD L1 with a CPS \geq 10.

Triple negative breast cancer (TNBC)

Pembrolizumab, in combination with chemotherapy as neoadjuvant treatment, and then continued as monotherapy as adjuvant treatment after surgery, is indicated for the treatment of adults with locally advanced, or early stage triple negative breast cancer at high risk of recurrence.

Pembrolizumab, in combination with chemotherapy, is indicated for the treatment of locally recurrent unresectable or metastatic triple negative breast cancer in adults whose tumours express PD L1 with a CPS \geq 10 and who have not received prior chemotherapy for metastatic disease.

Endometrial carcinoma (EC)

Pembrolizumab, in combination with carboplatin and paclitaxel, is indicated for the first-line treatment of primary advanced or recurrent endometrial carcinoma in adults who are candidates for systemic therapy.

Pembrolizumab, in combination with lenvatinib, is indicated for the treatment of advanced or recurrent endometrial carcinoma in adults who have disease progression on or following prior treatment with a platinum containing therapy in any setting and who are not candidates for curative surgery or radiation.

Cervical cancer

Pembrolizumab, in combination with chemoradiotherapy (external beam radiation therapy followed by brachytherapy), is indicated for the treatment of FIGO 2014 Stage III - IVA locally advanced cervical cancer in adults who have not received prior definitive therapy.

Pembrolizumab, in combination with chemotherapy with or without bevacizumab, is indicated for the treatment of persistent, recurrent, or metastatic cervical cancer in adults whose tumours express PD L1 with a CPS \geq 1.

Gastric or gastro-oesophageal junction (GEJ) adenocarcinoma

Pembrolizumab, in combination with trastuzumab, fluoropyrimidine and platinum-containing chemotherapy, is indicated for the first-line treatment of locally advanced unresectable or metastatic HER2-positive gastric or gastro-oesophageal junction adenocarcinoma in adults whose tumours express PD-L1 with a CPS \geq 1.

Pembrolizumab, in combination with fluoropyrimidine and platinum-containing chemotherapy, is indicated for the first-line treatment of locally advanced unresectable or metastatic HER2-negative gastric or gastro-oesophageal junction adenocarcinoma in adults whose tumours express PD L1 with a CPS \geq 1.

Biliary tract carcinoma (BTC)

Pembrolizumab, in combination with gemcitabine and cisplatin, is indicated for the first-line treatment of locally advanced unresectable or metastatic biliary tract carcinoma in adults.

Each single-use vial of 2.4 mL contains 395 mg of pembrolizumab, or 4.8 mL contains 790 mg of pembrolizumab as the active substance and it is given by subcutaneous injection.

The Recommended Dose of Pembrolizumab SC is:

- KEYTRUDA is administered as a subcutaneous injection into the abdomen or thigh over approximately 1-2 minutes. The recommended dose of KEYTRUDA solution for injection in adults is either:
 - 2.4 mL of KEYTRUDA (395 mg) every 3 weeks over 1 minute.
 - 4.8 mL of KEYTRUDA (790 mg) every 6 weeks over 2 minutes.

Further information about the evaluation of pembrolizumab's benefits can be found in pembrolizumab's European Public Assessment Report (EPAR), including in its plain-language summary, available on the EMA website, under the medicine's webpage link to product's EPAR summary landing page on the EMA webpage at the following link:

<https://www.ema.europa.eu/en/medicines/human/EPAR/keytruda>

II. Risks Associated With the Medicine and Activities to Minimise or Further Characterise the Risks

Important risks of pembrolizumab, together with measures to minimise such risks and the proposed studies for learning more about pembrolizumab's risks, are outlined below.

Measures to minimise the risks identified for medicinal products can be:

- Specific information, such as warnings, precautions, and advice on correct use, in the package leaflet and SmPC addressed to patients and healthcare professionals;
- Important advice on the medicine's packaging;
- The authorised pack size - the amount of medicine in a pack is chosen so to ensure that the medicine is used correctly;
- The medicine's legal status - the way a medicine is supplied to the patient (e.g. with or without prescription) can help to minimise its risks.

Together, these measures constitute *routine risk minimisation* measures.

In the case of pembrolizumab, these measures are supplemented with *additional risk minimisation measures* mentioned under relevant important risks, below. In addition to these measures, information about adverse reactions is collected continuously and regularly analysed, including PSUR assessment - so that immediate action can be taken as necessary. These measures constitute *routine pharmacovigilance activities*. If important information that may affect the safe use of pembrolizumab is not yet available, it is listed under 'missing information' below.

II.A List of Important Risks and Missing Information

Important risks of pembrolizumab are risks that need special risk management activities to further investigate or minimise the risk, so that the medicinal product can be safely administered. Important risks can be regarded as identified or potential. Identified risks are concerns for which there is sufficient proof of a link with the use of pembrolizumab. Potential risks are concerns for which an association with the use of this medicine is possible based on available data, but this association has not been established yet and needs further evaluation. Missing information refers to information on the safety of the medicinal product that is currently missing and needs to be collected (e.g. on the long-term use of the medicine).

Pembrolizumab has been on the market since July of 2015. Risk minimization activities recommending specific clinical measures to address the risks have become fully integrated into standard clinical practice, such as inclusion into treatment protocols or clinical guidelines. The identified risks in the EU RMP are: Immune-mediated adverse reactions.

The potential risks in the EU RMP include: an increased risk of severe complications of allogeneic stem cell transplantation (SCT) in patients who have previously received pembrolizumab and Graft versus host disease (GVHD) after pembrolizumab administration in patients with a history of allogeneic stem cell transplant (SCT).

Table II.A.1: List of Important Risks and Missing Information

List of Important Risks and Missing Information	
Important identified risks	Immune-mediated adverse reactions
Important potential risks	For hematologic malignancies: increased risk of severe complications of allogeneic stem cell transplantation (SCT) in patients who have previously received pembrolizumab Graft versus host disease (GVHD) after pembrolizumab administration in patients with a history of allogeneic stem cell transplant (SCT)
Missing information	None

II.B Summary of Important Risks

Table II.B.1: Important Identified Risk: Immune-Mediated Adverse Reactions

Evidence for linking the risk to the medicine	Review of pembrolizumab clinical trial data, post-marketing experience and literature regarding immune-mediated adverse reactions represent sufficient evidence of a causal association with pembrolizumab exposure. pembrolizumab KN001 Database Cutoff Date: 18APR2014 pembrolizumab KN001 Database Cutoff Date for Lung: 23JAN2015 pembrolizumab KN002 Database Cutoff Date: 28FEB2015 pembrolizumab KN006 Database Cutoff Date: 03MAR2015 pembrolizumab KN010 Database Cutoff Date: 30SEP2015 pembrolizumab KN013 Database Cutoff Date for Hodgkin Lymphoma: 28SEP2018 pembrolizumab KN024 Database Cutoff Date: 10JUL2017 pembrolizumab KN087 Database Cutoff Date: 21MAR2019 pembrolizumab KN045 Database Cutoff Date: 07SEP2016 pembrolizumab KN052 Database Cutoff Date: 01SEP2016 pembrolizumab KN021 Database Cutoff Date Cohort A: 07NOV2016, Cohort G/C: 31MAY2017 pembrolizumab KN189 Database Cutoff Date: 08NOV2017 pembrolizumab KN040 Database Cutoff Date: 15MAY2017 pembrolizumab KN012 Database Cutoff Date: 26APR2016 pembrolizumab KN055 Database Cutoff Date: 22APR2016 pembrolizumab KN054 Database Cutoff Date: 02OCT2017 pembrolizumab KN407 Database Cutoff Date: 03APR2018 pembrolizumab KN426 Database Cutoff Date: 24AUG2018 pembrolizumab KN048 Database Cutoff Date: 13JUN2018
---	---

Table II.B.1: Important Identified Risk: Immune-Mediated Adverse Reactions

	<p>pembrolizumab KN042 Database Cutoff Date: 26FEB2018</p> <p>pembrolizumab KN177 Database Cutoff Date: 19FEB2020</p> <p>pembrolizumab KN204 Database Cutoff Date: 16JAN2020</p> <p>pembrolizumab KN590 Database Cutoff Date: 02JUL2020</p> <p>pembrolizumab KN355 Database Cutoff Date: 11DEC2019</p> <p>pembrolizumab KN581 Database Cutoff Date: 28AUG2020</p> <p>pembrolizumab KN146 Database Cutoff Date: 18AUG2020</p> <p>pembrolizumab KN775 Database Cutoff Date: 26OCT2020</p> <p>pembrolizumab KN564 Database Cutoff Date: 14DEC2020</p> <p>pembrolizumab KN158 Database Cutoff Date Cohort K: 05OCT2020</p> <p>pembrolizumab KN164 Database Cutoff Date Cohorts A and B: 09SEP2019</p> <p>pembrolizumab KN826 Database Cutoff Date: 03MAY2021</p> <p>pembrolizumab KN522 Database Cutoff Date: 23MAR2021</p> <p>pembrolizumab KN716 Database Cutoff Date: 04DEC2020</p> <p>pembrolizumab KN811 Database Cutoff Date: 25MAY2022</p> <p>pembrolizumab KN091 Database Cutoff Date: 20SEP2021</p> <p>pembrolizumab KN859 Database Cutoff Date: 03OCT2022</p> <p>pembrolizumab KN966 Database Cutoff Date: 15DEC2022</p> <p>pembrolizumab KN671 Database Cutoff Date: 29JUL2022</p> <p>pembrolizumab KN869 Database Cutoff Date: 13MAR2023</p> <p>pembrolizumab KNA39 Database Cutoff Date: 08AUG2023</p> <p>pembrolizumab KN868 Database Cutoff Date: 16Dec2022 for dMMR participants and 06Dec2022 for pMMR participants</p> <p>pembrolizumab KNA18 Database Cutoff Date: 09JAN2023</p> <p>pembrolizumab KN483 Database Cutoff Date: 16SEP2022</p> <p>pembrolizumab KNA17 Database Cutoff Date: 21SEP2022</p> <p>pembrolizumab KN689 Database Cutoff Date: 25JUL2024</p> <p>pembrolizumab MK-3475A-D77 Database Cutoff Date: 12JUL2024</p>
--	--

Table II.B.1: Important Identified Risk: Immune-Mediated Adverse Reactions

<p>Risk factors and risk groups</p>	<p>Pneumonitis Patients with a history of non-infectious pneumonitis that required steroids or current pneumonitis were excluded from the clinical trials. These patients are considered to be a risk group for the development of pneumonitis; in the interim analysis of the KN001 NSCLC cohort, possible risk factors identified that might predispose subjects to pneumonitis were a documented history of prior thoracic radiation to the chest ($\geq 30\text{Gy}$). According to the literature, risk factors for interstitial lung disease may include occupational exposure to toxins, chest irradiation, some chemotherapies, smoking and advanced age.</p> <p>Colitis No specific risk factors for colitis and diarrhea associated with pembrolizumab were identified.</p> <p>Hepatitis Patients with moderate to severe liver dysfunction were excluded from clinical trials. No analysis of specific risk factors for immune-mediated hepatitis associated with pembrolizumab has been undertaken.</p> <p>Nephritis Patients with severe renal dysfunction were excluded from clinical trials. No specific risk factors for nephritis associated with pembrolizumab have been identified.</p> <p>Endocrinopathies No specific risk factors for endocrinopathies associated with pembrolizumab have been identified.</p>
<p>Risk minimisation measures</p>	<p>Routine risk minimisation measures:</p> <ul style="list-style-type: none"> ▪ The risk of the immune-mediated adverse reactions associated with the use of pembrolizumab is described in the SmPC, Section 4.2, 4.4, 4.8 and appropriate advice is provided to the prescriber to minimize the risk. <p>Additional risk minimisation measures:</p> <ul style="list-style-type: none"> ▪ Patient card
<p>Additional pharmacovigilance activities</p>	<p>Additional pharmacovigilance activities:</p> <ul style="list-style-type: none"> • Safety monitoring in all ongoing MAH-sponsored clinical trials for pembrolizumab in various tumor types.

Table II.B.2: Important Potential Risk: For Hematologic Malignancies: Increased Risk of Severe Complications of Allogeneic Stem Cell Transplantation (SCT) in Patients Who Have Previously Received Pembrolizumab

Evidence for linking the risk to the medicine	Review of pembrolizumab literature regarding increased risk of severe complications of allogeneic stem cell transplantation in patients who have previously received pembrolizumab represents scientific evidence of a possible causal association with pembrolizumab exposure.
Risk factors and risk groups	Patients with hematologic malignancies undergoing allogeneic SCT who were previously treated with a PD-1 inhibitor.
Risk minimisation measures	Routine risk Minimisation measures: <ul style="list-style-type: none"> ▪ For Hematologic malignancies: the increased risk of severe complications of allogeneic SCT in patients who have previously received pembrolizumab is described in the SmPC, Section 4.4, 4.8 and appropriate advice is provided to the prescriber to minimize the risk.
Additional pharmacovigilance activities	Additional pharmacovigilance activities: <ul style="list-style-type: none"> ▪ Safety monitoring in the ongoing HL trial (KN204)

Table II.B.3: Important Potential Risk: Graft Versus Host Disease (GVHD) After Pembrolizumab Administration in Patients With a History of Allogeneic Stem Cell Transplant (SCT)

Evidence for linking the risk to the medicine	Published literature Postmarketing data
Risk factors and risk groups	Patients with a history of allogeneic SCT treated with a PD-1 inhibitor.
Risk minimisation measures	Routine risk Minimisation measures: <ul style="list-style-type: none"> ▪ GVHD after pembrolizumab administration in patients with a history of allogeneic SCT is described in the SmPC, Section 4.4 and appropriate advice is provided to the prescriber to minimize the risk.
Additional pharmacovigilance activities	Routine pharmacovigilance activities Additional pharmacovigilance including: <ul style="list-style-type: none"> ▪ Safety monitoring in all other ongoing MAH-sponsored clinical trials for pembrolizumab in various tumor types

II.C Post-Authorisation Development Plan

II.C.1 Studies Which are Conditions of the Marketing Authorisation

The following studies are conditions of the marketing authorisation:

Table II.C.1.1: Studies Which are Conditions of the Marketing Authorisation

Study Title	Objectives
Efficacy studies which are conditions of the marketing authorisation	
A Phase III, Randomized, Open-label, Clinical Trial to Compare Pembrolizumab with Brentuximab Vedotin in Subjects with Relapsed or Refractory Classical Hodgkin Lymphoma (KN204) (on-going)	To compare overall survival (OS), progression free survival (PFS) and objective response rate (ORR) of pembrolizumab when compared to Brentuximab Vedotin in subjects with relapsed or refractory cHL and to examine the safety and tolerability between treatment groups.
Adjuvant immunotherapy with anti-PD-1 monoclonal antibody Pembrolizumab (MK-3475) versus placebo after complete resection of high-risk Stage III melanoma: A randomized, double-blind Phase 3 trial of the EORTC Melanoma Group (KN054) (on-going)	To prospectively assess whether post-operative adjuvant therapy with pembrolizumab improves recurrence-free survival (RFS) as compared to placebo in high-risk patients with complete resection of Stage IIIA (> 1 mm metastasis), IIIB and IIC melanoma. To prospectively assess whether in the subgroup of patients with PD-L1- positive tumor expression, pembrolizumab improves recurrence-free survival as compared to placebo (primary endpoint); distant metastasis free survival (DMFS) and overall survival (OS) in all-subjects and subjects with PD-L1-positive tumors (secondary endpoints for final study report).
A randomized, placebo-controlled, parallel-group, crossover/rechallenge, multi-center study of adjuvant pembrolizumab in participants 12 years of age and older with resected Stage IIB or IIC cutaneous melanoma (KN716) (on-going)	To compare RFS, DMFS and OS between pembrolizumab and placebo in patients with resected Stage IIB or IIC melanoma.
A randomized, Phase 3 trial with pembrolizumab versus placebo for patients with early stage NSCLC after resection and completion of standard adjuvant therapy (KN091) (on-going)	To compare OS between pembrolizumab and placebo in patients with early stage NSCLC after resection and completion of standard adjuvant therapy To assess data on treatment post-progression, and particularly on the uptake and activity of anti-PD(L)1 in patients previously treated with adjuvant pembrolizumab

II.C.2 Other Studies in Post-Authorisation Development Plan

There are no studies required for pembrolizumab.

REFERENCES

- [Ref. 4.2.3.7.2: TT151105FIN] Anti-Murine PD-1 Antibody (muDX400 L-005571333): Exploratory Multiple-Dose Subcutaneous Immunotoxicity Study in Mice with Hepatitis B Vaccine (L-005552770), 29-Sep-2015
- [Ref. 5.4: 00VMR9] Guleria I, Khosroshahi A, Ansari MJ, Habicht A, Azuma M, Yagita H, et al. A critical role for the programmed death ligand 1 in fetomaternal tolerance. *J Exp Med* 2005;202(2):231-7.
- [Ref. 5.4: 00VMXH] Wafula PO, Teles A, Schumacher A, Pohl K, Yagita H, Volk H-D, et al. PD-1 but not CTLA-4 blockage abrogates the protective effect of regulatory T cells in a pregnancy murine model. *Am J Reprod Immunol* 2009;62:283-92.
- [Ref. 5.4: 02CFKQ] Ansari MJ, Salama AD, Chitnis T, Smith RN, Yagita H, Akiba H, et al. The programmed death-1 (PD-1) pathway regulates autoimmune diabetes in nonobese diabetic (NOD) mice. *J Exp Med* 2003;198(1):63-9.
- [Ref. 5.4: 02CFKR] D'Addio FD, Riella LV, Mfarrej BG, Chabtini L, Adams LT, Yeung M, et al. The link between the PDL1 costimulatory pathway and Th17 in fetomaternal tolerance. *The Journal of Immunology* 2011;187:4530-41.
- [Ref. 5.4: 02CFKT] Habicht A, Dada S, Jurewicz M, Fife BT, Yagita H, Azuma M, et al. A link between PDL1 and T regulatory cells in fetomaternal tolerance. *The Journal of Immunology* 2007;179:5211-9.
- [Ref. 5.4: 02CFKV] Keir ME, Liang SC, Guleria I, Latchman YE, Qipo A, Albacker LA, et al. Tissue expression of PD-L1 mediates peripheral T cell tolerance. *J Exp Med* 2006;203(4):883-95.
- [Ref. 5.4: 02CFKW] Nishimura H, Nose M, Hiai H, Minato N, Honjo T. Development of lupus-like autoimmune diseases by disruption of the PD-1 gene encoding an ITIM motif-carrying immunoreceptor. *Immunity* 1999;11:141-51.

- [Ref. 5.4: 02CFKX] Nishimura H, Okazaki T, Tanaka Y, Nakatani K, Hara M, Matsumori A, et al. Autoimmune dilated cardiomyopathy in PD-1 receptor-deficient mice. *Science* 2001;291:319-22.
- [Ref. 5.4: 03MS03] Forsea AM, del Marmol V, de Vries E, Bailey EE, Geller AC. Melanoma incidence and mortality in Europe: new estimates, persistent disparities. *Brit J Dermatol* 2012;167:1124-30.
- [Ref. 5.4: 03MS0N] De Vries E, Arnold M, Altsitsiadis E, Trakatelli M, Hinrichs B, Stockfleth E, et al. Potential impact of interventions resulting in reduced exposure to ultraviolet (UV) radiation (UVA and UVB) on skin cancer incidence in four European countries, 2010-2050. *Brit J Dermatol* 2012;167(Suppl. 2):53-62.
- [Ref. 5.4: 03N0KS] Grann AF, Frøslev T, Olesen AB, Schmidt H, Lash TL. The impact of comorbidity and stage prognosis of Danish melanoma patients, 1987-2009: a registry-based cohort study. *Br J Cancer* 2013:1-7.
- [Ref. 5.4: 03QY3L] Cheson BD, Pfistner B, Juweid ME, Gascoyne RD, Specht L, Horning SJ, et al. Revised response criteria for malignant lymphoma. *J Clin Oncol* 2007;25:579-86.
- [Ref. 5.4: 03RC2F] Cheson BD, Pfistner B, Juweid ME, Gascoyne RD, Specht L, Horning SJ, et al. Revised response criteria for malignant lymphoma. *J Clin Oncol* 2007;25(6):579-86.
- [Ref. 5.4: 0439J6] Iachina M, Jakobsen E, Moller H, Luchtenborg M, Mellemegaard A, Krasnik M, et al. The Effect of Different Comorbidities on Survival of Non-small Cells Lung Cancer Patients. *Lung*. 2014 Dec 17. [Epub ahead of print].
- [Ref. 5.4: 043SX8] Guastadisegni C, Colafranceschi M, Ottini L, Dogliotti E. Microsatellite instability as a marker of prognosis and response to therapy: a meta-analysis of colorectal cancer survival data. *Eur J Cancer*. 2010 Oct;46(15):2788-98.

- [Ref. 5.4: 043VF3] Venderbosch S, Nagtegaal ID, Maughan TS, Smith CG, Cheadle JP, Fisher D, et al. Mismatch repair status and BRAF mutation status in metastatic colorectal cancer patients: a pooled analysis of the CAIRO, CAIRO2, COIN, and FOCUS studies. *Clin Cancer Res*. 2014 Oct 15;20(20):5322-30.
- [Ref. 5.4: 04BN0K] Znaor A, Lortet-Tieulent J, Laversanne M, Jemal A, Bray F. International variations and trends in renal cell carcinoma incidence and mortality. *Eur Urol*. 2015 Mar;67(3):519-30.
- [Ref. 5.4: 04D0C7] Landis SH, El-Hariry IA, van Herk-Sukel MP, van den Haak P, Janssen-Heijnen ML, Penning-van Beest FJ, et al. Prevalence and incidence of acute and chronic comorbidity in patients with squamous cell carcinoma of the head and neck. *Head Neck*. 2012 Feb;34(2):238-44.
- [Ref. 5.4: 04D76G] Marcos-Gragera R, Mallone S, Kiemeny LA, Vilardell L, Malats N, et al. Urinary tract cancer survival in Europe 1999-2007: Results of the population-based study EURO CARE-5. *Eur J Cancer*. 2015 Oct 5;51(15):2217-30.
- [Ref. 5.4: 04DDMM] Boje CR. Impact of comorbidity on treatment outcome in head and neck squamous cell carcinoma - a systematic review. *Radiother Oncol*. 2014 Jan;110(1):81-90.
- [Ref. 5.4: 04DGDP] Ernani V, Saba NF. Oral cavity cancer: risk factors, pathology, and management. *Oncology*. 2015;89(4):187-95.
- [Ref. 5.4: 04DGDY] Gillison ML, Chaturvedi AK, Anderson WF, Fakhry C. Epidemiology of Human Papillomavirus-Positive Head and Neck Squamous Cell Carcinoma. *J Clin Oncol*. 2015 Oct 10;33(29):3235-42.
- [Ref. 5.4: 04DP5M] Rettig EM, D'Souza G. Epidemiology of head and neck cancer. *Surg Oncol Clin N Am*. 2015 Jul;24(3):379-96.
- [Ref. 5.4: 04FF00] Fuji S, Takano K, Mori T, Eto T, Taniguchi S, Ohashi K, et al. Impact of pretransplant body mass index on the clinical outcome after allogeneic hematopoietic SCT. *Bone Marrow Transplant*. 2014 Dec;49(12):1505-12.

- [Ref. 5.4: 04FF0J] Fuji S, Kanda J, Kato S, Ikegame K, Morishima S, Miyamoto T, et al. Impact of HLA allele mismatch on the clinical outcome in serologically matched related hematopoietic SCT. *Bone Marrow Transplant.* 2014 Sep;49(9):1187-92.
- [Ref. 5.4: 04FF27] Ishida T, Hishizawa M, Kato K, Tanosaki R, Fukuda T, Takatsuka Y, et al. Impact of graft-versus-host disease on allogeneic hematopoietic cell transplantation for adult T cell leukemia-lymphoma focusing on preconditioning regimens: nationwide retrospective study. *Biol Blood Marrow Transplant.* 2013 Dec;19(12):1731-9.
- [Ref. 5.4: 04FF3L] Kanate AS, Mussetti A, Kharfan-Dabaja MA, Ahn KW, DiGilio A, Beitinjaneh A, et al. Reduced-intensity transplantation for lymphomas using haploidentical related donors vs HLA-matched unrelated donors. *Blood.* 2016 Feb 18;127(7):938-47.
- [Ref. 5.4: 04FFX4] Malard F, Chevallier P, Guillaume T, Delaunay J, Rialland F, Harousseau JL, et al. Continuous reduced nonrelapse mortality after allogeneic hematopoietic stem cell transplantation: a single-institution's three decade experience. *Biol Blood Marrow Transplant.* 2014 Aug;20(8):1217-23.
- [Ref. 5.4: 04FG08] Martin PS, Li S, Nikiforow S, Alyea EP 3rd, Antin JH, Armand P, et al. Infused total nucleated cell dose is a better predictor of transplant outcomes than CD34+ cell number in reduced-intensity mobilized peripheral blood allogeneic hematopoietic cell transplantation. *Haematologica.* 2016 Apr;101(4):499-505.
- [Ref. 5.4: 04FG2J] Martino R, Kerguelen A, Valcarcel D, Sureda A, Fachini L, Pinana JL, et al. Reduction of infection-related mortality after allogeneic PBSCT from HLA-identical siblings: longitudinal analysis from 1994 to 2008 at a single institution. *Bone Marrow Transplant.* 2011 May;46(5):690-701.
- [Ref. 5.4: 04FG2L] McClune BL, Ahn KW, Wang HL, Antin JH, Artz AS, Cahn JY, et al. Allogeneic transplantation for patients age ≥ 40 years with non-Hodgkin lymphoma: encouraging progression-free survival. *Biol Blood Marrow Transplant.* 2014 Jul;20(7):960-8.

- [Ref. 5.4: 04FG2Z] Merryman RW, Kim HT, Zinzani PL, Carlo-Stella C, Ansell SM, Halwani AS, et al. 2018 Safety and efficacy of allogeneic Hematopoietic Stem Cell Transplant (HSCT) after treatment with Programmed Cell Death 1 (PD-1) inhibitors [abstract]. Abstracts of the ASH 57th Annual Meeting & Exposition; 2015 Dec 5-8; Orlando, FL.: American Society of Hematology (ASH); 2015. p. 2.
- [Ref. 5.4: 04FG37] Nakane T, Fukuda T, Kanda J, Taniguchi S, Eto T, Ohashi K, et al. Age influences post-graft-versus-host disease non-relapse mortality in adults with acute graft-versus-host disease of varying severity following allogeneic hematopoietic cell transplant. *Leuk Lymphoma*. 2015;56(8):2392-7.
- [Ref. 5.4: 04FGZP] Aguilar-Guisado M, Jimenez-Jambrina M, Espigado I, Rovira M, Martino R, Oriol A, et al. Pneumonia in allogeneic stem cell transplantation recipients: a multicenter prospective study. *Clin Transplant*. 2011 Nov-Dec;25(6):E629-38.
- [Ref. 5.4: 04FGZW] Armenian SH, Sun CL, Vase T, Ness KK, Blum E, Francisco L, et al. Cardiovascular risk factors in hematopoietic cell transplantation survivors: role in development of subsequent cardiovascular disease. *Blood*. 2012 Nov 29;120(23):4505-12.
- [Ref. 5.4: 04FH09] Detrait M, Morisset S, Chalandon Y, Yakoub-Agha I, Dufosse F, Labalette M, et al. Suggestive evidence of a role of HLA-DRB4 mismatches in the outcome of allogeneic hematopoietic stem cell transplantation with HLA-10/10-matched unrelated donors: a French-Swiss retrospective study. *Bone Marrow Transplant*. 2015 Oct;50(10):1316-20.

- [Ref. 5.4: 04FH0G] EI Cheikh J, Sfumato P, Fegueux N, Mohamad M, Yakoub-Agha I, Vigouroux S, et al. Allogenic Hematopoietic Stem Cell transplantation Following Reduced-Intensity Conditioning Regimen for elderly Patients (60 years and older) with Hematological malignancies using Unrelated Donors: A Retrospective Study on Behalf of the French Society for Stem Cell Transplantation and Cell therapies (SFGM-TC). Abstract presented at: 56th ASH (American Society of Hematology) Annual Meeting and Exposition; 2014 Dec 6-9; San Francisco, CA. Abstract no.: 2472.
- [Ref. 5.4: 04FJ33] EI Jurdi N, Kumar A, Pidala J, Bazarbachi A, Djulbegovic B, Kharfan-Dabaja M. Assessment of efficacy (or lack thereof) of donor lymphocyte infusion in relapsed lymphoid malignancies following haematopoietic cell allografting: results of a systematic review [abstract]. Bone Marrow Transplant. 2012;47:S271-S3. Abstract no. P779.
- [Ref. 5.4: 04FJSZ] Ram R, Gooley TA, Maloney DG, Press OW, Pagel JM, Petersdorf SH, et al. Histology and time to progression predict survival for lymphoma recurring after reduced-intensity conditioning and allogeneic hematopoietic cell transplantation. Biol Blood Marrow Transplant. 2011 Oct;17(10):1537-45.
- [Ref. 5.4: 04FJT7] Soiffer RJ, Lerademacher J, Ho V, Kan F, Artz A, Champlin RE, et al. Impact of immune modulation with anti-T-cell antibodies on the outcome of reduced-intensity allogeneic hematopoietic stem cell transplantation for hematologic malignancies. Blood. 2011 Jun 23;117(25):6963-70.
- [Ref. 5.4: 04FJTB] Urbano-Ispizua A, Pavletic SZ, Flowers ME, Klein JP, Zhang MJ, Carreras J, et al. The Impact of Graft-versus-Host Disease on the Relapse Rate in Patients with Lymphoma Depends on the Histological Subtype and the Intensity of the Conditioning Regimen. Biol Blood Marrow Transplant. 2015 Oct;21(10):1746-53.
- [Ref. 5.4: 04FJTD] Wingard JR, Majhail NS, Brazauskas R, Wang Z, Sobocinski KA, Jacobsohn D, et al. Long-term survival and late deaths after allogeneic hematopoietic cell transplantation. J Clin Oncol. 2011 Jun 1;29(16):2230-9.

- [Ref. 5.4: 04FKV7] Carreras E, Diaz-Beya M, Rosinol L, Martinez C, Fernandez-Aviles F, Rovira M. The incidence of veno-occlusive disease following allogeneic hematopoietic stem cell transplantation has diminished and the outcome improved over the last decade. *Biol Blood Marrow Transplant.* 2011 Nov;17(11):1713-20.
- [Ref. 5.4: 04FR9W] Diaz-Padilla I, Romero N, Amir E, Matias-Guiu X, Vilar E, Muggia F, et al. Mismatch repair status and clinical outcome in endometrial cancer: a systematic review and meta-analysis. *Crit Rev Oncol Hematol.* 2013 Oct;88(1):154-67.
- [Ref. 5.4: 04GJ9J] U.S. Prescribing Information: OPDIVO (nivolumab) injection, for intravenous use: 2016.
- [Ref. 5.4: 04GXN0] Ording AG, Nielsen ME, Smith AB, Horvath-Puho E, Sorensen HT. Venous thromboembolism and effect of comorbidity in bladder cancer: A danish nationwide cohort study of 13,809 patients diagnosed between 1995 and 2011. *Urol Oncol.* 2016 Jul;34(7):292.e1-8.
- [Ref. 5.4: 04GXQB] Pakzad R, Mohammadian-Hafshejani A, Mohammadian M, Pakzad I, Safiri S, et al. Incidence and Mortality of Bladder Cancer and their Relationship with Development in Asia. *Asian Pac J Cancer Prev.* 2015;16(16):7365-74.
- [Ref. 5.4: 04H4GM] Goossens-Laan CA, Leliveld AM, Verhoeven RH, Kil PJ, de Bock GH, Hulshof MC, et al. Effects of age and comorbidity on treatment and survival of patients with muscle-invasive bladder cancer. *Int J Cancer.* 2014 Aug 15;135(4):905-12.
- [Ref. 5.4: 04HTK5] Lee JH, Kim Y, Choi JW, Kim YS. Prevalence and prognostic significance of Epstein-Barr virus infection in classical Hodgkin's lymphoma: a meta-analysis. *Arch Med Res.* 2014 Jul;45(5):417-31.
- [Ref. 5.4: 04KB3D] Najita JS, Swetter SM, Geller AC, Gershenwald JE, Zelen M, Lee SJ. Sex Differences in Age at Primary Melanoma Diagnosis in a Population-Based Analysis (US Surveillance, Epidemiology, and End Results, 2005-2011). *J Invest Dermatol.* 2016 Sep;136(9):1894-7.

- [Ref. 5.4: 04KCJW] Lowe KA, Danese MD, Gleeson ML, Langeberg WJ, Ke J, Kelsh MA. Racial and Ethnic Variability in the Prevalence and Incidence of Comorbidities Associated with Gastric Cancer in the United States. *J Gastrointest Cancer*. 2016 Jun;47(2):168-81.
- [Ref. 5.4: 04KJLL] Crocetti E, Mallone S, Robsahm TE, Gavin A, Agius D, Ardanaz E, et al. Survival of patients with skin melanoma in Europe increases further: Results of the EUROCORE-5 study. *Eur J Cancer*. 2015 Oct;51(15):2179-90.
- [Ref. 5.4: 04LJ0D] Robinson BM. Malignant pleural mesothelioma: an epidemiological perspective. *Ann Cardiothorac Surg*. 2012 Nov;1(4):491-6.
- [Ref. 5.4: 04PSFD] Herbaux C, Gauthier J, Brice P, Drumez E, Ysebaert L, Doyen H. et al. Efficacy and tolerability of nivolumab after allogeneic transplantation for relapsed Hodgkin lymphoma. *Blood*. 2017 May 4;129(18):2471-2478.
- [Ref. 5.4: 04Q8R4] Le DT, Durham JN, Smith KN, Wang H, Bartlett BR, Aulakh LK, et al. Mismatch-repair deficiency predicts response of solid tumors to PD-1 blockade. *Science*. 2017 Jun 8. [Epub ahead of print].
- [Ref. 5.4: 04QRV7] De Angelis R, Sant M, Coleman MP, Francisci S, Baili P, Pierannunzio D, et al. Cancer survival in Europe 1999-2007 by country and age: results of EUROCORE-5-a population-based study. *Lancet Oncol*. 2014 Jan;15(1):23-34.
- [Ref. 5.4: 04SJ3G] Fung-Kee-Fung M, Dodge J, Elit L, Lukka H, Chambers A, Oliver T. Follow-up after primary therapy for endometrial cancer: a systematic review. *Gynecol Oncol*. 2006 Jun;101(3):520-9.
- [Ref. 5.4: 04SRJJ] Guizard AV, Uhry Z, de Raucourt D, Mazzoleni G, Sanchez MJ, Ligier K, et al. Trends in net survival from head and neck cancer in six European Latin countries: results from the SUDCAN population-based study. *Eur J Cancer Prev*. 2017 Jan;26:S16-S23.

- [Ref. 5.4: 04SRJK] Le DT, Durham JN, Smith KN, Wang H, Bartlett BR, Aulakh LK, et al. Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. *Science*. 2017 Jul;357(6349):409-13.
- [Ref. 5.4: 04SXYB] Torre LA, Islami F, Siegel RL, Ward EM, Jemal A. Global cancer in women: burden and trends. *Cancer Epidemiol Biomarkers Prev*. 2017 Apr;26(4):444-57.
- [Ref. 5.4: 04TC0P] Vilar E, Gruber SB. Microsatellite instability in colorectal cancer-the stable evidence. *Nat Rev Clin Oncol*. 2010 Mar;7(3):153-62.
- [Ref. 5.4: 04TCGC] Bonneville R, Krook MA, Kautto EA, Miya J, Wing MR, Chen HZ, et al. Landscape of Microsatellite Instability Across 39 Cancer Types. *JCO Precision Oncology* 2017 :1, 1-15.
- [Ref. 5.4: 04TCK9] Hause RJ, Pritchard CC, Shendure J, Salipante SJ. Classification and characterization of microsatellite instability across 18 cancer types. *Nat Med*. 2016 Nov;22(11):1342-1350.
- [Ref. 5.4: 04VW4L] Sarfati D, Koczwara B, Jackson C. The impact of comorbidity on cancer and its treatment. *CA Cancer J Clin*. 2016 Jul-Aug;66(4):337-50.
- [Ref. 5.4: 04VW73] Segev Y, Pal T, Rosen B, McLaughlin JR, Sellers TA, Risch HA, et al. Risk factors for ovarian cancers with and without microsatellite instability. *Int J Gynecol Cancer*. 2014 May;24(4):664-9.
- [Ref. 5.4: 04VW8W] Seo JY, Jin EH, Jo HJ, Yoon H, Shin CM, Park YS, et al. Clinicopathologic and molecular features associated with patient age in gastric cancer. *World J Gastroenterol*. 2015 Jun 14;21(22):6905-13.
- [Ref. 5.4: 04VWCZ] Velho S, Fernandes MS, Leite M, Figueiredo C, Seruca R. Causes and consequences of microsatellite instability in gastric carcinogenesis. *World J Gastroenterol*. 2014 Nov 28;20(44):16433-42.

- [Ref. 5.4: 04VWL2] Tran B, Kopetz S, Tie J, Gibbs P, Jiang ZQ, Lieu CH, et al. Impact of BRAF mutation and microsatellite instability on the pattern of metastatic spread and prognosis in metastatic colorectal cancer. *Cancer*. 2011 Oct 15;117(20):4623-32.
- [Ref. 5.4: 04VWL8] Sinicrope FA, Sargent DJ. Clinical implications of microsatellite instability in sporadic colon cancers. *Curr Opin Oncol*. In press 2009.
- [Ref. 5.4: 04VYPZ] Hong SP, Min BS, Kim TI, Cheon JH, Kim NK, Kim H, et al. The differential impact of microsatellite instability as a marker of prognosis and tumour response between colon cancer and rectal cancer. *Eur J Cancer*. 2012;48:1235-43.
- [Ref. 5.4: 04VZ03] Clay MR, Allison KH, Folkins AK, Longacre TA. Risk of secondary malignancy (including breast) in patients with mismatch-repair protein deficiency. *Am J Surg Pathol*. 2014 Nov;38(11):1494-500.
- [Ref. 5.4: 04VZ0J] Gelsomino F, Barbolini M, Spallanzani A, Pugliese G, Cascinu S. The evolving role of microsatellite instability in colorectal cancer: a review. *Cancer Treat Rev*. 2016;51:19-26.
- [Ref. 5.4: 04W005] Limsui D, Vierkant RA, Tillmans LS, Wang AH, Weisenberger DJ, Laird PW, et al. Cigarette smoking and colorectal cancer risk by molecularly defined subtypes. *J Natl Cancer Inst*. 2010 Jul 21;102(14):1012-22.
- [Ref. 5.4: 04W02X] Phipps AI, Ahnen DJ, Campbell PT, Win AK, Jenkins MA, Lindor NM, et al. Family history of colorectal cancer is not associated with colorectal cancer survival regardless of microsatellite instability status. *Cancer Epidemiol Biomarkers Prev*. 2014 Aug;23(8):1700-4.
- [Ref. 5.4: 04W654] Horii A, Han HJ, Shimada M, Yanagisawa A, Kato Y, Ohta H, et al. Frequent replication errors at microsatellite loci in tumors of patients with multiple primary cancers. *Cancer Res*. 1994 Jul 1;54:3373-5.
- [Ref. 5.4: 04YDD6] Hillemanns P, Soergel P, Hertel H, Jentschke M. Epidemiology and early detection of cervical cancer. *Oncol Res Treat*. 2016;39:501-6.

- [Ref. 5.4: 04YDDS] LaVigne AW, Triedman SA, Randall TC, Trimble EL, Viswanathan AN. Cervical cancer in low and middle income countries: addressing barriers to radiotherapy delivery. *Gynecol Oncol Rep.* 2017;22:16-20.
- [Ref. 5.4: 04YVTB] Ibfelt EH, Kjaer SK, Hogdall C, Steding-Jessen M, Kjaer TK, Osler M, et al. Socioeconomic position and survival after cervical cancer: influence of cancer stage, comorbidity and smoking among Danish women diagnosed between 2005 and 2010. *Br J Cancer.* 2013;109:2489-95.
- [Ref. 5.4: 04YVTL] van der Aa MA, Siesling S, Kruitwagen RFP, Lybeert MLM, Coebergh JW, Janssen-Heijnen ML. Comorbidity and age affect treatment policy for cervical cancer: a population-based study in the south of the Netherlands, 1995-2004. *Eur J Gynaecol Oncol.* 2008;XXIX(5):493-8.
- [Ref. 5.4: 052ZM2] Shanbhag S, Ambinder RF. Hodgkin lymphoma: a review and update on recent progress. *CA Cancer J Clin.* 2018 Mar-Apr;68(2):116-32.
- [Ref. 5.4: 053GXR] Capitanio U, Bensalah K, Bex A, Boorjian SA, Bray F, Coleman J, et al. Epidemiology of renal cell carcinoma. *Eur Urol.* In press 2018.
- [Ref. 5.4: 053GXZ] Petejova N, Martinek A. Renal cell carcinoma: review of etiology, pathophysiology and risk factors. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub.* 2016 Jun;160(2):183-94.
- [Ref. 5.4: 053GY0] Rao A, Wiggins C, Lauer RC. Survival outcomes for advanced kidney cancer patients in the era of targeted therapies. *Ann Transl Med.* 2018;6(9):165.
- [Ref. 5.4: 053GY2] Rossi SH, Klatte T, Usher-Smith J, Stewart GD. Epidemiology and screening for renal cancer. *World J Urol.* 2018;36:1341-53.
- [Ref. 5.4: 053TKZ] Mangone L, Bossard N, Marcos-Gragera R, Pezzarossi A, Roncaglia F, Giorgi Rossi P. Trends in net survival from kidney cancer in six European Latin countries: results from the SUDCAN population-based study. *Eur J Cancer Prev.* 2017;26(suppl 1):S121-7.

- [Ref. 5.4: 054HBS] Domper Arnal MJ, Ferrandez Arenas A, Lanas Arbeloa A. Esophageal cancer: risk factors, screening and endoscopic treatment in Western and Eastern countries. *World J Gastroenterol*. 2015 Jul 14;21(26):7933-43.
- [Ref. 5.4: 054HM8] Launoy G, Bossard N, Castro C, Manfredi S. Trends in net survival from esophageal cancer in six European Latin countries: results from the SUDCAN population-based study. *Eur J Cancer Prev*. 2017;26(1):S24-31.
- [Ref. 5.4: 054HS0] Nandy N, Dasanu CA. Incidence of second primary malignancies in patients with esophageal cancer: a comprehensive review. *Curr Med Res Opin*. 2013 Sep;29(9):1055-65.
- [Ref. 5.4: 057K9G] SEER*Stat [Internet]. Bethesda (MD): National Cancer Institute. 2019. Cancer Stat Facts: Bladder Cancer: 2009-2015; [about 13 screens]. Available from: <http://seer.cancer.gov/statfacts/html/urinb.html>.
- [Ref. 5.4: 057SQL] SEER*Stat [Internet]. Bethesda (MD): National Cancer Institute (NCI). 2019. Cancer stat facts: oral cavity and pharynx cancer; [about 12 screens]. Available from: <https://seer.cancer.gov/statfacts/html/oralcav.html>.
- [Ref. 5.4: 057XM4] Kurnit KC, Ward KK, McHale MT, Saenz CC, Plaxe SC. Increased prevalence of comorbid conditions in women with uterine cancer. *Gynecol Oncol*. 2015;138:731-4.
- [Ref. 5.4: 057XQG] Williams GR, Mackenzie A, Magnuson A, Olin R, Chapman A, Mohile S, et al. Comorbidity in older adults with cancer. *J Geriatr Oncol*. 2016;7:249-57.
- [Ref. 5.4: 058TT3] SEER*Stat [Internet]. Bethesda (MD): National Cancer Institute (NCI). 2019. Cancer stat facts: lung and bronchus cancer; [about 12 screens]. Available from: <https://seer.cancer.gov/statfacts/html/lungb.html>.
- [Ref. 5.4: 059FM9] National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: bladder cancer; version 4. 2019. Plymouth Meeting (PA): National Comprehensive Cancer Network (NCCN); 2019. 108 p.

- [Ref. 5.4: 059PJJ] Liao HY, Zhang WW, Sun JY, Li FY, He ZY, Wu SG. The clinicopathological features and survival outcomes of different histological subtypes in triple-negative breast cancer. *J Cancer*. 2018 Jan 1;9(2):296-303.
- [Ref. 5.4: 059RH3] SEER*Stat [Internet]. Bethesda (MD): National Cancer Institute (NCI). 2019. Cancer stat facts: female breast cancer; [about 12 screens]. Available from: <https://seer.cancer.gov/statfacts/html/breast.html>.
- [Ref. 5.4: 059X3Q] Kumar P and Aggarwal R. An overview of triple-negative breast cancer. *Arch Gynecol Obstet*. 2016;293:247-69.
- [Ref. 5.4: 05B2QG] Cardoso F, Kyriakides S, Ohno S, Penault-Llorca F, Poortmans P, Rubio IT, et al. Early breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2019;30(8):1194-220.
- [Ref. 5.4: 05BFQQ] SEER*Stat [Internet]. Bethesda (MD): National Cancer Institute (NCI). 2019. Cancer stat facts: kidney and renal pelvis cancer; [about 12 screens]. Available from: <https://seer.cancer.gov/statfacts/html/kidrp.html>.
- [Ref. 5.4: 05DLL7] National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: colon cancer; version 3.2019. Plymouth Meeting (PA): National Comprehensive Cancer Network (NCCN); 2019. 182 p.
- [Ref. 5.4: 05DLL8] National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: cutaneous melanoma; version 2.2019. Plymouth Meeting (PA): National Comprehensive Cancer Network (NCCN); 2019. 199 p.
- [Ref. 5.4: 05DLLB] National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: head and neck cancers; version 3.2019. Plymouth Meeting (PA): National Comprehensive Cancer Network (NCCN); 2019. 231 p.

- [Ref. 5.4: 05DMDL] National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: Hodgkin lymphoma; version 2.2019. Plymouth Meeting (PA): National Comprehensive Cancer Network (NCCN); 2019. 86 p.
- [Ref. 5.4: 05DMDM] National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: kidney cancer; version 2.2020. Plymouth Meeting (PA): National Comprehensive Cancer Network (NCCN); 2019. 64 p.
- [Ref. 5.4: 05DMDQ] National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: non-small cell lung cancer; version 7.2019. Plymouth Meeting (PA): National Comprehensive Cancer Network (NCCN); 2019. 231 p.
- [Ref. 5.4: 05DMDS] National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: uterine neoplasms; version 4.2019. Plymouth Meeting (PA): National Comprehensive Cancer Network (NCCN); 2019. 103 p.
- [Ref. 5.4: 05DMHK] SEER*Stat [Internet]. Bethesda (MD): National Cancer Institute (NCI). 2019. Cancer stat facts: esophageal cancer; [about 12 screens]. Available from: <https://seer.cancer.gov/statfacts/html/esoph.html>.
- [Ref. 5.4: 05DMHL] SEER*Stat [Internet]. Bethesda (MD): National Cancer Institute (NCI). 2019. Cancer stat facts: Hodgkin lymphoma; [about 12 screens]. Available from: <https://seer.cancer.gov/statfacts/html/hodg.html>.
- [Ref. 5.4: 05DMHM] SEER*Stat [Internet]. Bethesda (MD): National Cancer Institute (NCI). 2019. Cancer stat facts: laryngeal cancer; [about 12 screens]. Available from: <https://seer.cancer.gov/statfacts/html/laryn.html>.
- [Ref. 5.4: 05DMLD] Miyazaki J, Nishiyama H. Epidemiology of urothelial carcinoma. *Int J Urol*. 2017;24:730-4.
- [Ref. 5.4: 05DMLF] Ratti M, Lampis A, Hahne JC, Passalacqua R, Valeri N. Microsatellite instability in gastric cancer: molecular bases, clinical perspectives, and new treatment approaches. *Cell Mol Life Sci*. 2018;75:4151-62.

- [Ref. 5.4: 05DMN4] U.S. Prescribing Information: OPDIVO (nivolumab) injection, for intravenous use: Sep 2019.
- [Ref. 5.4: 05DMN5] U.S. Prescribing Information: TECENTRIQ (atezolizumab) injection, for intravenous use: May 2019.
- [Ref. 5.4: 05DMN6] U.S. Prescribing Information: YERVOY (ipilimumab) injection, for intravenous use: Sep 2019.
- [Ref. 5.4: 05DN03] SEER*Stat [Internet]. Bethesda (MD): National Cancer Institute (NCI). 2019. Cancer stat facts: melanoma of the skin; [about 12 screens]. Available from: <https://seer.cancer.gov/statfacts/html/melan.html>.
- [Ref. 5.4: 05DNDC] U.S. Prescribing Information: BAVENCIO (avelumab) injection, for intravenous use: May 2019.
- [Ref. 5.4: 05DNNW] Li X, Yang J, Peng L, Sahin AA, Huo L, Ward KC, et al. Triple-negative breast cancer has worse overall survival and cause-specific survival than non-triple-negative breast cancer. *Breast Cancer Res Treat.* 2017;161:279-87.
- [Ref. 5.4: 05DNRG] Hwang SY, Park S, Kwon Y. Recent therapeutic trends and promising targets in triple negative breast cancer. *Pharmacol Ther.* 2019;199:30-57.
- [Ref. 5.4: 05DNTY] Howlader N, Altekruse SF, Li CI, Chen VW, Clarke CA, Ries LAG, et al. US incidence of breast cancer subtypes defined by joint hormone receptor and HER2 status. *J Natl Cancer Inst.* 2014 May 14;106(5):dju055. Supplemental tables; 5 p.
- [Ref. 5.4: 05DP3H] U.S. Prescribing Information: IMFINZI (durvalumab) injection, for intravenous use: Aug 2019.
- [Ref. 5.4: 05DPN4] De Angelis R, Minicozzi P, Sant M, Dal Maso L, Brewster DH, Osca-Gelis G, et al. Survival variations by country and age for lymphoid and myeloid malignancies in Europe 2000-2007: results of EURO CARE-5 population-based study. *Eur J Cancer.* 2015;51:2254-68.
- [Ref. 5.4: 05DPRK] Capitanio U, Bensalah K, Bex A, Boorjian SA, Bray F, Coleman J, et al. Epidemiology of renal cell carcinoma. *Eur Urol.* 2019;75:74-84.

- [Ref. 5.4: 05DPTW] Beebe-Dimmer JL, Fryzek JP, Yee CL, Dalvi TB, Garabrant DH, Schwartz AG, et al. Mesothelioma in the United States: a Surveillance, Epidemiology, and End Results (SEER)-Medicare investigation of treatment patterns and overall survival. *Clin Epidemiol*. 2016;8:743-50.
- [Ref. 5.4: 05DQKR] Bebe FN, Hu S, Brown TL, Tulp OL. Role, extent, and impact of comorbidity on prognosis and survival in advanced metastatic melanoma: a review. *J Clin Aesthet Dermatol*. 2019 Jan;12(1):16-23.
- [Ref. 5.4: 05DQP0] Anderson KN, Schwab RB, Martinez ME. Reproductive risk factors and breast cancer subtypes: a review of the literature. *Breast Cancer Res Treat*. 2014;144:1-10.
- [Ref. 5.4: 05DQP9] Barta JA, Powell CA, Wisnivesky JP. Global epidemiology of lung cancer. *Ann Glob Health*. 2019 Jan 22;85(1):8.
- [Ref. 5.4: 05DR9D] Ward WH, Farma, JM, editors. *Cutaneous Melanoma: Etiology and Therapy*. Brisbane (Australia): Codon Publications; c2017. 191 p.
- [Ref. 5.4: 05DRW2] Polamraju P, Haque W, Cao K, Verma V, Schwartz M, Klimberg VS, et al. Comparison of outcomes between metaplastic and triple-negative breast cancer patients. *Breast*. 2020;49:8-16.
- [Ref. 5.4: 05DRW3] Svedman FC, Pillas D, Taylor A, Kaur M, Linder R, Hansson J. Stage-specific survival and recurrence in patients with cutaneous malignant melanoma in Europe - a systematic review of the literature. *Clin Epidemiol*. 2016;8:109-22.
- [Ref. 5.4: 05DXHB] Ferlay J, Ervik M, Lam F, Colombet M, Mery L, Pineros M, et al. *Global Cancer Observatory 2018 cancer fact sheets: 17 selected cancers*. Lyon (France): International Agency for Research on Cancer (IARC); 2018. 35 p.
- [Ref. 5.4: 05FB0L] Carr PR, Alwers E, Bienert S, Weberpals J, Kloor M, Brenner H, et al. Lifestyle factors and risk of sporadic colorectal cancer by microsatellite instability status: a systematic review and meta-analyses. *Ann Oncol*. 2018;29(4):825-34.

- [Ref. 5.4: 05FVMP] James SL, Abate D, Abate KH, Abay SM, Abbafati C, Abbasi N, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018 Nov 10;392:1789-858. Erratum in: *Lancet*. 2019 Jun 22;393:e44.
- [Ref. 5.4: 05G5VD] PDQ Pediatric Treatment Editorial Board. PDQ Cancer Information Summaries [Internet]. Bethesda (MD): National Cancer Institute (NCI); 2002. Childhood Hodgkin lymphoma treatment (PDQ): health professional version; 2020 Jan 31 [cited 2020 Mar 18]; 63 p. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK65726/>.
- [Ref. 5.4: 05G7NG] Lee CC, Chang HH, Lu MY, Yang YL, Chou SW, Lin DT, et al. The incidence and risk factors of hepatic veno-occlusive disease after hematopoietic stem cell transplantation in Taiwan. *Ann Hematol*. 2019;98:745-52.
- [Ref. 5.4: 05G7NH] Lewis C, Kim HT, Roeker LE, Cutler C, Koreth J, Nikiforow S, et al. Incidence, predictors, and outcomes of veno-occlusive disease/sinusoidal obstruction syndrome after reduced-intensity allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transplant*. 2020;26:529-39.
- [Ref. 5.4: 05G88Y] Ijaz A, Khan AY, Malik SU, Faridi W, Fraz MA, Usman M, et al. Significant risk of graft-versus-host disease with exposure to checkpoint inhibitors before and after allogeneic transplantation. *Biol Blood Marrow Transplant*. 2019;25:94-9.
- [Ref. 5.4: 05H5SD] Torre LA, Siegel RL, Islami F, Bray F, Jemal A. Worldwide burden of and trends in mortality from gallbladder and other biliary tract cancers. *Clin Gastroenterol Hepatol*. 2018 Mar;16(3):427-37.
- [Ref. 5.4: 05MDND] National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: breast cancer; version 3.2020. Plymouth Meeting (PA): National Comprehensive Cancer Network (NCCN); 2020. 231 p.

- [Ref. 5.4: 05P545] Albayrak A, Garrido-Castro AC, Giannakis M, Umeton R, Devi Manam M, Stover EH, et al. Clinical pan-cancer assessment of mismatch repair deficiency using tumor-only, targeted next-generation sequencing. *JCO Precis Oncol.* 2020;4:1084-97.
- [Ref. 5.4: 05PFKM] Khan SA, Tavolari S, Brandi G. Cholangiocarcinoma: epidemiology and risk factors. *Liver Int.* 2019;39(suppl 1):19-31.
- [Ref. 5.4: 05PJ0F] Ahmad-Nielsen SA, Bruun Nielsen MF, Mortensen MB, Detlefsen S. Frequency of mismatch repair deficiency in pancreatic ductal adenocarcinoma. *Pathol Res Pract.* 2020;216:152985.
- [Ref. 5.4: 05PJ9H] Hu ZI, Shia J, Stadler ZK, Varghese AM, Capanu M, Salo-Mullen E, et al. Evaluating mismatch repair deficiency in pancreatic adenocarcinoma: challenges and recommendations. *Clin Cancer Res.* 2018 Mar 15;24(6):1326-36.
- [Ref. 5.4: 05PBJJ] Vanderwalde A, Spetzler D, Xiao N, Gatalica Z, Marshall J. Microsatellite instability status determined by next-generation sequencing and compared with PD-L1 and tumor mutational burden in 11,348 patients. *Cancer Med.* 2018;7(3):746-56. Erratum in: *Cancer Med.* 2018;7:2792.
- [Ref. 5.4: 05PJBW] Xia M, Singhi AD, Dudley B, Brand R, Nikiforova M, Pai RK. Small bowel adenocarcinoma frequently exhibits lynch syndrome-associated mismatch repair protein deficiency but does not harbor sporadic MLH1 deficiency. *Appl Immunohistochem Mol Morphol.* 2017 Jul;25(6):399-406.

[Ref. 5.4: 05Q3MX]

Global Cancer Observatory (GCO): Cancer Today [Internet]. Lyon (France): International Agency for Research on Cancer (IARC); c2020. Estimated number of new cases in 2020, Europe, females, all ages; Estimated number of new cases in 2020, Europe, females, ages 0-64; Estimated number of new cases in 2020, Europe, females, ages 0-69; [cited 2021 Feb 04]. Available from: https://gco.iarc.fr/today/online-analysis-table?v=2018&mode=population&mode_population=countries&population=900&populations=900&key=asr&sex=0&cancer=14_3_1&type=1&statistic=5&prevalence=0&population_group=5&ages_group%5B%5D=0&ages_group%5B%5D=17&nb_items=5&group_cancer=1&include_nmsc=1&include_nmsc_other=1. Estimation of median age at diagnosis of endometrial cancer in 2020.

[Ref. 5.4: 05QDW7]

Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. In press 2021.

[Ref. 5.4: 06C86L]

Global Cancer Observatory (GCO): Cancer Today [Internet]. Lyon (France): International Agency for Research on Cancer (IARC). Estimated number of new cases in 2020, kidney, both sexes, all ages; [cited 2021 Apr 13]. Available from: https://gco.iarc.fr/today/online-analysis-table?v=2020&mode=population&mode_population=continents&population=900&populations=900&key=asr&sex=0&cancer=29&type=0&statistic=5&prevalence=0&population_group=0&ages_group%5B%5D=0&ages_group%5B%5D=17&group_cancer=1&include_nmsc=1&include_nmsc_other=1#.

[Ref. 5.4: 06C8B2]

International Agency for Research on Cancer. Kidney. Lyon (France): International Agency for Research on Cancer (IARC); 2020. 2 p. Available from: <https://gco.iarc.fr/today/data/factsheets/cancers/29-Kidney-fact-sheet.pdf>.

- [Ref. 5.4: 06CL4Y] SEER*Stat [Internet]. Bethesda (MD): National Cancer Institute (NCI). 2021. Cancer stat facts: kidney and renal pelvis cancer; [about 16 screens]. Available from: <https://seer.cancer.gov/statfacts/html/kidrp.html>.
- [Ref. 5.4: 06CLG4] U.S. Prescribing Information: JEMPERLI (dostarlimab-gxly) injection, for intravenous use: Apr 2021.
- [Ref. 5.4: 06CLG5] U.S. Prescribing Information: LIBTAYO (cemiplimab-rwlc) injection, for intravenous use: Feb 2021.
- [Ref. 5.4: 06D898] International Agency for Research on Cancer. Stomach. Lyon (France): International Agency for Research on Cancer (IARC); 2020. 2 p. Available from: <https://gco.iarc.fr/today/fact-sheets-cancers>.
- [Ref. 5.4: 06D8FH] Yang L, Ying X, Liu S, Lyu G, Xu Z, Zhang X, et al. Gastric cancer: epidemiology, risk factors and prevention strategies. *Chin J Cancer Res*. 2020 Dec;32(6):695-704.
- [Ref. 5.4: 07Z653] Song W, Zhu ZG, Wu Q, Lv CG, Wang YG, Chen L, et al. A nomogram to predict overall survival for biliary tract cancer. *Cancer Manag Res*. 2018;10:1535-41.
- [Ref. 5.4: 07Z655] National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: hepatobiliary cancers; version 5.2021. Plymouth Meeting (PA): National Comprehensive Cancer Network (NCCN); 2021. 162 p.
- [Ref. 5.4: 07Z667] Ellington TD, Momin B, Wilson RJ, Henley SJ, Wu M, Ryerson AB. Incidence and mortality of cancers of the biliary tract, gallbladder, and liver by sex, age, race/ethnicity, and stage at diagnosis: United States, 2013 to 2017. *Cancer Epidemiol Biomarkers Prev*. 2021 Sep;30(9):1607-14.
- [Ref. 5.4: 07Z669] Kim BW, Oh CM, Choi HY, Park JW, Cho H, Ki M. Incidence and overall survival of biliary tract cancers in South Korea from 2006 to 2015: using the National Health Information Database. *Gut Liver*. 2019 Jan;13(1):104-13.

- [Ref. 5.4: 07Z66C] Minicozzi P, Cassetti T, Vener C, Sant M. Analysis of incidence, mortality and survival for pancreatic and biliary tract cancers across Europe, with assessment of influence of revised European age standardisation on estimates. *Cancer Epidemiol.* 2018;55:52-60.
- [Ref. 5.4: 07Z66G] Ouyang G, Liu Q, Wu Y, Liu Z, Lu W, Li S, et al. The global, regional, and national burden of gallbladder and biliary tract cancer and its attributable risk factors in 195 countries and territories, 1990 to 2017: a systematic analysis for the Global Burden of Disease Study 2017. *Cancer.* 2021 Jul 1;127:2238-50.
- [Ref. 5.4: 07Z66H] Rawla P, Sunkara T, Thandra KC, Barsouk A. Epidemiology of gallbladder cancer. *Clin Exp Hepatol.* 2019 May 23;5(2):93-102.
- [Ref. 5.4: 08097L] Tuo JY, Bi JH, Yuan HY, Jiang YF, Ji XW, Li HL, et al. Trends of stomach cancer survival: a systematic review of survival rates from population-based cancer registration. *J Dig Dis.* 2022;23(1):22-32.
- [Ref. 5.4: 08097V] Zhang T, Chen H, Zhang Y, Yin X, Man J, Yang X, et al. Global changing trends in incidence and mortality of gastric cancer by age and sex, 1990-2019: findings from Global Burden of Disease Study. *J Cancer.* 2021;12(22):6695-705.
- [Ref. 5.4: 08389Q] Surveillance, Epidemiology, and End Results Program [Internet]. Bethesda (MD): National Cancer Institute (NCI). Cancer stat facts: stomach cancer; [cited 2022 Jul 6]; [about 18 screens]. Available from: <https://seer.cancer.gov/statfacts/html/stomach.html>.
- [Ref. 5.4: 085XMZ] Gheorghe AS, Dumitrescu EA, Komporaly IA, Mihaila RI, Lungulescu CV, Stanculeanu DL. New targeted therapies and combinations of treatments for cervical, endometrial, and ovarian cancers: a year in review. *Curr Oncol.* 2022 Apr 17;29:2835-47.
- [Ref. 5.4: 085XN3] International Agency for Research on Cancer. Cervix uteri. Lyon (France): International Agency for Research on Cancer (IARC); 2020. 2 p. Available from: <https://gco.iarc.fr/today/fact-sheets-cancers>.

- [Ref. 5.4: 085XN9] Pimple S, Mishra G. Cancer cervix: epidemiology and disease burden. *Cytojournal*. 2022;19:21.
- [Ref. 5.4: 085XNF] Surveillance, Epidemiology, and End Results Program [Internet]. Bethesda (MD): National Cancer Institute (NCI). Cancer stat facts: cervical cancer; [cited 2022 Nov 7]; [about 19 screens]. Available from: <https://seer.cancer.gov/statfacts/html/cervix.html>.
- [Ref. 5.4: 086YC8] Global Cancer Observatory (GCO): Cancer Today [Internet]. Lyon (France): International Agency for Research on Cancer (IARC); c2020. Estimated number of new cases in 2020, WHO Europe (EURO), both sexes, ages 0-54; [cited 2023 Jan 3]. Available from: <https://gco.iarc.fr/today/home>.
- [Ref. 5.4: 086YC9] Global Cancer Observatory (GCO): Cancer Today [Internet]. Lyon (France): International Agency for Research on Cancer (IARC); c2020. Estimated number of new cases in 2020, WHO Europe (EURO), both sexes, ages 55+; [cited 2023 Jan 3]. Available from: <https://gco.iarc.fr/today/home>.
- [Ref. 5.4: 086YCC] Vaccarella S, Franceschi S, Zaridze D, Poljak M, Veerus P, Plummer M, et al. Preventable fractions of cervical cancer via effective screening in six Baltic, central, and eastern European countries 2017-40: a population-based study. *Lancet Oncol*. 2016 Oct;17:1445-52.
- [Ref. 5.4: 0874ZG] National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: Uterine Neoplasms; version 1.2023. Plymouth Meeting (PA): National Comprehensive Cancer Network (NCCN); 2022. 133 p.
- [Ref. 5.4: 087X38] Surveillance, Epidemiology, and End Results Program [Internet]. Bethesda (MD): National Cancer Institute (NCI). Cancer stat facts: uterine cancer; [cited 2023 Feb 16]; [about 4 screens]. Available from: <https://seer.cancer.gov/statfacts/html/corp.html>.

- [Ref. 5.4: 0880L0] Orbach D, Andre N, Brecht IB, Lopez Almaraz R, Ben-Ami T, Vermersch S, et al. Mesothelioma in children and adolescents: the European Cooperative Study Group for Pediatric Rare Tumors (EXPeRT) contribution. *Eur J Cancer*. 2020;140:63-70.
- [Ref. 5.4: 0885WD] Mirallas O, Lopez-Valbuena D, Garcia-Illescas D, Fabregat-Franco C, Verdaguer H, Tabernero J, et al. Advances in the systemic treatment of therapeutic approaches in biliary tract cancer. *ESMO Open*. 2022;7(3):100503.
- [Ref. 5.4: 0885WH] Rizzo A, Federico AD, Ricci AD, Frega G, Palloni A, Pagani R, et al. Targeting BRAF-Mutant biliary tract cancer: Recent advances and future challenges. *Cancer Control*. 2020;27:1-8.
- [Ref. 5.4: 0885WN] Tariq NU, McNamara MG, Valle JW. Biliary tract cancers: current knowledge, clinical candidates and future challenges. *Cancer Manag Res*. 2019;11:2623-42.
- [Ref. 5.4: 0885WP] Wardell CP, Fujita M, Yamada T, Simbolo M, Fassan M, Karlic R, et al. Genomic characterization of biliary tract cancers identifies driver genes and predisposing mutations. *J Hepatol*. 2018;68:959-69.
- [Ref. 5.4: 088R34] Alpert N, van Gerwen M, Taioli E. Epidemiology of mesothelioma in the 21st century in Europe and the United States, 40 years after restricted/banned asbestos use. *Transl Lung Cancer Res*. 2020 Feb;9(Suppl 1):S28-38.
- [Ref. 5.4: 088R4G] Brims F. Epidemiology and clinical aspects of malignant pleural mesothelioma. *Cancers*. 2021 Aug 20;13:4194.
- [Ref. 5.4: 088R5H] Danese MD, Daumont M, Nwokeji E, Gleeson M, Penrod JR, Lubeck D. Treatment patterns and outcomes in older patients with advanced malignant pleural mesothelioma: analyses of surveillance, epidemiology, and end results-medicare data. *Cancer Rep*. 2022;5(9):e1568.

- [Ref. 5.4: 088R5X] Kuryk L, Rodella G, Staniszezewska M, Pancer KW, Wieczorek M, Salmaso S, et al. Novel insights into mesothelioma therapy: emerging avenues and future prospects. *Front Oncol.* 2022 Jun 17;12:916839.
- [Ref. 5.4: 088R69] Lettieri S, Bortolotto C, Agustoni F, Lococo F, Lancia A, Comoli P, et al. The evolving landscape of the molecular epidemiology of malignant pleural mesothelioma. *J Clin Med.* 2021 Mar 3;10:1034.
- [Ref. 5.4: 088RW8] Schumann SO, Kocher G, Minervini F. Epidemiology, diagnosis and treatment of the malignant pleural mesothelioma, a narrative review of literature. *J Thorac Dis.* 2021 Apr;13(4):2510-23.
- [Ref. 5.4: 088RWP] Zhai Z, Ruan J, Zheng Y, Xiang D, Li N, Hu J, et al. Assessment of global trends in the diagnosis of mesothelioma from 1990 to 2017. *JAMA Netw Open.* 2021 Aug 11;4(8):e2120360.
- [Ref. 5.4: 088TMY] Centers for Disease Control and Prevention. Incidence of Malignant mesothelioma, 1999-2018. No. 27. Atlanta, GA: Department of Health and Human Services (HHS); 2022 Mar. 2 p.
- [Ref. 5.4: 088TP2] National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: Mesothelioma: Pleural; Version 1.2023. Plymouth Meeting (PA): National Comprehensive Cancer Network (NCCN); 2023. 60 p.
- [Ref. 5.4: 088TRN] SEER*Explorer [Internet]. Bethesda (MD): National Cancer Institute (NCI). Mesothelioma: SEER incidence rates by age at diagnosis, 2015-2019: observed SEER incidence rate by race/ethnicity, both sexes; [cited 2023 Mar 21]. Available from: https://seer.cancer.gov/statistics-network/explorer/application.html?site=111&data_type=1&graph_type=3&compareBy=sex&chk_sex_1=1&hdn_rate_type=1&race=1&advopt_precision=1&advopt_show_ci=on&advopt_show_count=on&hdn_view=0#resultsRegion0.

- [Ref. 5.4: 08BPP2] National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: cervical cancer; version 1.2023. Plymouth Meeting (PA): National Comprehensive Cancer Network (NCCN); 2023. 97 p.
- [Ref. 5.4: 08CJK3] Surveillance, Epidemiology, and End Results Program [Internet]. Bethesda (MD): National Cancer Institute (NCI). Cancer stat facts: cervical cancer; [cited 2023 Jul 18]; [about 18 screens]. Available from: <https://seer.cancer.gov/statfacts/html/cervix.html>.
- [Ref. 5.4: 08CRCF] Surveillance, Epidemiology, and End Results Program [Internet]. Bethesda (MD): National Cancer Institute (NCI). Cancer stat facts: bladder cancer; [cited 2023 Jul 30]; [about 18 screens]. Available from: <https://seer.cancer.gov/statfacts/html/urinb.html>.
- [Ref. 5.4: 08DSV2] Obermannova R, Alsina M, Cervantes A, Leong T, Lordick F, Nilsson M, et al. Oesophageal cancer: ESMO Clinical Practice Guideline for diagnosis, treatment and follow-up. *Ann Oncol*. 2022;33(10):992-1004.
- [Ref. 5.4: 08G0H3] Di Donato V, D'Oria O, Giannini A, Bogani G, Fischetti M, Santangelo G, et al. Age-adjusted Charlson comorbidity index predicts survival in endometrial cancer patients. *Gynecol Obstet Invest*. 2022;87:191-9.
- [Ref. 5.4: 08G0HC] Dork T, Hillemanns P, Tempfer C, Breu J, Fleisch MC. Genetic susceptibility to endometrial cancer: risk factors and clinical management. *Cancers*. 2020 Aug 25;12:2407.
- [Ref. 5.4: 08G0HL] Global Cancer Observatory (GCO). Corpus uteri. Lyon (France): International Agency for Research on Cancer (IARC); 2020 Dec. 2 p.
- [Ref. 5.4: 08G0HQ] Halla K. Emerging treatment options for advanced or recurrent endometrial cancer. *J Adv Pract Oncol*. 2022 Jan-Feb;13(1):45-59.
- [Ref. 5.4: 08G0J9] Nagle CM, Crosbie EJ, Brand A, Obermair A, Oehler MK, Quinn M, et al. The association between diabetes, comorbidities, body mass index and all-cause and cause-specific mortality among women with endometrial cancer. *Gynecol Oncol*. 2018;150:99-105.

- [Ref. 5.4: 08G0JV] Raglan O, Kalliala I, Markozannes G, Cividini S, Gunter MJ, Nautiyal J, et al. Risk factors for endometrial cancer: an umbrella review of the literature. *Int J Cancer*. 2019;145:1719-30.
- [Ref. 5.4: 08HXX0] International Agency for Research on Cancer. Incidence rate of mesothelioma worldwide in 2022. Lyon (France): International Agency for Research on Cancer (IARC); c1965-2024; [cited 2024 Feb 8]; [about 2 screens]. Available from: https://gco.iarc.fr/today/en/dataviz/tables?mode=cancer&cancers=18&multiple_populations=1&age_end=17&age_start=4&populations=900&group_populations=1&cancers_h=18.
- [Ref. 5.4: 08HXX6] International Agency for Research on Cancer. Incidence rate of mesothelioma in EU-27 in 2022. Lyon (France): International Agency for Research on Cancer (IARC); c1965-2024; [cited 2024 Feb 8]; [about 2 screens]. Available from: https://gco.iarc.fr/today/en/dataviz/tables?mode=cancer&cancers=18&multiple_populations=1&age_end=17&age_start=4&populations=940&group_populations=1&cancers_h=18.
- [Ref. 5.4: 08HXXC] International Agency for Research on Cancer. Incidence rate of mesothelioma in UK. Lyon (France): International Agency for Research on Cancer (IARC); c1965-2024; [cited 2024 Feb 8]; [about 2 screens]. Available from: https://gco.iarc.fr/today/en/dataviz/tables?mode=cancer&cancers=18&multiple_populations=1&age_end=17&age_start=4&populations=826&group_populations=1&cancers_h=18.
- [Ref. 5.4: 08HZXM] Siegel RL, Giaquinto AN, Jemal A. Cancer statistics, 2024. *CA Cancer J Clin*. 2024;74:12-49.
- [Ref. 5.4: 08J9LT] Malekzadeh P, Good M, Hughes MS. Cytoreductive surgery and hyperthermic intraperitoneal chemotherapy (HIPEC) with cisplatin in pediatric patients with peritoneal mesothelioma: a single institution experience and long term follow up. *Int J Hyperthermia*. 2021;38(1):326-31.

- [Ref. 5.4: 08J9N7] Scharf JB, Lees GM, Sergi CM. Malignant pleural mesothelioma in a child. *J Ped Surg Case Reports*. 2015;3:440-3.
- [Ref. 5.4: 08JD46] Zhang Y, Vaccarella S, Morgan E, Li M, Etxeberria J, Chokunonga E, et al. Global variations in lung cancer incidence by histological subtype in 2020: a population-based study. *Lancet Oncol*. 2023 Nov;24:1206-18.
- [Ref. 5.4: 08JJK5] Global Cancer Observatory (GCO): Cancer Today [Internet]. Lyon (France): International Agency for Research on Cancer (IARC); c1965-2024. Oesophagus; [cited 2024 Mar 4]; 2 p. Available from: <https://gco.iarc.who.int/media/globocan/factsheets/cancers/6-oesophagus-fact-sheet.pdf>.
- [Ref. 5.4: 08JJK6] Global Cancer Observatory (GCO): Cancer Today [Internet]. Lyon (France): International Agency for Research on Cancer (IARC); c1965-2024. Stomach; [cited 2024 Mar 4]; 2 p. Available from: <https://gco.iarc.who.int/media/globocan/factsheets/cancers/7-stomach-fact-sheet.pdf>.
- [Ref. 5.4: 08KSQF] Surveillance, Epidemiology, and End Results Program [Internet]. Bethesda (MD): National Cancer Institute (NCI). Cancer stat facts: stomach cancer; [cited 2024 Apr 26]; [about 17 screens]. Available from: <https://seer.cancer.gov/statfacts/html/stomach.html>.
- [Ref. 5.4: 08KZXV] Surveillance, Epidemiology, and End Results Program [Internet]. Bethesda (MD): National Cancer Institute (NCI). Cancer stat facts: esophageal cancer; [cited 2024 May 8]; [about 17 screens]. Available from: <https://seer.cancer.gov/statfacts/html/esoph.html>.
- [Ref. 5.4: 08LCQN] Bray F, Laversanne M, Sung H, Ferlay J, Siegel RL, Soerjomataram I, et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2024;74:229-63.
- [Ref. 5.4: 08LJMC] National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: gastric cancer; version 2.2024. Plymouth Meeting (PA): National Comprehensive Cancer Network (NCCN); 2024. 149 p.

- [Ref. 5.4: 08PKM9] Global Cancer Observatory (GCO): Cancer Today [Internet]. Lyon (France): International Agency for Research on Cancer (IARC); c1965-2024. Head and neck cancer, mortality, both sexes, in 2022; [cited 2024 Oct 16]. Available from: <https://gco.iarc.fr/today>.
- [Ref. 5.4: 08PQSB] National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: head and neck cancers; version 4.2024. Plymouth Meeting (PA): National Comprehensive Cancer Network (NCCN); 2024. 255 p.
- [Ref. 5.4: 08PQSD] Global Cancer Observatory (GCO): Cancer Today [Internet]. Lyon (France): International Agency for Research on Cancer (IARC); c1965-2024. Incidence, both sexes, in 2022 - Europe; [cited 2024 Oct 22]. Available from: https://gco.iarc.who.int/today/en/dataviz/tables-prevalence?mode=population&cancers=1_3_5_14&group_cancers=1&populations=100_112_191_196_203_208_233_246_250_276_300_348_352_372_380_40_428_440_442_470_498_499_528_56_578_616_620_642_643_68_8_70_703_705_724_752_756_8_804_807_826&types=2&multiple_cancers=1&multiple_populations=1.
- [Ref. 5.4: 08QGQ2] Global Cancer Observatory (GCO): Cancer Today [Internet]. Lyon (France): International Agency for Research on Cancer (IARC); c1965-2024. Lung Cancer - incidence, both sexes, UN regions; [cited 2024 Nov 21]; 4 p. Available from: <https://gco.iarc.fr/today/en>.
- [Ref. 5.4: 08QPK8] Matz M, Valkov M, Sekerija M, Luttmann S, Caldarella A, Coleman MP, et al. Worldwide trends in esophageal cancer survival, by sub-site, morphology, and sex: an analysis of 696,974 adults diagnosed in 60 countries during 2000-2014 (CONCORD-3). *Cancer Commun.* 2023;43:963-80.

ANNEXES

ANNEX 4 – SPECIFIC ADVERSE DRUG REACTION FOLLOW-UP FORMS

This RMP annex is not applicable.

ANNEX 6 – DETAILS OF PROPOSED ADDITIONAL RISK MINIMISATION ACTIVITIES

Risk Minimisation Activity: Educational material

The Marketing Authorisation Holder (MAH) shall ensure that all physicians who are expected to prescribe pembrolizumab are provided with the *Patient Cards*.

Overview of Implementation

- The MAH will ensure distribution of the educational material; details on the exact target audience for the communications/materials and the specifics of the risk minimisation materials will be agreed with the National Competent Authorities.
- In addition, the MAH will train the local MSD country staff on the patient card and on the need to provide the material to new prescribers. Records of the training will be maintained. The records of the mailer will be tracked and be used to report on the implementation of the risk minimization activity.
- Information from ongoing pharmacovigilance activities (routine pharmacovigilance) will assess appropriate recognition and management of immune-mediated adverse reactions and that the events observed in real world use are consistent with the cases observed in the clinical trial program and characterized in the SmPC and patient educational material.

Patient Card

Key elements:

- Information that pembrolizumab can cause serious side effects in many parts of the body that can get worse, can sometimes become life-threatening and lead to death, and need to be addressed immediately by the patient communicating with the treating physician
- Description of the main symptoms of immune-mediated adverse reactions and the importance of notifying their treating physician immediately if symptoms occur, persist, or worsen:
 - Lungs: shortness of breath, chest pain, or coughing
 - Intestines: diarrhoea or more bowel movements than usual, black, tarry, sticky stools or stools with blood or mucus, or severe stomach pain or tenderness
 - Liver: nausea or vomiting, pain on the right side of stomach, yellowing of skin or whites of eyes, dark urine, or bleeding or bruising more easily than normal
 - Kidneys: changes in the amount or colour of your urine

- Hormone glands (especially the thyroid, pituitary, and adrenal glands): rapid heartbeat, weight loss, increased sweating, weight gain, hair loss, feeling cold, constipation, deeper voice, muscle aches, dizziness or fainting, headaches that will not go away
- Type 1 diabetes, including diabetic ketoacidosis (acid in the blood produced from diabetes): feeling more hungry or thirsty, needing to urinate more often, weight loss, feeling tired or feeling sick, stomach pain, fast and deep breathing, confusion, unusual sleepiness, a sweet smell to your breath, a sweet or metallic taste in your mouth, or a different odour to your urine or sweat
- The importance of not attempting to self-treat any symptoms without consulting their Healthcare Professional first
- Placeholder including the weblink of the Package Leaflet on the EMA website
- The importance of carrying the wallet-sized Patient Card at all times, to show it at all medical visits to healthcare professionals other than the prescriber (e.g. emergency healthcare professionals). The Card reminds patients about key symptoms that need to be reported immediately to the physician/ nurse. It also contains prompts to enter contact details of the treating physician and to alert other physicians that the patient is treated with pembrolizumab.

The MAH shall agree to the format and content of the above material with the National Competent Authority prior to launch in the Member State.