



1 24 June 2022
2 EMA/CVMP/VICH/839/1999
3 Committee for Veterinary Medicinal Products (CVMP)

4 **VICH GL12 Efficacy of anthelmintics: specific**
5 **recommendations for bovines (Revision 1)**
6 **Draft**

Draft agreed by VICH Steering Committee	May 2022
Adoption by CVMP for release for consultation	15 June 2022
Start of public consultation	24 June 2022
End of consultation (deadline for comments)	1 November 2022

7
8

Comments should be provided using this [template](#). The completed comments form should be sent to vet-guidelines@ema.europa.eu

9





International Cooperation on Harmonisation of Technical Requirements
for Registration of Veterinary Medicinal Products

VICH GL12 (ANTHELMINTICS BOVINES)
May 2022
Revision at Step 9
For consultation at Step 4

EFFICACY OF ANTHELMINTICS: SPECIFIC RECOMMENDATIONS FOR BOVINES (REVISION 1)

Revision at Step 9

Recommended for Consultation at Step 4 of the VICH Process
in May 2022
by the VICH Steering Committee

This Guideline has been developed by the appropriate VICH Expert Working Group will be subject to consultation by the parties, in accordance with the VICH Process. At Step 7 of the Process the final draft is recommended for adoption to the regulatory bodies of the European Union, Japan and USA.

Secretariat: c/o HealthforAnimals, 168 Av de Tervueren, B-1150 Brussels (Belgium) - Tel. +32 2 543 75 72
e-mail : sec@vichsec.org - Website : <http://www.vichsec.org>

54 **EFFICACY OF ANTHELMINTICS:**
55 **SPECIFIC RECOMMENDATIONS FOR BOVINES**

56
57
58
59 **INTRODUCTION**
60

61 These guidelines for bovines were developed by the Working Group established by the
62 Veterinary International Cooperation on Harmonization (VICH), Anthelmintic Guidelines and
63 were subsequently revised in 2022. They should be read in conjunction with the VICH Efficacy
64 of Anthelmintics: General Requirements (VICH GL7) which should be referred to for
65 discussion of broad aspects for providing pivotal data to demonstrate product anthelmintic
66 effectiveness. The present document is structured similarly to VICH GL7 with the aim of
67 simplicity for readers comparing both documents.
68

69 The aim of the guidelines for bovines is (1) to be more specific for certain specific issues
70 for bovines not discussed in VICH GL7; (2) to highlight differences with VICH GL7 on
71 efficacy data requirements and (3) to give explanations for disparities with VICH GL7.
72

73 It is also important to note that technical procedures to be followed in the studies are not the
74 aim of this guideline. We recommend to the sponsors to refer to the pertinent procedures
75 described in detail in other published documents e.g. WAAVP Second Edition of Guidelines
76 for Evaluating the Efficacy of Anthelmintics in Ruminants (Bovine, Ovine, Caprine) Veterinary
77 Parasitology **58**: 181-213, 1995, and updated versions as they are published.
78

79 **A. General Elements**
80
81

82 **1 - The Evaluation of Effectiveness Data**
83

84 Only controlled tests based on parasite counts of adults/larvae are acceptable both for the
85 dose determination and dose confirmation studies, since critical tests generally are not
86 considered to be reliable for ruminants. Egg counts/larval identification is the preferred
87 method to evaluate the effectiveness in field studies. Long-acting or sustained-release
88 products should be subject to the same evaluation procedures as other therapeutic
89 anthelmintics. Adequate parasite infection should be defined in the protocol according to
90 regional prevalence or historical data and/or statistical analysis.
91

92 **2 - Use of Natural or Induced Infections**
93

94 Dose determination studies generally should be conducted using induced infections with
95 either laboratory strains or recent field isolates. Limited experience exists with induced
96 infections of *Toxocara vitulorum*, cestodes and *Dicrocoelium dendriticum*. For these parasites
97 the use of natural infections instead of induced infections may be justified.
98

99 Dose confirmation studies should be conducted using naturally infected animals, however,
100 induced infections or superimposed induced infections can also be used. This procedure will
101 allow a wide range of parasites to be present. For claims against 4th stage larvae, induced
102 infections must be used. For claims against hypobiotic larvae, only natural infections can
103 be considered. Sponsors should aim for a maximum period of accumulation of hypobiotic
104 larvae for the particular parasite species being targeted in trial animals. This will be area
105 or regionally dependent. Specific details on area or regional situations should be obtained
106 from experts on a case by case basis. In all cases, animals need to be housed (to preclude
107 reinfection) for a minimum of 2 weeks before treatment.

108 Persistent efficacy studies should be conducted using induced infections with recent field
109 isolates.

110
 111 The history of the parasites used in the induced infection studies should be included in the
 112 final report.
 113

114 **3 - Number of Infective Parasitic Forms Recommended for Induced Infections**

115
 116 The number to be used is approximate and will depend on the isolate that is used. The final
 117 number of larvae used in the infection should be included in the final report. Table 1 shows
 118 the range of numbers recommended for parasite species with existing infection models.
 119

120 **Table 1 - Number of Infective Stages Used to Produce Adequate Infections in Cattle for**
 121 **Anthelmintic Evaluation.**
 122

Parasite Anatomical Location Genus Species	Range of eggs/larvae
Abomasum	
<i>Haemonchus placei</i>	5,000 - 10,000
<i>Ostertagia ostertagi</i>	10,000 - 30,000
<i>Trichostrongylus axei</i>	10,000 - 30,000
Intestines	
<i>Cooperia oncophora</i>	10,000 - 30,000
<i>C. punctata</i>	10,000 - 15,000
<i>T. colubriformis</i>	10,000 - 30,000
<i>Nematodirus spathiger</i>	3,000 - 10,000
<i>N. helvetianus</i>	3,000 - 10,000
<i>N. battus</i>	3,000 - 6,000
<i>Oesophagostomum radiatum</i>	1,000 - 2,500
<i>O. venulosum</i>	1,000 - 2,000
<i>Chabertia ovina</i>	500 - 1,500
<i>Bunostomum phlebotomum</i>	500 - 1,500
<i>Strongyloides papillosus</i>	1,000 - 200,000
<i>Trichuris</i> spp.	1,000
Lungs	
<i>Dictyocaulus viviparus</i>	500 - 6,000
Liver	
<i>Fasciola hepatica</i> (metacercaria)	
Adult cattle	1,000
Young cattle	500-1,000

123
 124 **4 - Recommendations for the Calculation of Effectiveness**
 125

126
 127 **4.1 Criteria to Grant a Claim**

128
 129 To be granted a claim the following pivotal data should be included:
 130

- 131 a) Two dose confirmation studies conducted with a minimum of 6 adequately infected
 132 animals in the non-medicated control group in each study. The infection of the
 133 animals in the study will be deemed adequate based on historical, parasitological
 134 and/or statistical criteria.
 135
- 136 b) The differences in parasite counts between treated and control animals should be
 137 statistically significant ($p \leq 0.05$).
 138
- 139 c) Percent efficacy should be 90% or higher and calculated and interpreted using the
 140 procedures described in Section 4.2 of VICH GL7.
 141

4.2 Number of Animals (Dose Determination, Dose Confirmation and Persistency Trials)

The minimum number of animals required per experimental group is a critical point. Although the number of animals will depend on the possibility to process the data statistically according to the adequate statistical analysis, it has been recommended, to achieve harmonization, that the inclusion of at least 6 animals in each experimental group is a minimum.

In cases where there are several studies none of which have 6 adequately infected animals in the control group (for example, important rare parasites), the results obtained could be pooled to accumulate 12 animals in the studies; and statistical significance calculated. If the difference is significant ($p < 0.05$), effectiveness may be calculated and if the infection is deemed adequate, the claim may be granted. Sampling techniques and estimation of worm burden should be similar among laboratories involved in the studies to allow adequate and meaningful extrapolation of the results to the population.

4.3 Adequacy of Infection

The minimum adequate number of helminths in individual control animals should be defined in the protocol. However, final conclusions regarding adequacy of infection will be made as part of the final report based on statistical analysis, historical data, literature review, or expert testimony. The range of bovine helminths (adults) that has been considered adequate to grant a claim will vary according to the species. Generally, a minimum of 100 nematodes in individual control animals is considered an adequate infection. Lower counts are to be expected with *Bunostomum* spp, *Oesophagostomum* spp., *Trichuris* spp., and *Dictyocaulus* spp. For *Fasciola* spp. minimum counts of 20 adults are considered adequate.

Recommended worm counts (in individual control animals) to be considered adequate for specific parasites include:

Cooperia oncophora and *C. punctata*: 200 worms
All other Cooperia species: 100 worms
Haemonchus placei: 200 worms
Haemonchus contortus: 200 worms
Ostertagia ostertagi: 200 worms
Nematodirus helvetianus: 100 worms
Trichostrongylus axei, *T. colubriformis*, *T. longispicularis*: 100 worms
Bunostomum phlebotomum: 50 worms
Oesophagostomum radiatum: 50 worms
Dictyocaulus viviparus: 10 worms

4.4 Label Claims

For adult claims as a general rule, the treatment should not be administered earlier than 21 to 25 days after infection; optimum for most species is 28 to 32 days. Major exceptions are *Oesophagostomum* spp. (34 to 49 days), *Bunostomum* spp. (52 to 56 days), *Strongyloides papillosus* (14 to 16 days) and *Fasciola* spp. (8 to 12 weeks).

For L4 claims, treatments should be given on the following days after infection: 3 to 4 days for *Strongyloides papillosus*, 5 to 6 days for *Haemonchus* spp., *Trichostrongylus* spp. and *Cooperia* spp., 7 days for *Ostertagia* spp. and *Dictyocaulus viviparus*, 8 to 10 days for *Nematodirus* spp. and 15 to 17 days for *Oesophagostomum* spp. The term 'immature' on the labeling is not acceptable for these claims.

For early immature *Fasciola* spp., treatments should be given 1 to 5 weeks after infection and for late immatures at 6 to 9 weeks.

5 - Treatment Procedures

199
200 The method of administration (oral, parenteral, topical, slow-release etc.), formulation and
201 extent of activity of a product will influence the protocol design. It is advisable to consider the
202 weather and animal relationship with regard to effectiveness of topical formulations. Slow-
203 release products should be tested over the entire proposed effective time unless additional
204 information suggests that this is unnecessary, e.g. blood levels demonstrate steady state at
205 all points of the proposed therapeutic period.

206
207 When the drug is to be administered in the water or in a feed, it should be done as much as
208 possible following the labelling recommendations. Palatability studies may be required for
209 medicated feed. Samples of medicated water or medicated feed should be collected to confirm
210 drug concentration. The amount of medicated product provided to each animal should be
211 recorded to ensure that the treatment satisfies the label recommendations. For products used
212 topically, the impact of weather (e.g. rainfall, UV light), and coat length should be included in the
213 evaluation of the effectiveness of the product.

214 215 **6 - Animal Selection, Allocation and Handling**

216
217 Test animals should be clinically healthy and representative of the age, sex, and class for
218 which the claim of the test anthelmintic is to be made. In general, the animals should be
219 ruminating, and older than 3 months of age. Randomization to treatment group should be
220 performed using an adequate method that should be described in the protocol and final report.
221 Blocking should only be employed if it is expected to reduce residual error in the study. If
222 blocking is used, blocks should be included as a random effect in the statistical model.
223 Nevertheless, blocking is not always the most appropriate method for reducing residual error.
224 Alternative methods may therefore be considered e.g. a suitably selected covariate.

225
226 For induced infections, the use of helminth naive animals is recommended. Animals not
227 raised in a helminth-free environment should be treated with an approved anthelmintic,
228 chemically not related to the test drug, to remove pre-existing infections followed by faecal
229 examination to determine that the animals are helminth free.

230
231 Animal housing, feeding and care should follow strict requirements of welfare including
232 vaccination according to local practices. This information should be provided in the final
233 report. A minimum 7-day acclimatisation period is recommended. Housing and feed/water
234 should be adequate according to the geographical location. Animals should be monitored
235 daily to determine adverse reactions.

236 237 **B. Specific Evaluation Studies**

238 239 **1 - Dose Determination Studies**

240
241 No species specific recommendations.

242 243 **2 - Dose Confirmation Studies**

244
245 Confirmation studies are needed to support each claim: adult, larvae and when applicable
246 hypobiotic larvae.

247 248 **3 - Field Efficacy Studies**

249
250 The field studies should be replicated in different geographic locations and in animal/production
251 class(es) that represent the conditions of use for the indication being pursued. The protocol
252 should state the number of experimental units per treatment group (sample size), describe
253 allocation (proportion) to treatment groups, and include a brief description of how the sample
254 size was determined. The protocol should also describe procedures for random selection of
255 animals (number and percentage) to be sampled (if faecal samples will not be collected from
256

257 all available animals in the study), as appropriate, and the methods to be used for both faecal
258 collection and examination. Regardless of whether one or multiple parasites are being
259 evaluated within a study, an appropriate sample size calculation or justification is necessary
260 prior to study conduct.

261
262 Effectiveness against adult nematodes can be assessed by the reduction of faecal egg counts
263 and should be performed using samples from the same animal before and after treatment in
264 both study groups (control and treated). Post-treatment counts are generally made 10-14 days
265 after treatment, but the timing of post-treatment counts will depend on the parasite species and
266 class of anthelmintic evaluated. For example, due to the known effects of macrocyclic lactones
267 on nematode egg suppression, post-treatment counts should be delayed until at least 14 days
268 or longer. Efficacy should be calculated using post-treatment faecal egg counts from the treated
269 and control (typically placebo or untreated control) groups. Additionally, a calculation of efficacy
270 using pre- and post-treatment faecal egg counts may provide further information on field
271 effectiveness. Furthermore, additional endpoints for evaluating field effectiveness should be
272 considered as they are developed and generally accepted by experts in veterinary parasitology.

273
274 See also Sections 4.1 and 4.2 of VICH GL7.

275 276 **4 - Persistent Efficacy Studies**

277
278 Two basic study designs have been used to pursue persistent efficacy claims: one using a
279 single challenge, another using multiple daily challenges following treatment. For both
280 procedures, no standardised protocols have been developed. When conducting studies,
281 protocols details should include among other things: determination of larval viability
282 throughout the study, rationale for larval challenge and justification of slaughter-time. Parasite
283 naive cattle are recommended in these studies. A study design is recommended using
284 multiple daily challenges, as this most closely mimics what occurs in nature.

285
286 A minimum requirement for a persistent efficacy claim (for each duration and helminth claim)
287 should include 2 trials (with worm counts) each with a non-treated and one or more treated
288 groups. At least 6 animals in the control group shall be adequately infected. Persistent
289 efficacy claims will only be granted on a species-by-species basis.

290 In the protocol using multiple daily challenges, different groups of animals are treated and
291 exposed to a daily natural or induced challenge for 7, 14, 21 or more days after the treatment,
292 then at approximately 3 weeks after the last challenge (or earlier) the animals are examined
293 for parasite burden. The challenge interval and schedule may vary for longer acting
294 products, and should take into consideration the pharmacological properties of the product.

295
296 Persistent efficacy claims should be supported by a minimum 90% efficacy at each time
297 point and calculated and interpreted using the procedures described in Sections 4.1 and 4.2
298 of VICH GL7. Persistent efficacy claims should be granted for the longest period between
299 treatment and the last challenge where effectiveness criteria are met and all preceding time
300 points tested meet the criteria as well.