



1 24 June 2022  
2 EMA/CVMP/VICH/835/1999  
3 Committee for Veterinary Medicinal Products (CVMP)

4 **VICH GL19 Efficacy of anthelmintics: specific**  
5 **recommendations for canines (Revision 1)**  
6 **Draft**

Draft agreed by VICH Steering Committee	May 2022
Adoption by CVMP for release for consultation	15 June 2022
Start of public consultation	24 June 2022
End of consultation (deadline for comments)	1 November 2022

7  
8

Comments should be provided using this [template](#). The completed comments form should be sent to [vet-guidelines@ema.europa.eu](mailto:vet-guidelines@ema.europa.eu)

9





International Cooperation on Harmonisation of Technical Requirements  
for Registration of Veterinary Medicinal Products

**VICH GL19 (ANTHELMINTICS CANINES)**  
**May 2022**  
**Revision at Step 9**  
**For consultation at Step 4**

# **EFFICACY OF ANTHELMINTICS: SPECIFIC RECOMMENDATIONS FOR CANINES (REVISION 1)**

Revision at Step 9

Recommended for Consultation at Step 4 of the VICH Process  
in May 2022  
by the VICH Steering Committee

This Guideline has been developed by the appropriate VICH Expert Working Group will be subject to consultation by the parties, in accordance with the VICH Process. At Step 7 of the Process the final draft is recommended for adoption to the regulatory bodies of the European Union, Japan and USA.

Secretariat: c/o HealthforAnimals, 168 Av de Tervueren, B-1150 Brussels (Belgium) - Tel. +32 2 543 75 72  
e-mail : [sec@vichsec.org](mailto:sec@vichsec.org) - Website : <http://www.vichsec.org>

54  
55  
56  
57  
58

# EFFICACY OF ANTHELMINTICS: SPECIFIC RECOMMENDATIONS FOR CANINES

59

## INTRODUCTION

60  
61 The present guideline for canines was developed by the Working Group that was established  
62 by the Veterinary International Cooperation on Harmonization (VICH), Anthelmintic  
63 Guidelines. It should be read in conjunction with the VICH Efficacy of Anthelmintics: General  
64 Requirements (VICH GL7) which should be referred to for discussion of broad aspects for  
65 providing pivotal data to demonstrate product anthelmintic effectiveness. The present  
66 document is structured similarly to VICH GL7 with the aim of simplicity for readers comparing  
67 both documents.

68  
69 The aim of this guideline for canines is: (1) to be more detailed for certain specific issues for  
70 canines not discussed in VICH GL7; (2) to highlight differences with VICH GL7 on data  
71 requirements and (3) to give explanations for disparities with VICH GL7 guideline.

72  
73 It is important to note that technical procedures to be followed in the studies are not the aim  
74 of this guideline. We recommend that the sponsors refer to pertinent procedures described  
75 in detail in other published documents, e.g. WAAVP Guidelines for Evaluating the Efficacy  
76 of Anthelmintics for Dogs and Cats, Veterinary Parasitology **52**: 179-202, 1994, and updated  
77 versions as they are published.

78

## A. General Elements

79

### 1. The Evaluation of Effectiveness Data

80  
81  
82  
83 The evaluation of effectiveness data is based on parasite counts (adults, larvae) in dose  
84 determination and dose confirmation studies; egg counts/larval identification is the  
85 preferred method to evaluate effectiveness in field studies.

86  
87 The controlled test is the most widely accepted of the testing procedures for evaluation  
88 of anthelmintic drug effectiveness. However, the critical test may be appropriate for some  
89 intestinal species of parasites, e.g. ascarids.

90  
91 Adequate parasite infection should be defined in the protocol according to regional prevalence,  
92 historic data and/or statistical analysis.

93

### 2. Use of Natural or Induced Infections

94  
95  
96 Dose determination studies should be conducted using induced infections with either  
97 laboratory strains or recent field isolates.

98  
99 Dose confirmation studies should be conducted using naturally or artificially infected animals.  
100 Where possible, at least one study should be conducted in naturally infected animals; deviation  
101 from this requirement should be justified, e.g., applicable laws or regulations prohibit sourcing  
102 of naturally infected animals. Two studies should be conducted for each parasite claimed on  
103 the label. If both studies are conducted using experimentally infected animals, then parasites  
104 must have originated from naturally occurring infections from different geographical regions no  
105 older than 10 years prior to use for inducing infection. In addition to two dose confirmation  
106 studies, the efficacy and safety is generally confirmed by data from field studies. *Echinococcus*  
107 spp. and *Dirofilaria* spp. testing may be conducted using animals harbouring induced

108 infections due to public health considerations for echinococcosis and the complexity of the  
109 claims for heartworm. Due to the zoonotic potential of *Echinococcus* spp. trials conducted  
110 using this genus should be carried out under high biosecurity provisions.

111  
112 For the following helminths, induced infections may also be the only method to determine  
113 effectiveness of the product because of difficulties in obtaining a sufficient number of  
114 infected animals: *Filaroides milksi*, *F. hirthei*, *Diocotophyma renale*, *Capillaria aerophila*, *C. plica*,  
115 *Spirocerca lupi*, *Physaloptera* spp, *Mesocestoides* spp. and *Crenosoma vulpis*. For claims  
116 against larval stages, only studies with induced infections are acceptable.

117  
118 The history of the parasites used in the induced infection studies should be included in the  
119 final report.

### 120 121 **3. Number of Infective Parasitic Forms Recommended for Induced Infections**

122  
123 The number to be used is approximate and will depend on the isolate. The final number of  
124 larvae used in the infection should be included in the final report. Table 1 shows the range of  
125 numbers recommended for common helminths.

126  
127 **Table 1. Range of infective stages used to produce adequate infections in canines**  
128 **for anthelmintic evaluation**

129

<b>Parasite Anatomical Location</b>	<b>Range</b>
<b>Genus Species</b>	
<b>Small Intestine</b>	
<i>Toxocara canis</i>	100 – 500*
<i>Toxascaris leonina</i>	200 – 3,000
<i>Ancylostoma caninum</i>	100 – 300
<i>Ancylostoma braziliense</i>	100 – 300
<i>Uncinaria stenocephala</i>	1,000 – 1,500
<i>Strongyloides stercoralis</i>	1,000 – 5,000
<i>Echinococcus granulosus</i>	20,000 – 40,000
<i>Taenia</i> spp.	5 – 15
<b>Large Intestine</b>	
<i>Trichuris vulpis</i>	100 – 500
<b>Heart</b>	
<i>Dirofilaria immitis</i>	30 – 100 **

130

\* In suckling canines or canines less than 5 months of age.

131

\*\* For adulticidal or microfilaricidal testing 5 to 15 pairs of adult worms can be  
132 transplanted.

133

### 134 **4. Recommendations for the calculation of effectiveness**

135

#### 136 **4.1. Criteria to Grant a Claim**

137

138 To be granted a claim the following pivotal data should be included:

139

140 a) Two dose confirmation studies conducted with a minimum of 6 adequately infected non-  
141 medicated animals (control group) in each study. The infection of the animals in the study  
142 will be deemed adequate based on historical, parasitological and/or statistical criteria.

143

144 b) The differences in parasite counts between treated and control should be statistically  
145 significant ( $p \leq 0.05$ ).

146

147 Efficacy should be 90% or higher and calculated and interpreted using the procedure  
148 described in Section 4.2 of VICH GL7.

149  
150 For some parasites with public health, animal welfare/clinical implications, e.g. *E.*  
151 *granulosus* and *D. immitis*, respectively, higher efficacy standards (i.e. up to 100%) may  
152 be imposed. The regulatory authority of the region in which the product is intended to be  
153 registered should be consulted.

154  
155 c) Effectiveness against helminths will be evaluated examining for the presence or absence  
156 of parasitic elements in faecal material or blood. An *Echinococcus* spp. claim does not  
157 require field studies due to public health concerns.

#### 158 **4.2. Number of Animals (Dose Determination and Dose Confirmation Trials)**

159  
160 The minimum number of animals required per experimental group is a critical point. Although  
161 the number of animals will depend on the ability to process the data statistically according  
162 to the adequate statistical analysis it has been recommended, to achieve harmonization,  
163 that the inclusion of at least 6 animals in each experimental group is a minimum.

164  
165 In cases where there are several studies, none of which have 6 adequately infected animals  
166 in the control group (for example, important rare parasites), the results obtained could be  
167 pooled to accumulate 12 animals in the studies; and statistical significance calculated. If the  
168 differences are significant ( $p < 0.05$ ), effectiveness may be calculated and if the infection is  
169 deemed adequate, the claim may be granted. Sampling techniques and estimation of  
170 worm burden should be similar among laboratories involved in the studies to allow  
171 adequate and meaningful extrapolation of the results to the population.

#### 172 **4.3 Adequacy of Infection**

173  
174 The minimum adequate number of helminths in individual control animals should be  
175 defined in the protocol. However, final conclusions regarding adequacy of infection  
176 will be made as part of the final report based on statistical analysis, historical data, literature  
177 review, or expert testimony. Generally, a minimum of 5 nematodes in individual control  
178 animals is considered an adequate infection. -For *Dirofilaria immitis* microfilaria (mff) claims,  
179 300 mff/mL (blood) is considered an adequate infection. Recommended counts (in individual  
180 control animals) to be considered adequate for example cestodes include:

181  
182  
183  
184 *Echinococcus* spp. - 5 scolices  
185 *Taenia* spp. - 2 scolices  
186 *Dipylidium caninum* - 2 scolices

#### 187 **4.4 Label Claims**

188  
189 A claim for effectiveness against life stages of each parasite should refer to each stage in  
190 the case of natural infections, or age in days in the case of induced infection. Table 2 is  
191 provided as a guide for the recommended time of treatment of induced infections.

192  
193 With the majority of parasites approximately 7 days is a sufficient time period from the  
194 termination of treatment until the animals are necropsied. The following parasites are the  
195 exception to the above general recommendation:

- 196  
197  
198  
199 - *Physaloptera* spp., *S. lupi*, *C. plica*, *D. renale*, *E. granulosus*, *Taenia* spp., *D.*  
200 *caninum*, *Mesocestoides* spp.: 10 to 14 days;  
201 - *C. vulpis*: 14 days;  
202 - *F. milksi*, *F. hirthei*: 42 days;

- 203 - *F. osleri*: one-half of the animals at 14 days and the other half at 28 days;  
 204 - *D. immitis*: varies by trial design.

205  
206  
207 **Table 2. Recommended time of treatment after infection**  
208

Parasite	Adult Stages	Larval Stages
<i>S. stercoralis</i>	5 to 9 days	
<i>T. vulpis</i>	84 days	
<i>A. caninum</i>	> 21 days	
<i>A. braziliense</i>	> 21 days	6 to 8 days * (L4)
<i>U. stenocephala</i>	> 21 days	6 to 8 days (L4)
<i>T. canis</i>	49 days	6 to 8 days (L4)
		3 to 5 days (L3/L4)
<i>T. leonina</i>	70 days	14 to 21 days (L4/L5)
<i>D. immitis</i>	180 days	35 days (L4)
		2 days (L3), 20 to 40 days (L4)
<i>E. granulosu</i>	> 28 days	70 to 120 days (L5), 220 days (microfilariae)
<i>s Taenia spp.</i>	> 35 days	

209 \* For somatic larvae, treat within 2 days prior to  
 210 parturition.  
 211

212 For claims against transplacental and/or transmammary transmission of *T. canis* somatic  
 213 larvae of natural or artificially infected pregnant bitches should be treated prior to parturition  
 214 and the efficacy checked by counting the larvae in the bitch milk and/or the adult worms  
 215 in the small intestines of the litter.  
 216

217  
218 **5. Treatment Procedures**  
219

220 The method of administration (oral, parenteral, topical), formulation and extent of activity of  
 221 the product will influence the protocol design. It is advisable to consider the weather and  
 222 animal relationship and bathing with regard to effectiveness of topical formulations.  
 223

224 For oral formulations, palatability studies should always be included in the evaluation of  
 225 the effectiveness of the product. For products administered topically, the impact of weather  
 226 (e.g. rainfall, UV light), bathing and coat length should be included in the evaluation of the  
 227 effectiveness of the product.  
 228

229 **6. Animal Selection, Allocation and Handling**  
230

231 Approximately 6 month old canines are suitable for effectiveness studies. However there  
 232 are exceptions:

- 233 - *S. stercoralis*: less than 6 months;  
 234 - *A. caninum*, *A. braziliense*: 6 to 12 weeks;  
 235 - *T. canis*, *T. leonina*: 2 to 6 weeks;  
 236 - *D. caninum*: 3 months or older;  
 237 - *Mesocestoides* spp.: 8 weeks or  
 238 older;  
 239 - *T. vulpis*: dogs older than 6 months can be used.  
 240

241 Naturally infected animals are selected based on egg output or expelled proglottids for  
 242 gastrointestinal parasites, and parasitological and/or immunological methods for *D. immitis*.  
 243 Randomization to treatment group should be performed using an adequate method that should  
 244 be described in the protocol and final report. Blocking should only be employed if it is expected

245 to reduce residual error in the study. If blocking is used, blocks should be included as a random  
246 effect in the statistical model. Nevertheless, blocking is not always the most appropriate  
247 method for reducing residual error. Alternative methods may therefore be considered e.g. a  
248 suitably selected covariate. Animal housing, feeding and care should follow strict  
249 requirements of welfare for canines. Animals should be acclimated for at least 7 days to the  
250 experimental facilities and personnel. Animals should be monitored daily to determine adverse  
251 reactions.

252

## 253 **B. Specific Evaluation Studies**

254

### 255 **1. Dose Determination Studies**

256

257 No species-specific recommendation.

258

### 259 **2. Dose Confirmation Studies**

260

261 No species-specific recommendation.

262

### 263 **3. Field Efficacy Studies**

264

265 Field (clinical) studies should not be conducted with canines infected with *Echinococcus*  
266 spp.

267

### 268 **4. Persistent Efficacy**

269

270 Due to the differing biologies for the helminths of canines and the lack of experience  
271 with persistent efficacy for these parasites, no recommendations can be provided.